

**Table 5.7. Timing of Summer Steelhead Life Cycle Events for the Rogue Basin**

<b>Life Cycle</b>	<b>Oct</b>	<b>Nov</b>	<b>Dec</b>	<b>Jan</b>	<b>Feb</b>	<b>Mar</b>	<b>Apr</b>	<b>Ma y</b>	<b>June</b>	<b>July</b>	<b>Aug</b>	<b>Sept</b>
<b>Early-run adult migration</b>								X	X	X		
<b>Half- pounder migration</b>											X	X
<b>Late-run adult migration</b>	X										X	X
<b>Adult spawning</b>			X	X	X	X						
<b>Eggs/fry emerge</b>						X	X	X	X*			
<b>Fingerlings/ rearing</b>	X*	X	X	X	X	X	X	X	X*	X*	X*	X*
<b>Juvenile migration</b>		X	X	X	X	X	X	X	X*	X*		
<b>Smolt out- migration</b>							X	X	X			

X Indicates presence at the life cycle month stage.

X\* Indicates severe limiting factor condition in the life cycle month stage.

(Determination by technical committee.

Summer steelhead fry emerge from the gravel in April and May, then migrate to the mainstem in May and June as the natal streams dry up. Smolts generally migrate downstream from March through June.

This document was created with Win2PDF available at <http://www.daneprairie.com>.  
The unregistered version of Win2PDF is for evaluation or non-commercial use only.