

This report was funded by the Bonneville Power Administration (BPA), U.S. Department of Energy, as part of BPA's program to protect, mitigate, and enhance fish and wildlife affected by the development and operation of hydroelectric facilities on the Columbia River and its tributaries. The views in this report are the author's and do not necessarily represent the views of BPA.

For additional copies of this report, write to:

Bonneville Power Administration
Public Information Center - CKPS-1
P.O. Box 3621
Portland, OR 97208

Please include title, author, and DOE/BP number from the back cover in the request.

**UMATILLA BASIN NATURAL PRODUCTION
MONITORING AND EVALUATION**

ANNUAL PROGRESS REPORT 1994-1995

Prepared by:

Craig R. Contor
Eric **Hoverson**
Paul Kissner
Jed Volkman

Fisheries Program
Department of Natural Resources
Confederated Tribes of the
Umatilla Indian Reservation
Pendleton, OR

Contract Period
September 30, 1994 to September 29, 1995

Prepared for:

U.S. Department of Energy
Bonneville Power Administration
Environment, Fish and Wildlife
P. O. Box 3621
Portland, OR 97208-3621

Project Number 90-005-01
Contract Number **DE-BI79-92BP75349**

April 1996

ABSTRACT

This report summarizes the activities of the Umatilla Basin Natural Production Monitoring and Evaluation Project (UBNPME) from September 30, 1994 to September 29, 1995. This program was funded by Bonneville Power Administration and was managed under the Fisheries Program, Department of Natural Resources, Confederated Tribes of the Umatilla Indian Reservation.

An estimated 36.7 km (22.6 miles) of stream habitat were inventoried on the Umatilla River, Moonshine, Mission, Cottonwood and Coonskin Creeks. A total of 384 of 3,652 (10.5%) habitat units were electrofished. The number of juvenile fish captured follows: 2,953 natural summer steelhead (including resident rainbow trout; *Oncorhynchus mykiss*), one hatchery steelhead, 341 natural chinook salmon (*O. tshawytscha*), 163 natural coho salmon (*O. kisutch*), five bull trout (*Salvelinus confluentus*), 185 mountain whitefish (*Prosopium williamsoni*), and six northern squawfish (*Ptychocheilus oregonensis*). The expanded population estimate for the areas surveyed was 73,716 salmonids with a mean density of 0.38 fish/m².

The following number of non-salmonids were visually estimated: 7,572 speckled dace (*Rhinichthys osculus*), 5,196 sculpin (*Cottus spp.*), 532 suckers (*Catostomus spp.*) and 191 redbreast shiners (*Richardsonius balteatus*). The gross estimated density of all non-salmonids combined was 0.84 fish/m². The estimated ratio of non-salmonids to salmonids was 2.4: 1.

Relative salmonid abundance, seasonal distribution and habitat utilization were monitored at index sites throughout the basin. During index site monitoring, the following species were collected in addition to those listed above: american shad (*Alosa sapidissima*), smallmouth bass (*Micropterus dolomieu*), carp (*Cyprinus carpio*) and chiselmouth (*Acrocheilus alutaceus*). Thirty-nine sites were electrofished during the spring and summer seasons, while 36 sites were sampled in the fall season. Index sites with the highest mean salmonid catch/minute (fish/min.) during the three sample periods were located at the following sites: East Birch Creek (3.4 fish/min.), Boston Canyon Creek (3.2 fish/min.), Spring Creek (3.1 fish/min.) and upper Squaw Creek (3.0 fish/min.). The highest electrofishing catch rates were observed in the Umatilla River tributaries above river mile (RM) 70 in the August and September sample period (Table J-2 catalogs river miles with associated landmarks). During the November sample period, catch rates were highest in Birch Creek tributaries. Most salmonids were captured in slow water near the bank during the November and March sampling periods.

A study of the migration movements and homing requirements of adult salmonids in the Umatilla River was conducted during the 1994-95 return years. Radio telemetry was used to evaluate the movements of adult salmonids past diversion dams in the lower Umatilla River and to determine migrational movements of salmonids following upstream transport. Radio transmitters were placed in 30 summer steelhead, 15 spring chinook, nine fall chinook, and eight coho salmon. Salmon were released at Three Mile Falls Dam (TMD). An additional 11 summer steelhead and ten spring chinook salmon were tagged, hauled upstream, and released at either Barnhart, Nolin, Thornhollow, or Imeques C-mem-ini-kern. On average, summer steelhead required 36 days to successfully migrate from TMD to Stanfield Dam. Spring chinook required 18 days. Average passage times for summer steelhead (hours and minutes) at Westland, Feed Canal, and Stanfield Dams were 13:06, 83:24, and 2:58, respectively. Spring chinook salmon required 04:30 at Westland, 89:42 at Feed Canal, and 04:01 at Stanfield Dams. Migrational delays were observed at Feed Canal Dam at flows ranging from 563 to 1,601 cubic feet/second (cfs). Thirty-eight

percent of the fish used the fish ladder at **Westland** Dam, 75% at Feed Canal Dam, and 31% at Stanfield Dam. Average passage times at Feed Canal Dam (1995) were more than 15 times those at Stanfield Dam in 1994 and more than 20 times those-at Stanfield Dam in 1995.

Data related to homing and passage needs of Umatilla River salmonids was investigated in an attempt to maximize homing to the Umatilla River. Straying rates of adult summer steelhead and spring chinook salmon were found to be low while **coho** and fall chinook salmon stray rates were high in some groups, particularly adult returns from subyearling smolt releases of fall chinook salmon.

Attraction flows of from the mouth of the Umatilla River of at least 150 cfs were required to encourage migration and reduce straying of fall chinook and **coho** salmon. Significant numbers of summer steelhead entered when flows exceeded 500 cfs. Spring chinook salmon entry was variable with fish entering at flows ranging from 150 to more than 2,000 cfs.

Adult anadromous salmonids potentially available to spawn above TMD from August 26, 1994 to June 27, 1995 included: 593 adult and 530 jack fall chinook salmon (1994 brood), 879 adult and 54 jack **coho** salmon (1994 brood), 784 natural and 509 hatchery summer steelhead (1995 brood), and 378 adult and 62 jack spring chinook salmon (1995 brood). During escapement surveys (fall of **1994**), a total of 82 fall chinook salmon redds, 24 **coho** salmon redds and seven unidentified salmon redds (112 redds total, **2.6/mile**) were enumerated along 42.3 miles of the **mainstem** above TMD. In 1995, we enumerated and flagged 126 summer steelhead redds (3.6 redds/mile) along 35.3 miles of lateral tributaries of the Umatilla River. Also enumerated were 90 spring chinook salmon redds (1.6 redds/mile) along 55.8 miles of the mainstem. Ninety-six percent of the adult fall chinook salmon carcasses examined had spawned while 94% of the **coho** had spawned; 66.8% of the spring chinook salmon carcasses examined had spawned. A total of 49.3% of spring chinook salmon released above TMD were sampled during spawning ground surveys and 60 coded wire tags (**CWTs**) were recovered from 78 adipose clipped fish.

The rotary screw trap in the Umatilla River (**RM 76**) operated 63 of 113 days from September 21, 1994 to January 13, 1995. The trap captured 596 juvenile steelhead with a mean trap efficiency rate of 9.9%. A total of 1,368 juvenile chinook salmon were captured with a mean trap efficiency rate of 28.8%.

The rotary screw trap at the Imeques C-mem-ini-kern site (**RM 79.5**) operated 43 out of 43 days from May 5 through June 16, 1995. The trap captured 304 natural juvenile steelhead with a mean trap efficiency rate of 6.6%. A total of 102 natural juvenile chinook salmon were captured with a mean trap efficiency rate of 10.5%.

The rotary screw trap at the **Barnhart** site (**RM 42.2**) operated 87 out of 125 days from March 3 to June 1, 1995. The trap captured 105 natural juvenile steelhead, 247 natural juvenile chinook salmon, five natural **coho** salmon, 6,265 hatchery juvenile chinook salmon, 467 hatchery steelhead and 16,844 hatchery **coho** salmon. Mean trap efficiency rates ranged from 2.3 to 5.7%

Harvest monitors estimated that tribal anglers harvested 25 hatchery and five natural summer steelhead during the spring of 1995. There was no spring chinook salmon fishery in the Umatilla River during 1995 because of the low number of returning adults.

Scale analysis determined that over 85.0% of naturally produced juvenile summer steelhead sampled during biological and index surveys were age 0+ or 1 + . Naturally produced summer steelhead adults, returning to the Umatilla River in 1994-95, were mostly **from** the 1990 (46.4%) and 1991 (33.9%) brood years.

TABLE OF CONTENTS

| | |
|--|----|
| ABSTRACT | ii |
| INTRODUCTION | 1 |
| DESCRIPTION OF PROJECT AREA | 3 |
| MATERIALS AND METHODS | 4 |
| OBJECTIVE 1: Habitat Surveys | 4 |
| Task1.1: | 4 |
| Task1.2: | 5 |
| Tasks 1.3 through 1.5: | 6 |
| Task1.6: | 6 |
| OBJECTIVE 2: Biological Surveys | 6 |
| Task2.1: | 6 |
| Task2.2: | 7 |
| Task2.3: | 8 |
| Task2.4: | 8 |
| Task2.5: | 8 |
| Task2.6: | 9 |
| OBJECTIVES 3 and 4: Adult Passage Evaluations | 9 |
| Tasks3.1and4.1: | 9 |
| OBJECTIVE 5: Homing and Straying of Adult Salmonids | 10 |
| Task5.1: | 10 |
| OBJECTIVE 6: Spawning Surveys | 10 |
| Task 6.1: | 10 |
| Tasks6.2and6.3: | 11 |
| Task6.4: | 12 |
| Task6.5: | 12 |
| OBJECTIVE 7: Smolt Trapping | 12 |
| Task 7.1: | 12 |
| Task7.2: | 12 |
| Task7.3: | 12 |
| Task7.4: | 13 |
| Task7.5: | 13 |
| Task7.6: | 13 |
| OBJECTIVE 8: Tribal Harvest | 13 |
| Tasks8.1and8.2: | 13 |
| OBJECTIVE 9: Age and Growth | 14 |
| Tasks 9.1 and 9.2: | 14 |
| OBJECTIVE 10: Genetic and Ecological Effects of Supplementation | 14 |
| Task10.1: | 14 |
| Task10.2: | 14 |
| Task10.3: | 15 |
| Tasks 10.4 and 10.5: | 15 |

| | |
|--|-----------|
| OBJECTIVE 11: Supplementation Effects on Natural Steelhead | 15 |
| Task 11.1: | 15 |
| Task11.2: | 15 |
| RESULTS AND DISCUSSION | 16 |
| OBJECTIVE 1: Habitat Surveys | 16 |
| Task 1.1: | 16 |
| Task1.2: | 20 |
| Tasks 1.3 through 1.5: | 21 |
| Task1.6: | 21 |
| OBJECTIVE 2: Biological Surveys | 22 |
| Task2.1: | 22 |
| Task2.2: | 22 |
| Task2.3: | 25 |
| Task2.4: | 26 |
| Task2.5: | 27 |
| Task2.6: | 27 |
| OBJECTIVE 3: Adult Passage Evaluations | 27 |
| Task3.1: | 27 |
| OBJECTIVE 4: Adult Passage Evaluations Following Upstream Transport | 30 |
| Task4.1: | 30 |
| OBJECTIVE 5: Evaluate Homing and Straying of Adult Sahnionids | 31 |
| Task5.1: | 31 |
| OBJECTIVE 6: Spawning Surveys | 34 |
| Task6.1: | 34 |
| Tasks6.2and6.3: | 34 |
| Task6.4: | 36 |
| Task6.5: | 37 |
| OBJECTIVE 7: Smolt Trapping | 37 |
| Task 7.1: | 37 |
| Task7.2: | 38 |
| Task7.3: | 38 |
| Task7.4: | 38 |
| Task7.5: | 38 |
| Task7.6: | 38 |
| OBJECTIVE 8: Tribal Harvest | 38 |
| Tasks 8.1 and 8.2: | 38 |
| OBJECTIVE 9: Age and Growth | 39 |
| Tasks9.1 and9.2: | 39 |
| OBJECTIVE 10: Genetic and Ecological Effects of Supplementation | 40 |
| Task10.1: | 40 |
| Task10.2: | 40 |
| Task10.3: | 41 |
| Tasks 10.4 and 10.5: | 42 |
| OBJECTIVE 11: Supplementation Effects on Natural Steelhead | 42 |
| Task11.1: | 42 |
| Task11.2: | 43 |

| | |
|--|-----|
| ACKNOWLEDGEMENTS | 45 |
| LITERATURE CITED | 46 |
| APPENDICES: MAPS, TABLES AND FIGURES | A-1 |
| APPENDIX A: Maps | A-1 |
| APPENDIX B: Steelhead Returns and Flow | B-1 |
| APPENDIX C: Thermograph Locations and Temperatures | C-1 |
| APPENDIX D: Physical Habitat Surveys | D-1 |
| APPENDIX E: Biological Surveys | E-1 |
| APPENDIX F: Adult Passage Evaluations | F-1 |
| APPENDIX G: Spawning Survey Data | G-1 |
| APPENDIX H: Trapping | H-1 |
| APPENDIX I: Age and Growth | I-1 |
| APPENDIX J: Landmarks, River Miles, and Abbreviations | K-1 |

INTRODUCTION

The Umatilla Basin Natural Production Monitoring and Evaluation Project (UBNPME) was funded by Bonneville Power Administration (**BPA**) as directed by section 4(h) of the Pacific Northwest Electric Power Planning and Conservation Act of 1980 (**P.L.** 96-501) and pursuant of measure 703 (F)(1)(b) of **the** Northwest Power Planning Council's (NPPC) Columbia River Basin Fish and Wildlife Program (NPPC 1987). This report summarizes work completed during the contract year September 30, 1994 through September 29, 1995. Work was conducted by the Fisheries Program, Department of Natural Resources, Confederated Tribes of the Umatilla Indian Reservation (CTUIR) in cooperation with the Oregon Department of Fish and Game (ODFW, see Appendix J, Table J-2 for abbreviation definitions). This project was one of several subprojects of the Umatilla River Basin Fisheries Restoration Master Plan (CTUIR 1984, ODFW 1986) orchestrated to rehabilitate salmon and steelhead runs; subprojects include:

- Natural Production Monitoring and Evaluation, and Adult Passage Facility Evaluations (this project);
- Watershed Enhancement and Rehabilitation;
- Hatchery Construction and Operation;
- Satellite Facility Construction and Operations for Juvenile Acclimation and Release and Adult Holding and Spawning;
- Trapping and Hauling of Juvenile and Adult Salmonids Around Dry Reaches Below Irrigation Diversions;
- Juvenile Passage Facility Construction and Operation;
- Juvenile Passage Facility Evaluations;
- Evaluation of Juvenile **Salmonid** Outmigration and Survival in the Lower Umatilla River Basin;
- Adult Passage Facility Construction and Operation, and
- Flow Augmentation to Increase **Instream** Flows Below Irrigation Diversions.

The Umatilla River Basin Fisheries Restoration Master Plan identified the following four critical uncertainties that the UBNPME project addressed:

- 1) What was the observed natural production success and estimated natural production potential for spring chinook, fall chinook and **coho** salmon, and summer steelhead in **the** Umatilla River Basin?
- 2) How effective were the adult passage facilities?
- 3) was supplementation enhancing natural summer steelhead populations?
- 4) was supplementation impacting the genetic diversity and life history characteristics of native salmonids?

The approach to monitoring and evaluating the natural production in the Umatilla River Basin includes three phases. Phase one includes collecting baseline data relating to life histories, distribution, abundance, survival and **the** current and potential production of anadromous salmonids from the Umatilla Basin. Phase two involves the creation of a streamlined monitoring program developed and tested through completion of tasks in phases one and two. Phase three consists of risk containment monitoring where the monitoring program will be employed. Phase one of the UBNPME plan was scheduled for 1992-97. Phases two and three are scheduled to begin in 1997 and 2004 respectively.

The UBNPME program's 1994-95 goals were to evaluate the implementation of the Umatilla River Basin Fisheries Restoration Plan with respect to natural production, adult passage and tribal harvest. This report follows the outline of the task list from the statement of work as required postliminarily. Project objectives are listed below.

Objective 1: Estimate the amount of existing and potential spawning and rearing habitat for summer steelhead, spring and fall chinook and coho salmon.

Objective 2: Determine distribution, species composition and densities of fish species throughout the Umatilla Basin.

Objective 3: Utilize radio telemetry to evaluate the passage of adult salmonids past the major irrigation diversion dams and associated passage facilities on the lower Umatilla River.

Objective 4: Utilize radio telemetry to evaluate the movements of adult spring chinook salmon and summer steelhead trapped at Three Mile Falls Dam and transported upstream.

Objective 5: Evaluate factors that influence homing and straying of returning adult salmonids into or out of the Umatilla River Basin.

Objective 6: Determine natural spawning success, spawning habitat utilization, prespawning mortality, and number of redds/adult spring chinook salmon passed above Three Mile Falls Dam. Determine, if possible, spawning distribution and timing of steelhead, fall chinook salmon and coho salmon.

Objective 7: Estimate natural smolt production and survival rates of anadromous salmonids at various life history stages.

Objective 8: Estimate tribal harvest of returning adult salmon and steelhead.

Objective 9: Determine salmonid age, growth and life history characteristics.

Objective 10: Determine the genetic and ecological effects of supplementation on native steelhead and resident trout in the Umatilla Basin (as planned, this objective was not directly addressed during the 1994-95 contract year).

Objective 11: Determine if hatchery supplementation enhances production of natural steelhead (as planned, this objective was not directly addressed during the 1994-95 contract year).

DESCRIPTION OF PROJECT AREA

Summer steelhead, chinook and **coho** salmon-were abundant in the Umatilla River prior to **the** 1900's. Irrigation and agricultural development throughout the basin in the early 1900's was believed to be **the** primary cause of the decline of steelhead and the extinction of salmon (Bureau of Reclamation 1988). Since 1855, aquatic and riparian habitats have been degraded through irrigation diversions, water extractions, channelization, livestock grazing, logging, agriculture and urban development (Nielson 1950, NPPC 1987).

The Umatilla River Basin in northeast Oregon comprised **1,465,600** acres of the **6,400,000** acres of ceded CTUIR land (Figure A-1, A-2). The Umatilla River originated on the west slope of the Blue Mountains, east of Pendleton, and flows 115 miles in a northwesterly direction to the Columbia River at RM 289. The Umatilla River Basin, hydrologic unit number 17070103 (USGS **1989**), had a drainage area of 2,290 square miles. The mouth of the Umatilla River at Umatilla, Oregon, was at approximately 270 feet elevation (above mean sea level). The headwaters were as high as 4,950 feet. Mean annual precipitation ranged from ten inches/year at Umatilla to 50 inches/year in the headwaters (Taylor 1993).

The basin can be roughly divided into two physiographic regions. The lower river, west of Pendleton, has cut a low valley into a broad upland plain called the Deschutes-Umatilla Plateau. Parent geologic materials of the plain were dominated by multiple layers of middle Miocene basalt flows, specifically, the Wanapum and Grand Ronde **Basalts**, originating 14 to 17 million years ago. Basalt bedrock outcroppings were common in the river channel and act as hydraulic controls that delay the deepening of the river channel and valley floor. On top of the Miocene basalts were Pleistocene and Holocene **loess**, alluvial and glaciofluvial deposits (NPPC 1990, Walker and MacLeod 1991). Currently, vegetation on **the** broad Deschutes-Umatilla Plateau includes **dryland** crops and sagebrush-grass communities. Historically, deciduous trees were abundant in riparian areas on the valley floor; however, land-use practices over **the** last hundred years have cleared most of these areas for irrigated agricultural and urban uses. Approximately 70 percent of riparian areas in the Umatilla River Basin were reported to be in need of improvement (ODFW 1987).

The region east of Pendleton was dominated by foot hills and the Blue Mountains. The Blue Mountains were created by lifting, faulting and folding of volcanic, sedimentary and metamorphic rock. The middle Miocene basalts of the lower river were also the dominant parent materials in the headwaters. The river and streams have cut steep sided canyons into the layers of rock that form the higher elevations of the Blue Mountains. Exposed basalt fractured into blocks and plates while unexposed layers remain fairly impervious to water (Walker and **MacLeod** 1991). The combination of steep canyon walls and impervious bedrock lends to poor ground water recharge (NPPC 1990). U.S. Geological Survey (USGS) flow data from 1904 through 1994 show stream hydrographs that reflect the various features of the basin as described above. High flows regularly occur during rain storms and snow melt conditions. Extreme low flows were common during summer and dry conditions. This effect was less pronounced in the near pristine **North** Fork Umatilla Wilderness Area, apparently because of **the** lack of human disturbance, higher elevation of the headwaters, developed soils, large woody debris and climax plant communities. Vegetation distribution patterns upstream from Pendleton were typical for **the** Blue Mountains. Grasses and small shrubs dominated the drier, south facing slopes. Conifers dominated the north facing slopes, higher elevations and moderately wet areas.

MATERIALS AND METHODS

OBJECTIVE 1: Habitat Surveys

Task 1.1: Habitat Surveys.

Methods developed by **ODFW** (Moore et al. 1993) were used to inventory stream habitat. Habitat surveys were conducted from June 20 to September 11, 1995 on the Umatilla River (**RM** 81.8 to **89**), Moonshine Creek, Mission Creek, Cottonwood Creek and Coonskin Creek. A crew of two people worked upstream, dividing the valley into large scale reaches and the stream into individual habitat units. The same crew surveyed the entire stream to keep data as consistent as possible.

Reach classifications were made when major changes occurred in valley form, riparian composition or land use. A reach change could also be classified at fish passage barriers or when tributaries contributed a significant portion of flow to the stream being surveyed. **At the beginning** of a reach, we recorded specifics about land-form, valley-form, terrestrial vegetation, land use, water temperature, flow (high, medium or low) and valley floor width (**VWI**). **VWI** was the ratio of active channel width to valley floor width. Photographs were taken of **the** riparian areas and the reach. Notes and additional photographs were taken throughout the survey to document landmarks, habitat problems, passage concerns, irrigation diversions and surface springs. The locations of landmarks such as bridges or tributaries were marked **with** a unit number on a photocopy of a 7.5 minute quadrangle topographic map. A record was kept with detailed information on each photograph. An Oregon Water Resources map of the Umatilla River Basin was used to approximate river miles.

Stream habitat units were classified with more detail than were the reaches. A habitat unit was a section of stream that had a distinct hydraulic characteristics from adjacent stream sections (exception: dry channel classification). Each unit was numbered sequentially then identified as a riffle with pockets, lateral scour pool or glide, etc. Surveyors overestimated the **width** of dry channel units which inflated area calculations of dry units. Normally **the** width of a habitat unit was the wetted channel width which was narrower than active channel width (wet during bank full flows). When dry units were measured, **the** entire active channel width was measured as there was no water/shore interface.

If a unit was overlooked by a habitat crew but identified by **electrofishers**, the area was measured and recorded as an unclassified unit. Side channels with springs contributing **the** majority of the water were classified as spring seeps. Water temperatures were recorded from springs and tributaries and from the **mainstem** up and downstream. Crews estimated the percentage of **mainstem** flow contributed by each spring and tributary.

The following data were recorded at each habitat unit: estimated mean length, width, depth (maximum for slow water units and mean for fast water units), slope, aspect, shade, substrate composition, boulder count (**>** 0.5 m in diameter), wood rating (based on benefit to fish), bank stability, bank composition, percent undercut bank, percent flow in channel(s) and channel type. The primary channel measurements were kept separate from secondary channels measurements. The percent composition of gravel substrate was multiplied by the total wetted area surveyed to estimate potential spawning habitat.

At every tenth unit the following data were also recorded: unit length and width, active channel height and width, **VWI** and terrace characteristics. The starting point of every tenth unit was marked with an orange flag by **the** habitat survey-crew to enhance locating selected units during electrofishing. The number, habitat type and length of the unit was written on the flag.

Riparian communities were inventoried and photographed every 30 habitat units and at **the** start of each reach. A measuring tape was extended 30 m into the riparian zone, perpendicular to the stream, halfway between the upper and lower unit boundaries, and from **the** margin of **the** wetted and active channel. Three lateral transects measuring ten m long by five m wide were inventoried on both sides of the stream. Within each transect, the following data were recorded: geomorphic surface features, ground slope; canopy closure; percent shrub cover; percent grass; tree groups (conifer or hardwood); tree count by breast height diameter (**DBH**) class, and pertinent notes. Grain fields and stubble were tallied as grasses. The percentages of exposed soil, rock, roads, secondary stream channels were noted.

Woody debris were tallied and described if they met minimum length (3 m) and diameter (15 cm) requirements. Root wads were tallied if they met **the** minimum diameter requirement (15 cm). Crews recorded tree group (conifer or hardwood), length class, diameter, configuration and location in the channel for woody debris.

Task 1.2: Monitor stream temperatures in the Umatilla Basin, and examine USGS flow data from active gages in the basin.

Temperatures

CTUIR, ODFW, U.S. Forest Service (**USFS**) and U.S. Bureau of Reclamation (**BOR**) coordinated **the** deployment of 32 thermographs and four HYDROMET stations in the Umatilla River Basin to maximize consistency and coverage without duplicating effort. Specifics regarding **the** location and deployment of these thermographs were summarized in Tables C-1 through C-5. CTUIR thermographs were initialized, downloaded and deployed in the field with the use of a portable computer. New batteries were installed and the seals and clamps were cleaned, inspected and changed as needed. Thermographs were sealed inside a waterproof housing and placed inside a small cage made of expanded steel. Steel chains or cables anchored the units to a large tree or boulder on the shore. Thermographs and cables were concealed to minimize tampering. Photographs were taken and detailed descriptions of the location of each thermograph were written at **the** time of deployment. Detailed vicinity maps were drawn and 7.5 minute topographic maps were marked.

Flow

We examined the correlation between flow and the number of adult natural summer steelhead returning to the Umatilla River (two years later) for 16 years of flow and return records (Hubbard et al. 1995, Suzanne Miller, USGS, personal communication). Adult steelhead returns prior to 1982-83 were not correlated to flows because counts were considered to be rough estimates (Jim Phelps, ODFW, personal communication). The number of returning adult natural steelhead was compared to mean annual and monthly flows at the Umatilla gage (**RM 1.2**). The flow year and steelhead return years were designated differently by convention and can be confusing. For example, **the** comparison between flows in Water Year 1990 (October 1989 to September 1990) and steelhead returns in 1992-93 (fall 1992 through spring 1993) was denoted as a two year lag. However, the actual number of months between spring flows during juvenile emigration and when the adult steelhead actually return to the river may range from 30 to 35

months. Correlation coefficients were calculated by using Pearson's product-moment correlation with Bonferroni adjustments on multiple tests (SYSTAT 1984).

Tasks 1.3 through 1.5: Obtain habitat data collected by other agencies. Digitize and summarize habitat data. Estimate total usable habitat by stream reach, drainage and entire basin.

Data from Habitat surveys conducted by ODFW were obtained on computer diskette. No additional data entry or summarization was required. Raw habitat data collected and recorded in the field by CTUIR was entered into a database program. Original data were copied and archived. Data were validated before and after entry. After the second validation, summary charts and tables were created and examined for a final validation.

Estimates of total usable habitat by stream reach, drainage and basin were calculated from surveys conducted during summer low flow periods (1993-95). Usable habitat was defined as the area of a stream surveyed **that** had adequate water with suitable temperatures ($<24^{\circ}\text{C}$ Brett 1952, Black 1953). Expansions were made for reaches not surveyed by using data from adjacent streams of similar type. Wildhorse Creek, Butter Creek and several ephemeral streams were estimated to provide no anadromous **salmonid** habitat even though we have observed a few **salmonids** near spring seeps (Table B-1).

Task 1.6: Coordinate water quality monitoring efforts in the Lower Umatilla River with the Oregon Department of Environmental Quality.

Total maximum daily load (TMDL), water temperature monitoring, suspended sediment monitoring and water quality monitoring efforts in the basin were coordinated among Department of Environmental Quality (**DEQ**), ODFW, BOR, USFS, and CTUIR. Coordination was facilitated by **the** Umatilla Monitoring Evaluation and Oversight Committee (UMEOC) and **the** Umatilla Total Maximum Daily Load Technical Advisory Committee.

OBJECTIVE 2: Biological Surveys

Task 2.1: Conduct **salmonid presence/absence surveys in the Umatilla River Basin.**

Emphasis in conducting **salmonid** presence/absence surveys was minimized to allow completion of index site and quantitative biological surveys. Presence/absence surveys were conducted as time allowed to determine **salmonid** distribution. Several presence/absence sites were sampled in tributaries of the North Fork Umatilla River.

One electrofishing pass was made intermittently through several hundred meters of stream. Crews concentrated on areas where the probability of capturing salmonids was highest. The distance sampled was variable and could include multiple areas of a stream. Surveyors took photographs, marked the site on a map, recorded species and lengths of the catch, recorded site conditions and dimensions, and recorded effort (seconds of electrofishing).

Task 2.2: Electrofish and estimate salmonid densities in streams surveyed for habitat.

Backpack electroshockers and blocknets were used to sample fish from streams recently inventoried for habitat. Crews began electrofishing within several weeks of habitat surveys to best record relationships between habitat conditions and **salmonid** abundance. The units sampled for fish were selected in the field by **the** biological survey crew leader. Field selection was necessary because some units could not be sampled due to excessive depth, width, **instream** cover or absence of water. Every effort was made to minimize selective bias by stratifying the samples throughout the reach and by sampling approximately ten percent of the wetted area. Units with a variety of physical characteristics (i.e. braided and single channels, shaded or unshaded, cover or lack of cover) were sampled to represent the stream's habitat complexity. Care was taken to avoid startling fish from a unit before securing block nets. Water temperatures were recorded in all units sampled.

Salmonids were captured with dip-nets and removed on successive electrofishing passes until a depletion rate of at least 50% was achieved. The same individual electrofished in a similar manner for the same number of seconds (or slightly more) as the previous pass. This maximized equality of sampling effort between removal passes. Electroshocker settings (i.e. volts, pulse) remained constant for each pass. A second pass was not done if salmonids were neither captured nor observed during the first pass.

Captured salmonids were placed in a **livewell** until the completion of each pass. Fish were identified to species, measured (fork length, mm) and inspected for fin clips. Indicators of fish condition such as injuries, signs of disease or stress were noted. Bird bites were delineated as either puncture or scissor wounds.

Juvenile spring chinook salmon were not differentiated from juvenile fall chinook salmon nor were juvenile steelhead differentiated from resident rainbow trout. After examination, salmonids were released where captured or into a nearby area if conditions were significantly better.

Scale samples were taken from a portion of **the** total salmonids captured. A wide variety of sizes were sampled for age determination. Approximately 6-12 scales were removed from an area above the lateral line, posterior to the dorsal fin, and anterior to the adipose fin. Scale samples were taken from all **salmonid** mortalities. Scales were placed in clear mylar envelopes labeled with stream name, unit number, date, species and length.

Captured northern squawfish were sacrificed. Stomach contents were examined to determine the extent of predation on juvenile salmonids. Scale samples were taken from each **squawfish** and placed in mylar envelopes. Numeric estimates of all other non-salmonids observed during the first pass were recorded.

Estimates of **salmonid** abundance were calculated with a maximum-likelihood model (Van Deventer and **Platts** 1989) from the number of salmonids captured during successive electrofishing removal passes. Densities were estimated by dividing estimated **salmonid** abundance with estimated wetted channel area (estimated from habitat data). Low sample sizes required us to pool ***Oncorhynchus*** species to generate **salmonid** abundance estimates. Estimates for each species were calculated by multiplying the percent species composition by the expanded estimate for all salmonids. Mean density for a specific habitat type was calculated by dividing the sum of population estimates for each unit type by the area electrofished. The population estimates for each habitat type were added together to estimate the total population of the stream. **Salmonid** densities were also estimated for slow and fast water units. Densities for whitefish and **squawfish** were estimated only for habitat types where they were captured. Densities were also calculated from actual catch rather than from expanded abundance estimates. Densities of other **non-**

salmonids were based on the number observed (not captured) divided by area. Expanded estimates of non-salmonid abundance were calculated by multiplying the total wetted habitat area by the estimated density.

Task 2.3: Electrofish permanent index sites during November, April and August.

We **electrofished** 40 permanent index sites located throughout the Umatilla River Basin to monitor **salmonid** relative abundance, seasonal distribution and habitat utilization. (Figure A-3). Stable sites were chosen with the intent to monitor changes in **salmonid** populations rather than salmonid's response to changes in habitat. Habitat at each site was evaluated using the same methodology as in our habitat surveys (Task 1.1).

A typical index site consisted of fast and slow water habitat type. A few sites had more than two habitat types. Meacham Creek (site 30) was the only site with only one habitat type.

The lower and upper boundary of each site was marked in the field with numbered tags to assist consistent sampling. Most tags were placed on living trees or on wooden posts outside of the active channel to avoid tag loss during high flows. Site measurements, photographs and a detailed description of tag and site location were taken to expedite locating **the** site. Each index site location was also marked on an Oregon Water Resources map of the Umatilla River Basin (Figure A-3).

Index sites were sampled during March, August and November. Specific time periods for sampling varied depending on environmental conditions. Floods, cold weather, de-watering and inaccessibility occasionally prevented the sampling of some sites. During each sampling period, the length, width and depth of each habitat unit was measured at each index site. We measured mean depth in fast water units and maximum depth in slow water units. The habitat was measured to monitor physical changes which may effect catchability, abundance and species composition. Crews took photographs and recorded water and air temperatures, **weather**, stream flow (low, medium or high), water clarity, visibility, and electrofishing effort and settings (voltage, pulse).

Index sites were **electrofished** upstream (single pass) without blocknets. One person operated a backpack electroshocker with a netted electrode while a second person captured fish with a dip-net. Methods for collecting fish data were consistent with the methods described in Task 2.2. **Salmonid** catch rate (**fish/min.**) was calculated for each index site. Except northern squawfish, non-salmonids were counted but not captured.

Task 2.4: Evaluate the use of snorkeling for enumerating salmonids.

We evaluated snorkeling as a technique to enumerate juvenile salmonids. We examined the comparability of snorkeling data to electrofishing data, suitability of snorkeling techniques to stream conditions, and expense and time of obtaining gear and training snorkelers.

Task 2.5: Scale Analysis

See Task 9.1.

Task 2.6: Estimate total number of salmonids in each stream reach, stream, and subbasin.

The total populations of juvenile summer steelhead and spring chinook salmon for the Umatilla River Basin were estimated by expanding quantitative electrofishing and habitat data collected during the summers of 1993-95 (as detailed in Tasks 1.1-1.6 and 2.1-2.3). Additional population estimates were made by comparing streams with empirical data to those not yet sampled quantitatively (Table B-1). We estimated populations for summer steelhead ages 0+ through 3 + and for spring chinook salmon ages 0+ and 1+ (age 1 + denoting a fish having one **annulus** and in its second season of **growth**).

OBJECTIVES 3 and 4: Adult Passage Evaluations

Tasks 3.1 and 4.1: Evaluate the upstream migration of radio tagged adult salmon and summer steelhead past the irrigation diversions in the lower Umatilla River, and evaluate movements of radio tagged adult spring chinook salmon and summer steelhead following upstream transport.

CTUIR initiated a study in 1992 to evaluate adult **salmonid** passage in the lower Umatilla River with radio telemetry. The first year of the project was intended to function as a feasibility study and was conducted on a small scale. This project has since expanded. Fixed-site receivers were installed at key locations and **salmonid** movement following upstream transport was evaluated.

Radio telemetry work on the Umatilla River encompassed the entire Umatilla River and tributaries upstream of TMD. Primary emphasis was given to five major irrigation diversion dams. These include Maxwell Dam (**RM 15.2**), Dillon Dam (**RM 24.6**), Westland Dam (**RM 27.2**), Feed Canal Dam (**RM 28.2**), and Stanfield Dam (**RM 32.4**; Figure A-2).

The radio telemetry portion of this project involves two separate evaluations of adult **salmonid** movements. The "passage evaluation" (Task 3.1) evaluates migration of adult summer steelhead, **coho**, and spring and fall chinook salmon from Three Mile Falls Dam (**TMD**) to above Stanfield Dam. The "upstream transport evaluation" (Task 4. 1), evaluates the movements of summer steelhead and spring chinook salmon following upstream transport and release.

Fish utilized for the radio telemetry project were captured in the TMD adult trapping facility (east-side) and anesthetized **with** carbon-dioxide. Radio transmitters were inserted into the stomach. Individually tagged fish were either released in the **forebay** directly above TMD (passage evaluation) or placed in a truck for transport upstream (upstream transport evaluation). Transported fish were released at either Nolin (**RM 33.6**), Barnhart (**RM 42.2**), Thornhollow (**RM 73.5**), or Imeqes C-mem-ini-kern (Fred Grays, RM 80).

Fish were radio tagged at various times depending on numbers returning to TMD. An attempt was made to radio tag a representative sample throughout the adult return period at low, medium, and high river flows. Coded transmitters were purchased from Lotek Engineering in Newmarket, Ontario, Canada. Radio transmitters were high frequency 150 MHz and varied in size depending on the species being tagged. Summer steelhead and **coho** salmon received transmitters measuring 4.5 centimeters long and 1.7 centimeters in diameter. Fall and spring chinook salmon transmitters were 8.2 centimeters long and 1.7 centimeters in diameter. All radio transmitters had a minimum operating life of approximately 250 days.

Tagged fish were radio-tracked **with** Lotek SRX 400 radio telemetry receivers. Both mobile and fixed-site tracking efforts were employed during the study. Fixed-site receivers (with memory capabilities) were installed at Westland, Feed Canal, and **Stanfield** Dams. An additional receiver was installed near the ODFW district office in Pendleton at RM 56 (ODFW site). Each fixed-site receiver (at diversion dams) included two antennas; one underwater antenna in the fish ladder, and one three-element yagi antenna. Receivers were programmed to alternately scan each antenna for six seconds. This arrangement allowed migrational route (fish ladder or over the dam crest) and arrival and departure times of individual fish at each diversion dam to be determined. Passage times at diversion dams for individual fish were calculated by comparing arrival and departure times. Passage duration through the diversion areas were found by comparing the release time at TMD to the last recorded time at Stanfield Dam (the uppermost diversion).

Most of the mobile radio tracking was conducted in a vehicle equipped with a four-element antenna. On occasion, particularly in areas inaccessible to vehicles, portions of the river were walked with a receiver and hand-held three-element antenna. Once determined, radio tagged fish locations were recorded to the nearest tenth of a river mile.

Migrational movements of radio tagged summer steelhead and spring chinook salmon in relationship to water temperatures and river flows were included in the study. Temperature and flow data were provided by Zimmerman and Duke (1995).

OBJECTIVE 5: Homing and Straying of Adult Salmonids

Task 5.1: Determine factors essential for homing and upstream migration of maturing salmonids.

Available data on returning adult **coho**, fall and spring chinook salmon, and summer steelhead were analyzed in an attempt to understand conditions necessary for successful homing to the Umatilla River. All information related to known Umatilla River origin fish was considered in the search. This included juvenile release data, CWT recoveries, and radio telemetry data. Water flow and temperature data were obtained from Zimmerman and Duke (1995). Homing and straying information represents estimated CWT recoveries from **Rowan** (1995).

OBJECTIVE 6: Spawning Surveys

Task 6.1: Determine final disposition of adult anadromous salmonids released above TMD.

Trap and Haul Project records were reviewed to determine the disposition of all salmonids enumerated at TMD and to determine if adult salmonids released at TMD, after being **caudal** punched, fell back over the dam. Radio telemetry data were also reviewed to determine if radio tagged adult salmonids fell back over TMD after tagging.

Tasks 6.2 and 6.3: Conduct prespawning, spawning, and post spawning surveys throughout the basin for each anadromous species and run;- Estimate the number of successful redds and the adult/redd ratios (female/&d,-female/male) of fish passed above TMD (adjusted for harvest and fall-back, if possible).

Spawning ground surveys to enumerate summer steelhead, spring and fall chinook and coho salmon redds and to sample mortalities were conducted in various reaches of the Umatilla River Basin. Repeated surveys were conducted in areas found to be important for spawning or holding. Other areas were surveyed fewer times or not at all because of low fish abundance observed during previous years or poor survey conditions. Surveyors wore polarized glasses to maximize fish observing capabilities. To minimize stress on prespawning salmonids, crews did not attempt to drive adults from cover for observation by probing debris jams or throwing rocks into pools. The majority of the surveys were conducted by two people, with additional surveyors paired with experienced surveyors during post spawning die-off. Three to four river miles were generally surveyed daily by each person, walking either along the margins of the smaller lateral tributaries. In larger tributaries, surveyors often traversed from bank to bank cover spawning areas and find carcasses.

Redds were judged to be complete (and thus spawning probably successful) based on redd size, depth, location and amount and size of rock moved. All redds were reviewed by our most experienced surveyors for consistency. Redds were marked with orange and white striped flagging. The date, location, species and number of males and females observed on or near the redd were written with permanent marker on the flagging. Writing on the flagging was at least three inches above the lower end of the flag because wind whip caused the ends of the flagging to deteriorate. Flags were placed in trees as close to the redd as possible and at least five feet off the ground to minimize disturbance by wildlife and livestock. In a data book, the surveyors recorded each redd as well as the stream name, location, date, sex and number of fish on or near the redd, carcasses sampled near the redd, and habitat type. Carcasses found during the survey were measured from the middle of the eye to the hypural plate (MEHP). Fork lengths were measured if severe caudal fin erosion had not occurred. Obvious injuries were described and attempts were made to determine the cause of death in prespawning salmonids.

Salmon and steelhead carcasses were cut open to determine egg retention of the females and spawning success of the males. We defined prespawning mortality as death before any spawning had occurred. We classified carcasses as prespawning mortalities only for females with intact skeins and 100% eggs retention and for males with full, corpulent, gonads. Tails of sampled fish were removed at the caudal peduncle to prevent re-sampling. Snouts were removed behind the orbit to recover CWTs from steelhead with both adipose and left ventral (pelvic or pectoral) fin clips, and salmon with adipose fin clips. Snouts were placed in plastic bags and given an individual snout number for identification. The snout card number linked the snout with other biological data collected from the individual fish. Snouts and accompanying biological data were sent to ODFW's Mark Process Center in Clackamas, Oregon, for CWT extraction and reading.

Task 6.4: Calculate fecundity of fish found on spawning grounds. Estimate the number of eggs/redd and total eggs deposited.

The potential egg deposition for natural spring chinook salmon in the Umatilla River was determined from fecundity data from Carson National Fish Hatchery multiplied by redds observed. Estimates of egg retention were subtracted from the total estimated egg deposition. Fecundity of summer steelhead, fall chinook and **coho** salmon were estimated by calculating mean fecundity of salmonids returning to the Umatilla River. Length versus fecundity data were not available for Umatilla River adult returns because eggs were pooled.

Task 6.5: Compare Umatilla Basin spawning survey findings with other salmonid populations in the region.

The standard unit of comparison of adult spawning success in Columbia River tributaries was the total number of redds observed per mile surveyed in index areas, by species.

OBJECTIVE 7: Smolt Trapping

Task 7.1: Install and operate rotary screw traps in Umatilla River below the mouth of Squaw Creek (RM 76) and below the mouth of Birch Creek (RM 48).

We employed two rotary screw traps, five-foot diameter, (E.G. Solutions, Inc. design) to capture emigrating juvenile salmonids. One trap was installed in the Umatilla River on September 21, 1994 at Tumla (RM 76) and was operated from September 21, 1994 to January 13, 1995. After the river channel at the Tumla site was altered by high flows, the trap was moved to the Imeqes C-mem-ini-kern site (RM 79.5) where it was operated from May 5 to June 16, 1995. The second trap was installed in the Umatilla River near **Barnhart (RM 42.2)**. The **Barnhart** trap operated from March 7 to June 1, 1995. The following data were recorded: trap site, date, time, number and species of fish captured, lengths, marks, clips, number of fish marked and released and comments regarding weather, stream flows and trap effectiveness. Scales were subsampled arbitrarily from captured salmonids. Non-salmonid species were counted. We estimates the number of **dace** and shiners when large numbers were trapped. During two occasions at **the Barnhart** site, **the** number of hatchery **coho** captured was estimated volumetrically with a small **dip-net**. We determined the number of **coho/net** from subsamples.

Task 7.2: Install and operate modified pipe traps in Birch Creek.

Pipe traps were not installed or operated in Birch Creek.

Task 7.3: Estimate trap efficiencies.

Trap efficiency rates were estimated by marking salmonids with one of 12 temporary marks. Fish were marked by clipping a notch in the margins of the **caudal** fin, anal fin, dorsal fin or a combination of clips. Marked salmonids were released approximately **100** to 300 m above the rotary traps. Recaptured salmonids were counted, measured and released below the trap. Additional marked juvenile salmonids were placed in the **livewell** for 24 hours to determine

containment rates. Minimizing escapement from the **livewell** through containment monitoring (and immediate repair when necessary) increased effective catch rates. Depending on availability, we used one to 100 fish of a given species and size class for mark-recapture and containment trials.

Trap **efficiency** estimates and total migrants were calculated utilizing two methods. The first method estimated an average capture rate by dividing the number marked fish recaptured by the total number of marked fish released. An estimate of total fish migrating past the trapping site was calculated by dividing total catch by the mean catch rate. Using mean migration rates/day, estimates were generated for times when the trap was not operating. The second method used the average of multiple running means from catch, mark and recapture trials of three to 13 days. The estimate was expanded for times when the trap was not operating by incorporating flow and temperature data and using interpolation techniques.

Assumptions used to estimate trap catch rates and the number of salmonids migrating past the traps include: 1) marked and unmarked salmonids were actively migrating past the trap; 2) fish downstream of the trap did not return to risk capture again; 3) previously captured, handled and marked fish released upstream of the trap had an equal probability of capture as naive unmarked fish; 4) recaptured fish escaped from the **livewell** at the same rate as naive fish; 4) marks on recaptured fish were correctly recognized and recorded by samplers, and 6) no mortality of marked fish occurred between the release site and the trap.

Task 7.4: Freeze brand fish for interrogation in the lower Umatilla and Columbia Rivers in coordination and cooperation with ODFW and the Fish Passage Center.

In agreement with ODFW, freeze branding fish for interrogation in the lower Umatilla and Columbia Rivers was postponed until the fall of 1995. Information will be reported in the **1995-96** progress report.

Task 7.5: Reconstruct emigration timing and minimum survival rates.

Emigration timing was estimated from trapping operations during the past several years. Survival rates were not estimated because Task 7.4 was postponed until the 1995-96 trapping season.

Task 7.6: Design and conduct a mark retention study.

The mark retention study was postponed until the fall of 1995 as it was linked to Tasks 7.4 and 7.5.

OBJECTIVE 8: Tribal Harvest

Tasks 8.1 and 8.2: Design and implement creel and phone surveys to estimate tribal harvest of adult anadromous salmon.

CTUIR fisheries personnel monitored the tribal harvest of adult steelhead in the Umatilla River from December through April, 1995. A roving creel survey was incorporated for harvest monitoring. Survey design followed **the** work of Malvestuto et al. (1978) and Malvestuto (1983). Surveyors recorded the time, location and number of anglers, and the number of fish caught. In

addition, we conducted a selective phone survey with tribal steelhead anglers after the season. There was no tribal season on spring chinook salmon during 1995. Harvest of fall chinook and **coho** salmon was not monitored systematically during the 1994-95 contract year because of the low number of adult salmon and minimal angler effort.

OBJECTIVE 9: Age and Growth

Tasks 9.1 and 9.2: Age analysis of adult and juvenile salmonids.

From adult salmon and steelhead we collected approximately five scales from the preferred area (two rows above the lateral line on the left side of the fish in a diagonal line between **the** posterior edge of the dorsal fin and the anterior edge of **the** anal fin). Additional scales were taken two rows below the lateral line and from the right side of the fish in **the** same areas. Adult scales were mounted on gum cards and pressed in cellulose acetate. In addition to MEHP lengths, we measured fork lengths of adult fish without severe **caudal** fin erosion. Approximately ten scales were collected from juvenile salmonids sampled in **the** preferred area. Scales were mounted between strips of mylar that had been folded in half. Species, fork **length**, date and area captured were written on the left hand edge of the mylar strips with permanent marker. Adult and juvenile scales were analyzed under a microfiche reader at magnifications of 42x and/or 72x. Scales were aged using the European Method of age designation (i.e. age 1.2 was a **fish** that migrated from freshwater during its second year of life, spent two winters rearing in the ocean, and returned to freshwater to spawn at total age four). Scales were read by one or two scale readers. Both readers reviewed scales that were difficult to interpret. Differences in age interpretation were discussed, and if the readers could not agree on an interpretation, the scale was eliminated from the sample. The numbers of circuli to the freshwater **annulus** were determined for 20 known hatchery and 20 unmarked spring chinook salmon in the 1995 escapement in an attempt to separate hatchery from natural returning fish. Age data were collected from a sample of juvenile salmonids captured during biological surveys (all fish were measured). We estimated ages of all juvenile salmonids captured (by five mm increments) from the length and age data of fish subsampled.

OBJECTIVE 10: Genetic and Ecological Effects of Supplementation

Task 10.1: Establish a genetic baseline database from native steelhead.

CTUIR, and Currens and Schreck (1993 1995) sampled juvenile steelhead from 14 locations in **the** Umatilla River during the fall of 1992 and 1994. Workers collected 20-75 steelhead from each location. Currens and Schreck (1995) examined numerous allozymes, mitochondrial DNA, and meristic characteristics.

Task 10.2: Review literature on effects of hatchery-reared salmonids on naturally produced salmonids

Literature regarding **salmonid** interactions was examined,

Task 10.3: Identify acceptable levels of impact from hatchery supplementation on natural steelhead and native trout.

Researchers and managers worked in cooperation during UMEOC meetings to identify methods for measuring, developing criteria for, and monitoring impacts on natural steelhead from supplementation activities.

Tasks 10.4 and 10.5: Examine the utility and feasibility of observing behavior and performance response of naturally produced salmonids in treatment and control areas before and after, and with and without releases of hatchery smolts. Examine the need to study residualization of hatchery smolts and the potential effects on naturally produced salmonids.

Researchers and managers, during several UMEOC meetings, examined the utility and feasibility of conducting residualization studies and monitoring behavioral responses of naturally produced salmonids subjected to hatchery releases in comparison to control groups. Findings of similar work recently conducted in the Columbia River Basin were discussed.

OBJECTIVE 11: Supplementation Effects on Natural Steelhead

Task 11.1: Combine, examine and summarize data gathered in objectives 1-10 that would indicate enhancement of natural steelhead through hatchery supplementation.

We examined production and release data of hatchery steelhead in the **Umatilla** Basin and examined the numbers of returning natural and hatchery adult steelhead. We estimated **the** number of additional natural steelhead that would have been produced if natural adult spawners had not been taken for hatchery brood stock. Production of natural adults was based on ratios of natural adult spawners to resultant natural adult returns to TMD from 1981 through the spring of 1995 (36% to 500% .). No compensatory factors were applied to the estimate as only a five to ten percent increase in adult spawners would have occurred. The proportion of the progeny of each brood year recruiting to subsequent brood years was derived from adult steelhead age data (Table H-2, and I-1, CTUIR et al 1994, Contor et al. 1995).

Task 11.2: Examine potential tests to better evaluate supplementation.

Potential methods to evaluate the effects of supplementation were examined and discussed with experts throughout the pacific northwest and at the UMEOC meetings.

RESULTS AND DISCUSSION

OBJECTIVE 1: Habitat Surveys

Task 1.1: Habitat surveys.

Umatilla River

Habitat surveys were conducted from the upper Umatilla Indian Reservation Boundary (RM 81.8) to the mouth of the North Fork of the Umatilla River (RM 89.6) from July 18 to August 7, 1995 (Tables D-1 through D-8). Habitat crews surveyed 151,949 m² of stream area. Elevation ranged from 1,880 feet at the upper reservation boundary to 2,320 feet at the forks (56 feet/mile). Crews classified and inventoried 639 habitat units. Nine additional habitat units totaling 2,053 m² were identified later by electrofishing crews. These obscure units were isolated pools lateral to the mainstem. The streambed slope averaged 1.4%. The highest water temperature recorded during habitat surveys was 32°C (89.6°F) at Bingham Hot Springs near RM 86.6. The second highest water temperature recorded was 21°C (70°F) near RM 84.8 while the lowest was 10°C (50°F) near RM 85.6. Water temperature and habitat conditions were suitable for salmonids throughout the river section excluding Bingham Hot Springs.

Fast water habitat accounted for 60.3% of the wetted area surveyed. Riffle habitat comprised the most fast water habitat followed by riffles with pockets, rapids over boulders and rapid over bedrock. The average depth of fast water habitat was 0.27 m. Slow water habitat comprised 38.5% of the area. Lateral scour pools comprised the most slow water habitat followed by straight scour pools, glides, and isolated pools. The average maximum depth of slow water habitat types was 0.65 m. Dry channel accounted for 0.3% of the area surveyed (Table D-3).

Secondary (braided) channels accounted for 31.4% of the channel length and 12.8% of the total area surveyed. The average width of the active channel was 2.0 times that of the wetted channel width. The average width to depth ratio of the wetted channel was 22.6:1. The width to depth ratio for riffles was 35.4: 1. The streambank was undercut 8.6% and eroded 7.1% (by length; Table D-2). Gravel (2-64 mm) was the most abundant type of substrate, comprising 35% (53,182 m²) of the wetted streambed area. Spawning gravel abundance does not limit salmonid natural production.

The ground cover in the riparian zone was 39% shrubs, 35% grasses and 26% bedrock and exposed soil (Table D-6). Low terraces were dominant and high terraces were secondary in riparian transects. Many of the high terraces were roads and dikes. The artificial terraces constrain the channel and disrupt the meandering and energy distribution of the river. The stream's power was no longer diffused throughout the flood plain during floods. The concentration of flows by channelization contributes to increased scour and bank erosion. Scouring of redds was suspected to frequently cause mortality of fall chinook and coho salmon eggs in the mainstem Umatilla River.

Hardwoods were the most abundant trees in the riparian zone (71.8%), but tree density was low (3.3 trees/100 m²). Most trees (77%) were 3-15 cm in diameter at breast height (DBH) while only 14.9% were 30 cm DBH or more (Table D-6). Riparian canopy ranged 28 to 31% while percent open sky averaged 49%. The harvest and clearing of trees reduced canopy in this reach. Large woody debris in the river channel averaged only 1.5 pieces/100 m and provided little fish habitat (Table D-5).

A total of 27 surface springs (**3.5/mile**) were observed. Nineteen provided off channel **salmonid** habitat. Eight smaller springs contributed cold water to the mainstem. The highest concentration of springs (**9.1/mile**) was between RM 85.5 and 86.6. Bingham Hot Springs (**RM 86.6; 36°C; 96.8°F**) contributed about 2% (one cfs) of the **mainstem** flow. Five small, screened, irrigation pumps extracted water directly from the river (**RM 81.9 to 87.6**; Tables D-7 and D-8).

Moonshine Creek

Habitat surveys were conducted on Moonshine Creek from the mouth to the forks (**RM 4.4**) from August 28 to September 5, 1995 (Tables D-1, D-2 and D-9 through D-13). The total stream area surveyed was **11,213 m²**. Elevation ranged **from** 1,400 feet at the mouth to 2,590 feet at the forks (270 feet/mile). Crews classified and inventoried 594 habitat units. Streambed slope averaged 2.7%. The highest water temperature recorded during habitat surveys was 23°C (**73.4°F**) while **the** lowest was **10°C (50°F)**. Habitat was marginal for salmonids throughout the entire 4.4 miles.

The stream channel was mostly dry (58% by area), followed by slow and fast water habitat (23 and 18% respectively). Lateral scour pools were the most abundant slow water habitat, followed by beaver dam pools, glides, straight scour pools and puddled areas (0.24 mean maximum depth). Riffles were the most abundant fast water habitat followed by riffles with pockets and rapids over boulders (0.07 m mean depth).

The stream was often confined by terraces and had few braided channels (3.9% by length **2.1%** by wetted area). The active channel width was 3.4 times the wetted channel width. The wetted width to depth ratio averaged 8.9: 1 for all units and 20.0: 1 for riffles. The streambank was undercut 6.0% and eroded 6.0% (by length). Gravel was abundant and comprising 36% (**4,037 m²**) of the wetted streambed area. Spawning gravel abundance does not limit **salmonid** natural production (Table D-1 1).

Ground cover in the riparian zone was 5 1% grasses, 44% shrubs, and 4% exposed soil. Grain fields and stubble were recorded as grasses so the riparian area was in poorer condition than indicated. Agricultural soils are often exposed during winter and spring when erosion potential is highest. Erosion from agricultural fields appeared to be the primary source of sediment to the creek. Riparian canopy was lowest (6 to 27%) farther from the stream. The ground farthest from **the** stream (riparian transect zones two and three) had often been cleared for agricultural uses. Percent open sky averaged 44%. High terraces were the most abundant **landform** within the riparian zone. Most terraces were recently formed by bank erosion and down-cutting (Tables **D-11** and **D-12**).

The trees in the riparian area (**3.2 trees/100 m²**) were mostly hardwoods (99%). Most trees were small (**68%**, 3-15 cm DBH), only 16.3% were 30 cm DBH or more (Table D-12). The low tree density in **the** riparian zone correlated **with** the low woody debris count (1.2 **pieces/100 m**) and the deficiencies of **instream** structure and **salmonid** habitat (Table D-1 1). A total of 27 surface springs were identified (**6.1/mile**; Table D-). These springs contributed cold water to the stream but were too small to provide any off-channel **salmonid** habitat.

The following three passage barriers were found: a natural bedrock step 0.9 m in height (**RM 0.4**); a 0.7 m step formed by a concrete road bridge support near RM 1.0, and a 0.9 m step formed by a log near RM 1.3 (**Table E-23**). Fish passage might be improved with channel or structure modifications at **these** locations.

Mission Creek

Habitat surveys were conducted on Mission Creek from the mouth to the forks RM (4.3) from August 15 to September 11, 1995 (Tables **D-1**; D-2 and D-14 through D-18). The **total** stream area surveyed was 9,994 m². Elevation ranged from 1,270 feet at the mouth to 2,200 feet at the forks (216 feet/mile.). Crews classified and inventoried 872 habitat units. The average slope was 2.8 %. The highest water temperature recorded during habitat surveys was 14°C (**57.2°F**) while the lowest was 6°C (**42.8°F**). Habitat was marginal for salmonids throughout the entire stream.

Dry channel accounted for 76.3% of **the** area surveyed. Slow water habitat accounted for 12.0% of the area surveyed. Lateral scour pools were the most abundant slow water type, followed by straight scour pools and puddled channels. Maximum depth of slow water habitat averaged 0.18 m. Fast water habitat accounted for 11.4% of **the** area. Riffles comprised the most area, followed by rapids over boulders and riffles **with** pockets. The average depth of fast water habitat types was 0.05 m (Table D-14).

Secondary (braided) channels accounted for 3.0% of the channel length and 2.3% of the wetted area. Active channel width averaged 2.5 times wetted channel width. Width to **depth** ratio of all units averaged **9.3:1** and **32.9:1** for riffles. The streambank was undercut 8.2% and eroded 21.3% (by length). Gravel was the most abundant wetted substrate (4,394 m², 44% of the area; Tables D-15 and D-16). Fines comprised 24% of the wetted area. Spawning gravel abundance does not limit **salmonid** natural production.

The ground cover in the riparian transects averaged 58% grasses, 18% shrubs and 24% exposed soil. Grain fields and stubble were recorded as grasses so the riparian area was in poorer condition than indicated. Agricultural fields are **often** exposed during winter and spring when erosion potential is highest. Erosion from agricultural fields and effects from livestock grazing appeared to be the primary source of sediment. Riparian canopy was lowest (42%) farther from the stream. The percent open sky averaged 38% (Table). High terrace and hill-slope were the most abundant **landform** in **the** riparian zone (Tables D-16 and D-17). Most high terraces were recently formed by bank erosion and down-cutting.

Hardwoods were the most abundant tree type (94.6%) in the riparian area, but tree densities were low (2.9 **trees/100 m²**). Most trees (77.3%) were in the **3-15cm** DBH range, only 10.0% were 30 cm DBH or more (Table D-17). Low tree density in the riparian zone correlated with **the** low woody debris count (6.6 **pieces/100 m**) and inadequate **instream** structure for **salmonid** habitat (Table D-16). Twenty-one surface springs were identified (**4.9/mile**). The springs were too small to provide off-channel **salmonid** habitat but contributed cold water to the stream (Table D-18).

No water diversions were observed. However, two wells near RM 0.5 and 4.1 may affect **instream** flows. The temperature of the well water was **10.5°C (50.9°F)**, whereas the temperature of the creek was **12.5°C (54.5°F)**. The impacts of these wells to stream flows remains unknown.

Seven potential passage barriers were found. Four were artificial structures and three were natural (Table E-23). It appeared that **the** barriers would significantly impede migration at moderate to high flows and completely block it at low flow. Improvements in fish passage might be achieved through installation of log check dams or structure modification. The most severe artificial barriers were at the bridge near RM 1.4 and at the culvert near RM 3.3.




Figure C-1. Maximum and Minimum **Temperatures** Recorded in the Umatilla River, Near Rieth, RM 49.5, December 94 through May 1995 (TGUR9412.CH3).




Figure C-2. Maximum and Minimum **Temperatures** Recorded in the Umatilla River, Bamhart, RM 42.5, February Through June, 1995 (TCUB9502.CH3).

3.1 %. The highest water temperature (29°C) was recorded at the mouth of an un-named tributary near RM 0.9 while the lowest (11°C) was recorded -in three springs (RM 0.8, 1.2 and 3.7).

Fast water habitat accounted for 63.2% of the area. Riffles were the most abundant fast water habitat, followed by riffles with pockets and rapids over boulders. The depth of fast water habitat types averaged 0.10 m. Slow water habitat accounted for 36.2% of the area. Lateral scour pools comprised the most area, followed by straight scour pools and glides. The maximum depth of slow water habitat types averaged 0.28 m (Table D-24). Only 0.2% of the stream area was dry. Sampling Coonskin Creek earlier in the summer than the adjacent tributaries may explain the low percent of dry channel area. Water temperature and habitat was marginal for salmonids throughout the stream.

Secondary (braided) channels accounted for 7.9% of the channel length and 10.4% of the wetted area. The width of the active channel was 2.5 times the wetted width. The width to depth ratio of all units averaged **7.6:1** but averaged **19.2:1** for riffles. The streambank was undercut 11.2% and eroded 13.2% (by length). Gravel was the most abundant type of substrate and comprised 34% (1,992 **m²**) of the wetted streambed area followed by fines (31%; Table D-25 and D-26). Spawning gravel abundance does not limit **salmonid** natural production.

The ground cover in the riparian zone was 49% grasses, 43% shrubs and 8% exposed soil. Many of the grasses were actually grain crops. While crops stabilize fields during the growing season, agricultural soils are often exposed during winter and spring when erosion potential is highest. Erosion from agricultural fields appeared to be the primary source of sediment. Riparian canopy (15-31%) was lower further from the stream. Clearing of trees from the riparian area for agricultural uses was common. Percent open sky averaged 41% (Tables D-26 and D-27).

Low and high terraces were the most common **landform** in the riparian transects. Many of the terraces recently formed from bank erosion and down-cutting. Hardwoods were the most abundant trees (98.8%) but tree density was low (2.8 **trees/100 m²**). Most trees (73.5%) were in the **3-15cm** DBH range, and only 15.7% were 30 cm DBH or more (Table D-27). The lack of trees in the riparian zone correlated with the lack of large woody debris (1.6 pieces/100 m) and the deficiencies in fish habitat (Table D-26). Crews observed 17 springs contributing cold water to the stream (**8.5/mile**; Table D-28). The springs were too small to provide off-channel **salmonid** habitat.

Eleven passage barriers were found. Most barriers resulted from down-cutting of the channel below clay layers. We estimate that the barriers impeded migration at high and moderate flows and completely blocked migration at low flow. The barriers ranged from 0.65 m to 1.65 m in height. Near RM 0.4 a concrete structure (0.8 m high) protecting Pendleton's water pipe was recently modified so that it further diminished fish passage (Table E-23).

Task 1.2: Stream temperatures and stream flow in the Umatilla Basin.

Temperatures

Stream temperature profiles collected throughout the Umatilla River Basin were plotted in Appendix C (Figures C-1 through C-9). Water temperatures became unsuitable (above **20°C**, **68°F**) for salmonids during the summer below RM 70 in the Umatilla River and in the lower ends of many of the tributaries. For example, in the Umatilla River at RM 42.5 and 49, water temperatures were well above **20°C** (Figures C-1 through C-3). In Wildhorse Creek at RM 1.5, water temperatures were above 25°C (**77°F**) in July and August. Higher in the basin, temperatures were suitable for salmonids throughout the year. In Mission Creek, at RM 3, water temperatures did not exceed 16°C (**61°F**) during July and August 1995. In several locations, a spring or cool

tributary infused enough cool water to provide suitable flows and temperatures for several hundred feet to several miles downstream. The North Forks of the Umatilla River and Meacham Creek are examples of this.

The riparian canopy along many reaches in the Umatilla River Basin was minimal and provided little shade to the streams. Direct solar radiation and total water volume play the greatest roles in stream temperature dynamics (Brown 1983). Removing large trees from stream areas has been shown to increase maximum stream temperatures in test streams from a maximum of **15.6°C** (60°F) before vegetation removal to **30°C (86°F)** after removal. Control reaches had no significant changes during the same time period (Brown and Krygier 1970). Shallow, unshaded pools and glides are typical to much of the Umatilla River and function as efficient solar energy collectors and water temperatures can become too warm for salmonids (Brett 1952, Black 1953).

Flow

A strong correlation existed between mean annual (**$r=0.913$**) and spring flows (**$r=0.869$**) at the Umatilla gage (**RM 1.2**) and the number natural adult steelhead returning two years later from return years 1982-83 to 1994-95 (Figures B-1 and B-2). Assuming the relationship between spring **instream** flows and the number of returning adult steelhead remains consistent, approximately 2,000 adult natural and hatchery steelhead will return during the 1995-96 season with 1,400 and 1,800 steelhead expected to return during the 1996-97 and 1997-98 seasons respectively.

Tasks 1.3 through 1.5: Obtain habitat data collected by other agencies. Digitize and summarize habitat data. Estimate total usable habitat by stream reach, drainage and entire basin.

Data from habitat surveys conducted by ODFW in 1991 and 1992 on Umatilla River Basin tributaries were obtained on computer diskette. No additional data entry or summarization was required. Raw habitat data collected and recorded in the field were entered into a data base program. Habitat data summaries were listed in Appendix D.

Estimates of **salmonid** summer rearing habitat by stream reach, drainage and basin were summarized in Table B-1. Approximately 30% (233 of 770 stream miles) of the **salmonid** habitat in the Umatilla River Basin is suitable for natural production. De-watering, sedimentation, poor water quality and/or excessive water temperatures were the primary reasons 70% of the 770 miles were rated unsuitable. We do not know how much habitat was available historically for **salmonid** production. We speculate that 70% (540 of 770 stream miles) of the drainage may have been suitable for summer rearing of salmonids. The remaining 30% of the streams include portions of subbasins such as Wildhorse Creek, Butter Creek, Alkali Canyon, Spear Canyon and Coombs Canyon. Currently, these streams (many are ephemeral) flow from desert uplands and presumably never supported salmonids during the summer.

Task 1.6: Coordinate water quality monitoring efforts in the Lower Umatilla River with the Oregon Department of Environmental Quality.

Water quality monitoring is currently being conducted by CTUIR, ODFW, USFS, DEQ and BOR. CTUIR monitors temperatures and sediment through this project, the Habitat Project and the Artificial Production Program (Appendix C). ODFW, BOR and USFS also monitor water temperatures in the Umatilla River Basin. DEQ monitors several sites in the Umatilla River for 45

heavy metals, conductivity, **pH**, total alkalinity, nitrogen, total organic carbon, phosphorous, hardness and others. DEQ and CTUIR, in cooperation with the Umatilla Basin Watershed Council, will begin more intensive water quality monitoring in April, 1996. As data are collected and examined, recommendations regarding point source and non-point source pollution allocation and management for reducing pollutants will come from the newly formed Umatilla River Total Maximum Daily Load Technical Advisory Committee.

OBJECTIVE 2: Biological Surveys

Task 2.1: Conduct presence/absence surveys in the Umatilla River Basin.

A fish survey was conducted in Coyote Creek and in an un-named tributary that enters the North Fork Umatilla River from the north at RM 1.5 (March 24, 1995). Time and personnel constraints limited additional presence/absence surveys.

Coyote Creek (4°C; **39.2°F**) was electrofished for 380 seconds from the mouth to approximately 300 m upstream. Pools with adequate cover for **fish** were sampled. Crews captured seven steelhead (61 to 148 mm) in poor condition. Approximately ten sculpin were sighted. Stream and riparian habitat conditions appeared excellent for salmonids. Pools and large **instream** woody debris were abundant.

The un-named tributary (5°C; 41°F) was electrofished for 180 seconds from the mouth to 200 m upstream. Pools and pockets were sampled. **One** steelhead was captured (99 mm). No other fish were sighted. Riparian conditions appeared good and stream habitat appeared fair for salmonids. Rapids were the most common habitat type.

Task 2.2: Estimate salmonid densities in streams where habitat has been surveyed by electrofishing.

Umatilla River

The Umatilla River was subsampled for fish from the upper Umatilla Indian Reservation Boundary (**RM 8 1.8**) to the mouth of the North Fork of the Umatilla River (**RM 89.6**) from August 8 to August 25, 1995. Salmonids were captured from RM 81.9-89.3. The highest water temperature recorded in the **mainstem** during fish surveys was **19°C (66.2°F)** near RM 83.2 while the lowest was **9.5°C (49°F; RM 88.3)**. Based on **salmonid** densities, this section of the Umatilla River appeared to be an important rearing area for juvenile steelhead, chinook salmon and mountain whitefish.

We sampled 72 of 643 habitat units (11.1% by units, 6.7% by area). Thirteen of 17 habitat types were electrofished (dry units and steps were excluded). A total of 2,234 of the following salmonids were captured: 1,899 (78.5%) natural steelhead trout; 327 (13.5%) juvenile natural chinook salmon; 185 (7.6%) mountain whitefish, and five (0.2%) bull trout. The bull trout were captured from pools or pocket water between RM 87.7 and 89.2.

The expanded population estimate was 69,116 salmonids with a mean density of 0.45 **salmonids/m² (s/m²)**; Tables E-1 and E-1 1). Juvenile **salmonid** densities in slow water units averaged 0.52 **s/m²** and averaged 0.40 **s/m²** in fast water units (Table E-6). Lateral scour pools had a mean density of 0.87 **s/m²**, and a single dam pool had a density of 1.77 **s/m²**. An increase in pool and pocket water habitat would likely increase natural production of salmonids.

Fork lengths of captured salmonids ranged from 29-258 mm for natural steelhead trout, 65-127 mm for natural juvenile chinook salmon, **116-440** mm for mountain whitefish, and 170-265 mm for bull trout (Table E-12, Figures E-1 and E-2). Fifty-six percent of the whitefish captured were from slow water habitat where mean density was twice as high as in fast water habitat. The highest mean density of whitefish was estimated in plunge pool habitat (0.12731 m^2). Whitefish were captured from RM 82.2-88.7, most were near RM 87.7.

Electrofishing and handling caused observed mortality of 2.8% of the captured natural chinook salmon juveniles, 1.9% of natural steelhead and 0.5% of mountain whitefish. Scissor and puncture wounds from avian predators were observed on a few salmonids (0.11 to 2.2%) including three chinook (mean length 88 mm), two steelhead (208 mm), and four mountain whitefish (336 mm).

The population estimate of non-salmonid was 151,511 fish. The ratio of non-salmonid to **salmonid** was 2.2: 1. Speckled **dace** and **redside** shiners were the most abundant of non-salmonids (comprising **98.9%**, Table E-17). Six northern squawfish (112-170 mm) were captured in an isolated pool with a spring seep; their stomachs contained insects, sculpins and snails.

Moonshine Creek

Salmonids were captured by electrofishing in Moonshine Creek from the mouth to RM 4.4 (September 18 to 21, 1995). The highest water temperature recorded was **18.5°C (65.3°F)** near RM 1 while the lowest (**11.5°C, 52.7°F**) was recorded from a spring near RM 0.1. Moonshine Creek appeared to be an important rearing area for steelhead and of lesser importance to **coho** and chinook salmon.

The following numbers of juvenile salmonids were captured: 369 (97.4696, 48-240 mm) natural steelhead trout; six (**2.4%**, 88-95 mm) natural **coho** salmon, and one (**0.3%**, 88 mm) natural chinook salmon (Tables E-2, E-13 and Figure E-3). Juvenile **coho** and chinook salmon likely migrated into the creek from the **mainstem** Umatilla River. All salmon were captured from one scour pool near RM 0.2.

Fourteen habitat types and 89 of 526 habitat units were sampled (15.0% by units and 9.9% by area). The expanded population estimate was 1,169 salmonids and mean density was 0.10 s/m^2 (Table E-7). The **salmonid** density of slow water units was 2.1 times higher than in fast water units. Plunge and trench pools had mean densities of 2.22 and 1.86 s/m^2 , respectively. The density of salmonids in riffles with pockets was 12.5 times as high as in riffles. Increase in pool and pocket water habitat would likely increase **salmonid** production.

Electrofishing and handling caused observed mortality of 0.81% of the captured natural steelhead. A scissor bite was observed on one steelhead (165 mm). The expanded population estimate of non-salmonids was 10,340 fish. The ratio of non-salmonid to **salmonid** was **8.8:1** (Table 18). Suckers were the most abundant non-salmonids and were concentrated near the confluence with the Umatilla River. Sculpins and speckled **dace** were not as numerous, but were distributed throughout the stream.

Mission Creek

Fish surveys were conducted in Mission Creek from the mouth to the forks (**RM 4.3**) from September 5 to 13, 1995. Salmonids were captured from RM **0.4-4.2**. The maximum water temperature recorded was 21°C (**70°F**) near RM 0.6 while the lowest was (**11.5°C, 52.7°F**) from a spring near RM 4.1. Mission Creek appeared to be important for juvenile steelhead and of moderate value to **coho** salmon. Ten habitat types and 65 of 641 habitat units were sampled (7.5% by units and 4.4% by area). The expanded population estimate was 903 salmonids with

mean **salmonid** density of 0.093 **s/m²** (Table E-3). The density of slow water units was 14 times as high as in fast water units. Plunge pools had the highest density of any habitat type with an estimated density of 1.62 **s/m²** (Table E-8). **Salmonid** density in riffles with pockets was six times higher than in riffles. Increasing pool and pocket water habitat would likely increase the **salmonid** natural production.

Crews captured 202 natural steelhead trout (90.2%; **56-290** mm), 21 natural **coho** salmon (**9.4%**, 88-95 mm) and one hatchery steelhead (**0.4%**, 230 mm). This was the only hatchery steelhead captured during any of the biological surveys conducted from June 29 to September 21, 1995 (Table E-14 and Figure E-4). All **coho** salmon were captured in pools near RM 0.5. Juvenile **coho** and chinook salmon presumably migrated into the creek from the **mainstem** Umatilla River where spawning has been documented.

Electrofishing and handling caused observed mortality of 0.50% of the captured natural steelhead. The population estimate of non-salmonids was 10,326. The ratio of non-salmonid to **salmonid** was 11.1: 1 (Table E-19). Speckled **dace** (76.9%) were the most abundant non-salmonid followed by sculpins and **redside** shiners.

Cottonwood Creek

Fish surveys were conducted in Cottonwood Creek from the mouth to the forks (**RM** 4.1) from July 5 to August 1, 1995. Salmonids were captured from RM 0.0-3.1. The highest water temperature recorded was 24°C (**75.2°F**) near RM 2.9 while the lowest was **8.5°C** (47.3°F) from a spring near RM 0.2. Cottonwood Creek appeared to be an important rearing area for steelhead and of moderate value to **coho** salmon.

The following juvenile salmonids were captured: 172 natural steelhead trout (**78.2%**, 37-340 mm); 47 natural **coho** salmon (**21.4%**, **69-103** mm), and one natural chinook salmon (**0.46%**, 63 mm). Juvenile **coho** and chinook salmon may migrate from the **mainstem** Umatilla River where spawning has been documented. Ninety-eight percent of the salmon captured were found in pools in the lower 1.1 miles of the creek (Table E-4, E-15 and Figure E-5).

Fourteen habitat types were sampled from 70 of 769 units (7.7% by number and 18.3% by area). The expanded population estimate was 626 salmonids. The mean density estimated for the entire area of stream was 0.04 **s/m²** (Table E-9). The mean **salmonid** density in slow water units was 2.1 times higher than in fast water units. The density of salmonids in riffles with pockets was 4.2 times higher than in riffles. This suggested that an increase in the amount of pool and pocket water could increase the number of salmonids in the stream section.

Electrofishing and handling caused observed mortality of 1.74% of the captured natural steelhead. A scissor bite was observed on one steelhead (211 mm). The population estimate of non-salmonids in the survey section was 8,937. The ratio of non-salmonid to **salmonid** was 11.9: 1 (Table E-20). Speckled **dace** (85.1%) were the most abundant non-salmonid followed by sculpins, **redside** shiners and suckers.

Coonskin Creek

Salmonids were captured in Coonskin Creek from the mouth to RM 3.7 (June 29 to July 18, 1995). The highest water temperature recorded was **27.5°C (81.5°F)** near RM 0.8 while the lowest was 11°C (**51.8°F**) near RM 0.4. Near RM 0.1, the water temperature was 11.5°C (**52.7°F**) under a developed canopy but was **17.5°C (63.5°F)** only 30 m upstream where a wheat field directly bordered the stream. Coonskin Creek appeared to be an important rearing area for steelhead and **coho** salmon and of moderate value to chinook salmon (Table E-5).

The following numbers of juvenile salmonids were captured: 311 natural steelhead trout (**76.0%**, 42-327 mm); 86 natural **coho** salmon (**21.0%**, 64-90 mm), and 12 natural chinook salmon (2.9% **74-90** mm). Eighty-one percent of the salmon captured were found in pools between RM 0.1 and 0.2 (Table E-10, E-16 and Figure E-6). Juvenile **coho** and chinook salmon may migrate into the creek from the **mainstem** Umatilla River where spawning has been documented.,

Twelve habitat types were sampled from 88 of 592 units (14.1% by number and 15.4% by area). The population estimate in the survey area was 1,875 **salmonids**. The mean density estimate for the entire stream was 0.320 **s/m²** (Table E-10). The mean **salmonid** density in slow water units was 5.9 times higher than in fast water units. The density of salmonids in riffles with pockets averaged 1.8 times higher than riffles. Increasing in the amount of pool and pocket water might increase **salmonid** natural production.

Electrofishing and handling caused observed mortality of 8.33% of the captured natural chinook salmon juveniles, 2.32% of natural **coho** salmon juveniles and 0.64% of natural steelhead. A puncture wound was observed on one natural steelhead (151 mm). The population estimate of non-salmonids was 1,955 fish. The ratio of non-salmonids to salmonids was 1: 1 (Table E-21). Speckled **dace** (71.2%) were the most abundant non-salmonid followed by sculpins.

Task 2.3: Electrofish permanent index sites during November, April and August.

Index sites with the highest average catch rate during the three sample periods were: East Birch Creek (3.4 **fish/min.**); Boston Canyon Creek (3.2 **fish/min.**); Spring Creek (3.1 **fish/min.**), and Squaw Creek (site 27, 3.0 **fish/min.**). Ryan Creek had a high catch rate (5.1 **fish/min.**) but was only sampled once (Table E-22). In general, the highest catch rates during August were in the upper tributaries of the Umatilla River. During November, tributaries of Birch Creek had the highest catch rates. Most salmonids were captured in slow water, near the bank, during March and November.

During index surveys, crews captured steelhead, chinook salmon, **coho** salmon, mountain whitefish, **american** shad, speckled **dace**, **redside** shiners, northern squawfish, chiselmouth, suckers, sculpins, smallmouth bass and carp. Several passage barriers were found during index surveys and were listed in Table E-23. Modifications to some barriers would allow salmonids access to additional rearing area.

March and April

Field conditions were generally poor for sampling at most sites during March and April because of moderate to high flows. Sampling was often restricted to the stream margins. Low catch rates were frequent. The Ryan Creek index site (37) was not sampled because of poor accessibility.

Natural steelhead were not collected in the spring at index sites downstream of RM 74 (site 8) nor were natural chinook salmon collected below RM 88 (site 10). No natural **coho** salmon were observed; however, 44 hatchery **coho** salmon were collected at RM 9 (site 2). One mountain whitefish (167 mm) was collected at RM 25 (site 3). The highest **salmonid** catch rates were in Line Creek (3.3 **fish/min.**), Boston Canyon Creek (2.7 **fish/min.**), East Birch Creek (1.9 **fish/min.**), and the Umatilla River, RM 9.0 (site 2; 1.9 fish/min.).

August and September

Field conditions were good for sampling during August and September. The Ryan Creek site (37) was not sampled. Seventy-eight young-of-the-year (YOY) shad, 33 YOY carp and 14 smallmouth bass were captured at RM 1.5 (site 1). Five naturally produced **coho** juveniles were captured from an isolated pool with a spring seep at RM 38 (site 4).

During summer index monitoring, natural steelhead were not observed below RM 50 (site 5) nor were natural chinook salmon collected below RM 88 (site 10). Natural **coho** salmon were not collected below RM 67.5 (site 7). The highest catch **salmonid** rates were in Squaw Creek (site 27; 6.7 fish/mitt.), Meacham Creek (site 34; 5.3 **fish/min.**), East Meacham Creek (4.0 **fish/min.**), and the South Fork Umatilla River (site 13; 4.0 fish/min.). Boulders to improve **salmonid** habitat altered the site in East Birch Creek (**RM** 4.5, site 19).

November

Field conditions were poor for sampling during November due to high flows. In most cases, sampling was restricted to the stream margin. Most salmonids were captured in slow water, with undercut, root wads or woody debris. Many of the fish appeared to have been actively feeding. The following sites were not sampled in November due to flooding: South Fork Umatilla River (site **13**), North Fork Meacham Creek (site **33**), East Fork Meacham Creek (site 35) and Shimmiehorn Creek (site 40). Four adult fall chinook salmon, one adult steelhead, three mountain whitefish and many adult suckers were present in the isolated pool at site one. We did not electrofish over the salmon redds at site one. Many large cottonwood trees in the riparian area at site three had been cut down and removed. An adult fall chinook salmon was observed at site three. A fall chinook or **coho** salmon was occupying a redd at site four. Numerous YOY squawfish were rearing in the backwater pool with a spring seep at site four.

During fall sampling, natural steelhead were not observed below RM 50 (site 5) nor were natural chinook salmon collected below RM 88 (site 10). Natural **coho** salmon were not collected below RM 67.7 (site 7). **The** streams with the highest catch rates were Ryan Creek (5.1 fish/min.), Bear Creek 5.0 **fish/min.**), East Birch Creek (4.9 **fish/min.**), and Pearson Creek (4.4 **fish/min.**). **Salmonid** habitat improvement projects (gravel removal and boulder placements) altered the index sites in Birch Creek (**RM** 10, site 16) and West Birch Creek (**RM** 2, site 17).

Task 2.4: Evaluate the use of snorkeling for enumerating salmonids.

Snorkeling as a technique to enumerate juvenile salmonids has been used successfully by a researchers in Oregon, Washington and Idaho (Petrosky and Holubetz 1987, Bugert et al. 1990, Kucera et al. 1991, Angradi and Contor 1989, **Hillman** and **Mullan** 1989, **Mullan** et al. 1992, Cannamela 1993, Contor and Griffith 1995). However, we found that snorkeling techniques would not meet our data needs and were impractical for many of the streams in the basin. **Salmonid** density estimates from snorkeling techniques would not be directly comparable to existing electrofishing data. Many of the juvenile salmonids captured by electrofishing were extracted from substrate interstitial spaces and would not have been visible to snorkelers estimating **salmonid** abundance. Water was often too shallow (often less than 15 cm) or too turbid for snorkeling enumeration techniques. Snorkeling would also require extensive training and evaluation, yet not provide opportunities to take scales, lengths and weights from salmonids.

Task 2.5: Scale Analysis.

See Task 9.1.

Task 2.6: Estimate total number of salmonids in each stream reach, stream, and subbasin.

The populations of natural juvenile summer steelhead (ages 0+ to 3+) and spring chinook salmon (ages 0+ to 1+) in the Umatilla River Basin were estimated to be near 725,000 and 52,000 respectively. The majority of steelhead rear in Birch Creek (170,000), Meacham Creek (265,000), Squaw Creek (40,000), and the upper Umatilla River (216,000). Natural chinook reared primarily in the North Fork and the upper mainstem (RM 70 to 89.6) of the Umatilla River (41,000) and Meacham Creek (10,000). The estimates should not be considered static or accurate and were based on limited quantitative data (Table B-1). More refined estimates will be possible as additional data are collected. Recognize, that the available habitat and associated salmonid populations expand and contract depending on factors such as, snow pack, summer precipitation, flow and water temperatures.

OBJECTIVE 3: Adult Passage Evaluations.

Task 3.1: Evaluate the upstream migration of radio tagged adult salmon and steelhead past the irrigation diversions in the lower Umatilla River.

Fall Chinook Salmon and Coho Salmon

A total of nine fall chinook salmon were radio tagged and released at TMD between October 6 and 20, 1994. Of these, three successfully migrated over Westland Diversion Dam and one (of the three) successfully negotiated Feed Canal and Stanfield Dams. The remaining six salmon all remained below Westland Dam (RM 27.2).

Between October 12 and 26, 1994, a total of eight coho salmon were radio tagged and released at TMD. Three of these passed Westland Dam and one of the three passed Feed Canal and Stanfield Dams. Of the remaining five coho salmon, one regurgitated the radio transmitter and four remained below Westland Dam.

Peak migration for fall chinook and coho salmon over McNary Dam on the Columbia River has typically occurred in September. Entry dates at TMD have varied but generally follow flows exceeding 150 cfs (Volkman 1994). Umatilla River coho and fall chinook salmon broodstock have typically spawned in early November (Rowan, CTUIR, personal communication). In 1994, flows in the Umatilla River began to increase in early October and most fall chinook and coho salmon arrived in mid to late October. By this time, coho and fall chinook salmon were entering advanced stages of maturation and reduced physical condition. The potential for these fish to successfully migrate to headwater sections of the Umatilla River Basin was remote.

Telemetry data collected in 1994 were indicative of sexually mature fish and portrayed the movements of fish at or near spawning. Evidence that these fish were near spawning was demonstrated by ripe adults at TMD and numerous fall chinook and coho salmon spawning below TMD each fall. If fall chinook and coho salmon are released at TMD in October and November, most will spawn within 20 miles of the release point. Unfortunately, most of the lower Umatilla River does not contain quality spawning and rearing conditions, particularly for coho salmon. If natural production of these species is desired, trapping and hauling may be the best solution until flows are made available in early September.

Summer Steelhead

A total of 30 summer steelhead were radio tagged between October 31, 1994 and May 16, 1995. Of these, 16 provided data past all of the major diversion dams (TMD to above Stanfield Dam), seven could not be located after release, and seven regurgitated the radio transmitter. On average, 36 days were required to migrate from TMD to above Stanfield Dam (Table F-1). Twenty-five days were required to complete this distance in 1993-94. Average migrational passage time (hours and minutes) required to negotiate Westland, Feed Canal, and Stanfield dams were **13:06**, **83:24**, and **2:58** respectively (Table F-1). This compares to **1:30**, **48:54**, and **1:23** in 1993-94 (Figure F-1). Percent of fish migrating through the ladder at each diversion was 38% at Westland, 75% at Feed Canal, and 31% at Stanfield (Table F-1, Figure F-2).

Average migrational passage time between TMD and **Westland** Dam, Feed Canal Dam, Stanfield Dam, and the ODFW site, were 27.2, 29.2, 36.4, and 48.5 days, respectively (Table F-2). Passage times between diversion areas are provided in Figure F-3.

Flow ranges encountered during adult passage were 707 to 2650 cfs at **Westland** Dam, 531 to 2448 cfs at Feed Canal Dam and 662 to 3420 cfs at Stanfield Dam. Migrational delays were documented at Feed Canal Dam at flows ranging from 563 to 1,601 cfs (Table F-1). Some minor delays also occurred at **Westland** and Stanfield Dams in the 1,200 to 1,400 cfs range (Table F-1). Water temperatures encountered during passage for each diversion are presented in Table F-1.

During the last three years, average passage times required to migrate from TMD to above Stanfield Dam have been similar. In 1993, 1994, and **1995**, 30 days, 25 days, and 27 days were required, respectively. Passage times through the Umatilla River were longest for summer steelhead entering early in the migrational period (September through December). Fish entering later in the period, and thus closer to spawning, such as in March or April, migrated through the system more quickly (Figures F-6 and F-7).

In the last two years, nine summer steelhead (22%) could not be located following release at TMD. Although it's possible the radio transmitter failed or the fish were captured, fall-back out of the system is more likely. This may suggest that TMD counts for summer steelhead were **inflated**. Several studies have been conducted at TMD to evaluate fall-back levels. Unfortunately, these experiments only enumerate recaptures. In an effort to understand this uncertainty, CTUIR will install an additional telemetry receiver downstream of TMD for the 1995-96 evaluation.

Migrational delays were again observed at Feed Canal Dam. Passage times in 1994-95 (**83:25**) were considerably longer than those observed in 1993-94 (**48:54**). Although some increased delay was likely in response to high flows and gravel accumulations at the dam, poor facility design remains the primary problem. Feed Canal Dam was designed for water diversion, not fish passage. The large apron on the downstream side of the dam creates false attraction for ascending adults and prevents fish from jumping over the crest of the dam. Because of this, the ability of fish to locate the fish ladder entrance at Feed Canal Dam was of paramount importance. In 1994-95, 75% of the radio tagged summer steelhead passing the facility used the fish ladder. In comparison, 38 % used the ladder at **Westland** Dam and 3 1% at Stanfield Dam.

Data indicated that upstream migrants could not locate the ladder entrance at Feed Canal Dam. The large expanse of the dam compared to the small fish ladder entrance was likely responsible. Strong attraction flows toward the fish ladder may reduce this problem. This, however, would only be a solution during low flows. During high flows, water spills over the entire crest, thus creating attraction away from the fish ladder and again passage delays.

The effect of delay below Feed Canal Dam on upstream migrants is unknown. For summer steelhead returning early in the migrational period, a small. delay is probably insignificant. Late returning steelhead, however, and spring chinook, fall chinook, and **coho** salmon were likely

impacted. Timing for these fish is critical. Migrational delay and repeated attempts to negotiate the structure may be tapping into vital energy reserves needed for spawning. This, in turn, may promote prespawn mortality and impact distance migrated and spawning sites chosen. It should be noted that passage times for Feed Canal Dam only represent fish that successfully negotiate the structure. In each of the last two consecutive years, several radio tagged fish have been unable to negotiate Feed Canal Dam. These fish were thus forced to choose spawning sites downstream of the dam.

Several solutions concerning delays at Feed Canal Dam have been suggested. These include various combinations of additional spill gates, jump pools and fish ladders. Given the continual problems associated with Feed Canal Dam, however, reconstruction or dam removal is likely the best option. In **1994-95**, Feed Canal Dam experienced severe gravel accumulation problems. Gravel accumulations compounded existing passage concerns and required the Irrigation District to conduct **instream** work several times during the migrational period. Its important to understand that gravel accumulations were not directly responsible for passage delays at Feed Canal Dam but rather facility design. Until major modifications are made to Feed Canal Dam, most upstream migrants will be severely delayed with some migrants completely unable to negotiate the structure.

Figure F-3 illustrates that the reach of river did not cause delay but rather the diversion dams within the reach. Clearly, summer steelhead display little difficulty ascending sections of the river without diversion dams. Once encountering sections with dams, migrational movements were considerably reduced. It's interesting that summer steelhead appeared willing to migrate at marginal water temperatures of 4.4 to **6.1°C (40 to 43°F)** through sections of the river without diversion dams, but upon encountering sections with dams, migration either stops or passage time increases.

Spring Chinook Salmon

Between April 10 and 26, 1995 a total of 15 spring chinook salmon were radio tagged at TMD. Of these, nine provided data past Stanfield Dam, two regurgitated the radio tag, three fell back and were recaptured at TMD, and one migrated up to but not past Stanfield Dam. Average time needed to migrate from TMD to above Stanfield Dam was 18 days (Table F-3). Twelve days were needed to complete this distance in 1993-94. Average passage times (hours and minutes) at Westland, Feed Canal, and Stanfield dams were **04:30, 89:42, and 04:01**, respectively (Table F-3). In 1993-94, **01:30, 48:54, and 01:23** were required to complete this distance (Figure F-4). Forty percent of the fish chose to use the fish ladder at Westland, 60% at Feed Canal, and 11% at Stanfield (Table F-3, Figure F-5).

Flows encountered during passage were 796 to 911 cfs at **Westland** Dam, 689 to 2772 cfs at Feed Canal Dam, and 675 to 3,781 cfs at Stanfield Dam. Migrational delays occurred at Feed Canal Dam at flows ranging from 700 to 2,772 cfs. **One** chinook salmon was also delayed at **Westland** Dam at average flows of 796 cfs (Table F-3). No flow-related delays were documented for spring chinook salmon at Stanfield Dam. Water temperature information is provided in Table F-3.

In 1995, spring chinook salmon required an average of 18 days to migrate through the diversion areas (**TMD** to above Stanfield Dam) compared to 36 days for summer steelhead. Most of the difference in passage time occurred between TMD and **Westland** Dam. Spring chinook salmon required on average six days to complete this section while summer steelhead required 27 days.

Like summer steelhead, it appears that gravel accumulations coupled with increased flows greatly affected spring chinook salmon passage at Feed Canal Dam in 1995. In 1994, average passage time (hours and minutes) for spring chinook salmon at Feed Canal Dam was **11:58**. This number increased to **89:42** in 1995. It's interesting that average passage time for summer steelhead at Feed Canal Dam was nearly identical at **83:24**. During 1994, flows (encountered during passage) at Feed Canal Dam ranged from 346 to 1,563 cfs. In 1995, flows ranged from 689 to 2,772 cfs. During moderate to high flow events, such as those experienced in 1995, much of the flow spilled over the crest of the dam and was directed away from the fish ladder. By itself, false attraction will increase passage times. Compound this with gravel accumulations that prevent migration toward the fish ladder and passage times increase dramatically. This occurred at Feed Canal Dam in 1995. During low flow events, as in 1994, most of the flow was directed toward the irrigation canal headworks and toward the fish ladder. Under these circumstances, ascending adults homed in on the fish ladder and passage times reduced accordingly. This does not suggest that spring chinook were without migrational difficulty at Feed Canal Dam during low flow conditions. Average passage times at Feed Canal Dam were more than **15** times higher than those at Stanfield Dam in 1994, and more than 20 times those at Stanfield Dam in 1995.

OBJECTIVE 4: Adult Passage Evaluations Following Upstream Transport.

Task 4.1: Evaluate movements of radio tagged adult spring chinook salmon and summer steelhead following upstream transport.

Summer Steelhead

A total of 11 summer steelhead were radio tagged between November 10, 1994 and April 7, 1995 as part of the upstream transport evaluation. Following release at either **Barnhart** or Nolin, nine migrated upstream (seven into the Umatilla River, one into Birch Creek, one into McKay Creek), one fell back below TMD and was recaptured and hauled upstream, and one regurgitated the radio transmitter. On average, fish released at TMD traveled at a rate of 4.1 miles/day (5.9 miles/day in 1993-94) between Stanfield Dam and the fixed-site at ODFW (Table F-4). By comparison, fish hauled upstream traveled an average of 1.7 miles/day (5.2 miles/day in 1993-94) between the release site (Barnhart or Nolin) and the ODFW site (Table F-5).

In 1995, ten summer steelhead provided data following upstream transport and release. All but one migrated upstream following release at either **Barnhart** or Nolin. Although similar in 1994, migrational rates through the same section of river for fish released at TMD versus those hauled upstream were different in 1995. Some discrepancy in miles moved per day can be explained by differences in release dates. Variation between years was likely a result of changing flows and water temperatures. Migrational differences in these two release groups was not critical but does provide a means of comparison. What does matter is whether summer steelhead successfully migrate to spawning locations following upstream transport. In the last two years, 94% (17 out of 18) of the summer steelhead evaluated successfully migrated upstream following upstream transport and release.

Spring Chinook Salmon

Beginning on May 16 and concluding on June 16, 1995, a total of ten spring chinook salmon were radio tagged at TMD and released at either Thomhollow (**RM 73.5**) or Imeques **C**-mem-ini-kern (**RM 80**). After release, six remained at or near the release location until time of

spawning, one fell back to **Stanfield** Dam and then returned upstream (above the ODFW site, RM **56**), two fell back to **Westland** Dam and then returned upstream, and one regurgitated the radio transmitter.

Because all spring chinook salmon were released above the uppermost receiver (ODFW site), no 1994-95 migrational comparisons of upstream transport versus passage evaluation are available. Comparisons for 1993-94 and passage evaluation information for 1994-95 is provided in Tables F-6 and F-7.

During the last two years, a total of 18 spring chinook salmon (nine each year) have provided migrational data following upstream transport and release. All 18 have successfully migrated to or remained at spawning locations. Most salmon in 1995 (six out of nine) remained at or near the release location (**Thornhollow**, Imeqes C-mem-ini-kern) until spawning. Three, however, fell back into the diversion sections of the Umatilla River (one to Stanfield Dam and two to **Westland** Dam) before returning upstream. Although some fall-back following release was expected, these fish fell back an average of 46.5 miles. All three fish fell back during late **May** and early June. At this time, flows in the lower section of the river, particularly below the major diversion points, were extremely low and water temperatures were extremely high.

In recent years, adult counts on spawning surveys in relationship to release numbers at TMD have suggested spring chinook salmon are falling back into the lower Umatilla River and potentially out of the basin. As recent as 1993, an estimated 43% of the spring chinook salmon released above TMD were unaccounted for (CTUIR 1994). It's possible that the Umatilla River received strays from other systems. Once released above TMD, they fell back over the dam to continue migration to their stream of origin. To better understand these questions, this project will focus on the movements of spring chinook salmon in 1996.

OBJECTIVE 5: Evaluate Homing and Straying of Adult Salmonids

Task 5.1: Determine factors essential for homing and upstream migration of maturing salmonids.

Fall chinook Salmon and Coho Salmon

Consistent with **mainstem** passage information (Table **F-8**), CWT data demonstrate that Umatilla River fall chinook salmon first enter the John Day Pool during the period of August 24 to 30 with peak migration occurring in mid September (**Kissner** 1992, Wagner 1990). In 1992, significant numbers of fall chinook salmon entered the Umatilla River when flows reached 150 cfs (Figure F-8). Large numbers of fall chinook salmon entered at 200 cfs in 1993 and 1994 (Figures F-9 and F-10).

Homing rates for Umatilla River fall chinook salmon (all release groups) during the last four return years have ranged from a low of 24% in 1992 to a high of 59.5% in 1990 (Table F-9). Average attraction flows exiting the Umatilla River in early September (September 1-15, 1990-94) ranged from a low of 1.5 cfs in 1992 to a high of 78 cfs in 1993 (Table F-9). Acclimated versus direct release experiments of fall chinook salmon (Table F-10) show weighted average homing rates of 52.1% and 55.3 % respectively. Homing rates versus age at release for Umatilla River fall chinook salmon were highest for age 1+ fish. Age 1 + fish had weighted average homing rates of 67.9% while spring and fall releases of subyearlings (O+,O+ +) averaged 48.4% (Tables F-11 and F-12).

Although **coho** salmon enter the Columbia River later than fall chinook salmon, entry timing at TMD was similar. In 1992, **coho** entered TMD when flows reached 150 cfs (Figure F-8). Two-hundred cfs was required to encourage significant numbers in 1993 and 1994 (Figures F-9 and F-10).

Many **coho** salmon released in the Umatilla River return to their rearing facility at Bonneville Complex (Table F-13). Stray rates above McNary Dam were essentially zero. Homing rates for **coho** salmon (all release groups) during the 1987-91 return years have ranged from a high of 100% to a low of 58.3%. Weighted average homing rate for these same years was 73.1% (Table F-13). Weighted average homing rates to the Umatilla River for acclimated versus direct releases of **coho** salmon were 70.4% and **72.1%**, respectively (Table F-14).

Entry for fall chinook salmon at TMD hinges on availability of attraction flows. Phase I of the Umatilla Basin Project provided minimum flow levels below TMD beginning in 1993. These flows, however, have not been significant enough to encourage migrational entry. Data clearly demonstrate that at least 150 cfs was required to encourage movement of both fall chinook and **coho** salmon into the Umatilla River. Without attraction flows from the mouth of the Umatilla River in late August and early September, straying and late entry of fall chinook salmon is inevitable.

Regardless of attraction flow levels, it may be discovered that some fall chinook salmon naturally migrate upstream of the mouth of the Umatilla River. Migrational behavior of this type has been documented for both Umatilla River origin summer steelhead and spring chinook salmon at attraction flows far exceeding those experienced during the fall chinook salmon migration (Volkman 1994). Fall chinook salmon above the mouth of the Umatilla River may simply be “testing” for Umatilla River water with the intention of dropping back if the Umatilla River is not detected. Once over McNary Dam however, they find passage back through the dam **difficult** and thus spend days if not **weeks** in the McNary pool and **forebay** before successfully falling back and entering the Umatilla River. Typically, a Umatilla River origin fall chinook salmon above McNary Dam was considered to be straying. In reality, this may be a natural part of the migrational process of these fish.

It would be interesting to observe entry **dates** of fall chinook salmon at flows exceeding 500 cfs in early September. Given these conditions, **mainstem** straying and thus delay may be significantly reduced. One might argue that historically flows at the mouth of the Umatilla River were not 500 cfs in early September. Historically, however, the Columbia River was not a reservoir as it is today. Lake-like conditions and thus poor water mixing in the **mainstem** may demand attraction flows far greater than previously required. The construction of **mainstem** dams has also made it more difficult for fish to ascend and fall-back to their respective tributaries. At this time, attraction flow levels in the Umatilla River are not fully understood. Until more information is gathered, minimum attraction flows should not be set.

Summer Steelhead

Coded wire tag data analyzed by Kissner (1992), found summer steelhead in the **mainstem** Columbia River (Zone 6) from August 1 through October 31. Entry timing at TMD varies and may extend over ten months. Though large numbers of summer steelhead have entered the Umatilla River in November and December, typically the largest number of **fish** enter in February, March, and April.

In each of the last three return years, peaks of over 500 cfs (over 1,000 cfs in some years) were necessary to encourage significant numbers of summer steelhead to enter TMD (Figures F-11, F-12 and F-13). Water temperatures above **4.4°C (40°F)** generally do not delay entry. Stray

rates for summer steelhead were low. Coded wire tag data analyzed by **Rowan** (1994) uncovered one Umatilla River origin summer steelhead above **McNary** Dam. However, some Umatilla River summer steelhead were known to migrate over **McNary** Dam prior to falling back and ascending the Umatilla River (Wagner 1990, Wagner and **Hillson** 1991).

Entry timing for summer steelhead at TMD can begin as early as late August and extend into late May. Native summer steelhead have survived in the Umatilla River because of their ability to wait long periods of time, if necessary, between **mainstem** entry and spawning (**Kissner** 1992). Stray rates associated with summer steelhead were extremely low. Unlike salmon, summer steelhead migrating above McNary Dam can have as long as ten months to fall-back, relocate, and successfully ascend the Umatilla River.

Large flows were necessary to attract significant numbers of summer steelhead into the Umatilla River. Flows exceeding 500 cfs were required in most cases and as much as 1,500 cfs in some years. This does not suggest migrational entry will not occur at flows less than 500 cfs. Summer steelhead will enter the Umatilla River under low flow conditions, but when available, most enter during moderate to high flows.

Spring Chinook Salmon

Spring chinook salmon migration in the Umatilla River begins in early April and typically peaks in May. Migrational entry of spring chinook salmon versus flows varies greatly year to year (Figures F-14, F-15 and F-16). Migration to TMD will occur at flows ranging from 200 cfs to over 10,000 cfs (Volkman 1994). In both 1993 and 1995, 2,000 cfs was necessary to encourage migration (Volkman 1993). In 1994, 500 cfs was required.

Umatilla River spring chinook salmon stray rates remain low. Coded-wire tag homing data (all release groups) for the recovery years of 1990-94 have ranged from 92.4% in 1994, to 99.9% in 1991 (Table F-15).

Recommendations

Modification of Feed Canal Dam is the highest priority. Telemetry data have identified this dam as the only significant barrier to upstream migrants (from above TMD to above **Stanfield** Dam) under adequate flow conditions. In the absence of modifications at Feed Canal Dam, large delays and impasse will occur. As mentioned previously, additional jump pools and fish ladders may help. The design of this facility, however, encourages false attraction and will likely continue to cause problems. Complete reconstruction or removal of the dam is likely the best option for upstream migrants at this facility.

Plans for the 1995-96 Adult Passage Evaluation

Radio telemetry has provided valuable information regarding the migrational movements of adult salmonids in the Umatilla River. Each year, a better understanding of the movements of anadromous fish is being assembled. For 1995-96, CTUIR will conduct a study similar in size and scope to the study conducted previously. An additional receiver will be installed below TMD. Migrational patterns following release at TMD will be evaluated for all four species of anadromous salmonids in the Umatilla River. Summer steelhead and spring chinook salmon will be evaluated following upstream transport. Greater effort will be designated to increasing the sample size for both evaluations.

OBJECTIVE 6: Spawning Surveys

Task 6.1: Determine the final disposition of adults salmonids released above TMD.

Summer Steelhead

The estimated disposition of 875 natural and 656 hatchery summer steelhead trapped at TMD from September 26, 1994 and June 22, 1995, follows: 86 natural and 68 hatchery adults taken for broodstock; 33 hatchery adults sacrificed for **CWTs**, five natural and 25 hatchery adults harvested by tribal members (Task **8.2**), and 21 hatchery adults harvested by non-tribal anglers (Mike Hayes, ODFW, personal communication). The remaining 784 natural and 509 hatchery adult steelhead were available for spawning. Prior to release at TMD, adult steelhead were marked. Five marked summer steelhead fell back over the dam and were recaptured again.

Spring Chinook Salmon

The disposition of 388 adult and 108 jack spring chinook salmon trapped at TMD from March 29 to June 27, 1995 entails ten adults and 46 jacks sacrificed for **CWTs** and 378 adults and 62 jacks released above TMD for spawning (Table G-5). Prior to release at TMD, adult salmon were marked. Seven marked spring chinook salmon fell back over the dam and were recaptured again.

Fall Chinook and Coho Salmon

At the adult trap at TMD, 688 adult and 604 jack fall chinook and 984 adult and 62 jack **coho** salmon were trapped between August 26 and December 5, 1994. Crews collected **CWTs** from 95 adult and 74 jack fall chinook and 105 adult and eight jack **coho** salmon. The remaining salmon were released above TMD to spawn and included 593 adult and 530 jack fall chinook and 879 adult and 54 jack **coho** salmon.

Tasks 6.2 and 6.3: Conduct prespawning, spawning, and post spawning surveys throughout the basin for each anadromous species and run. Estimate the number of successful redds and the adult/redd ratios (female/redd, female/male) of fish passed above TMD (adjusted for harvest and fall-back, if possible).

Summer Steelhead

During summer steelhead escapement surveys, we observed 35 adults on redds, six adults holding (peak counts) and 87 redds (**3.3/mile**) along 26.5 miles of lateral tributaries of the upper Umatilla River (Table G-1). ODFW conducted escapement surveys on 8.8 miles of Birch Creek tributaries and enumerated 39 redds (**4.4/mile**; Tim Bailey, ODFW, personal communication). Scales were sampled from three carcasses, three adults trapped in the rotary screw trap (**RM 42.2**) and three from the water intake at TMD. Most biological data (age, sex, length and scales) were obtained from the natural brood trapped at TMD and held at Minthom Springs. If desirable, additional adults could be sampled at **Westland** when the Trap and Haul Project operates.

Conditions for surveys were generally excellent in the smaller tributaries from March 8 through April 18. Heavy rains and high water in late April made survey conditions poor through May. A survey of Squaw Creek (May 18) indicated that previously marked redds were no longer visible. Escapement surveys of summer steelhead were terminated for the year.

Summer steelhead redd data can not be utilized as an **annual** index of abundance because conditions for observing the escapement vary too much from year to year. Summer steelhead redds are perhaps the most difficult of *Oncorhynchus* to enumerate because of the variation in the size of spawning fish and the number of false redds. Resident rainbow trout also spawn at the same time and often in similar substrates.

Steelhead escapement surveys in years with low snow pack and low precipitation can yield valuable information. Some trends can be documented for smaller systems and surveys can assist biologists in quantifying fishery values of streams. Single surveys once a year to enumerate steelhead redds were of limited value in the Umatilla River Basin. Detection of redds has been difficult just two weeks after redd construction. Furthermore, substrate movement during freshets can conceal redds. Because of the variables discussed above, and factors such as harvest, there was not a good correlation between summer steelhead released above TMD and redds/mile (Table G-2).

Surveys during low flow years indicate that Meacham Creek and tributaries are probably the most important summer steelhead spawning areas in the Umatilla River Basin followed by Squaw Creek (Table G-3, Figure A-4). Based on CTUIR and ODFW surveys, East Birch Creek and Pearson Creek are also important summer steelhead spawning tributaries.

Spring Chinook Salmon

During spring chinook salmon escapement surveys, we enumerated 90 redds (**1.6/mile**) sampled 217 carcasses along 55.8 miles of the Umatilla River Basin between May 30 and October 2, 1995 (Table G-4, Figure A-4). We recovered 49.3% of the 440 spring chinook salmon released above TMD. A total of 60 **CWTs** were removed from 78 adipose clipped spring chinook salmon found during surveys. Dispositions of spring chinook salmon enumerated at TMD from 1989-95 are presented in Table G-5.

Survival to spawning of spring chinook salmon above Pendleton varied greatly between areas. Survival of adults to spawning was again highest in the colder headwaters and decreased downstream as water temperatures increased. Survival to spawning (based on carcass examination) was 92.9% in the North Fork of the Umatilla River, 81.4% between the Forks and Fred Gray's Bridge (**RM 90-80**), 63.2% from Fred Gray's Bridge to the **Meacham** Creek confluence (**RM 80-79**), and 37.7 % from the confluence of Meacham Creek to Thornhollow Bridge (**RM 79-73.5**) (Tables G-6 and G-1 1). The percentage of the carcasses sampled this year that had successfully spawned was the lowest observed to date, 66.8%. Zimmerman (CTUIR, personal communication) noted that approximately 33% of the spring chinook salmon enumerated at TMD during April through June, 1995, were injured. To assist the rapid development of a naturally sustaining population of spring chinook salmon, adults should be hauled to Corporation (**RM 89**) for the next five years (one cycle). Spring chinook salmon released in the lower river have often failed to migrate to the cold, relative pristine, headwaters. Many chinook died before spawning because of high water temperatures (Brett 1952, Black 1953). Others spawned in locations where survival of their progeny was likely poor because of high incubation temperatures. This has been especially evident in Meacham Creek and the **mainstem** Umatilla River below Meacham Creek. Hauling adults to the headwaters would increase egg deposition into quality habitat. Egg to fry and **fry** to parr survival would improve because of the cooler incubation temperatures and better rearing conditions.

Fall Chinook and Coho Salmon

Adult returns in the fall of 1994 included 711 fall chinook salmon (greater than 610 mm) (688 at TMD and 23 below) and 1,003 **coho** salmon adults (greater than 457 mm; 984 at TMD and 19 below; Table G-7, Figure A-4). Fall chinook and **coho** salmon escapement surveys were conducted from October 27 through December 19, 1994. Eighty-two fall chinook redds, 24 **coho** salmon redds and seven unidentified salmon redds (112 total redds, **2.6/mile**) were enumerated. Forty-nine fall chinook and 41 **coho** salmon carcasses were sampled along 42.3 miles of the **mainstem** Umatilla River above TMD (Table G-8). During past years, the majority of adult fall chinook and **coho** salmon were nearly ripe when captured at TMD. After being hauled to the Yokum or **Barnhart** release sites, most spawned immediately in the general area. The fall of 1994 was the first year significant numbers of adult fall chinook and **coho** salmon were released above TMD well before reaching maturity. The majority of fall chinook and **coho** redds were observed from Mission to Thornhollow Bridge (**RM 60.0-73.5**) with the highest concentration from Mission to **Minthorn Springs (RM 60.0-63.8)**. Fall chinook and **coho** salmon still spawned in the vicinity of Bamhart and Yokum, but water clarity was poor for accurate surveys. Surveys were not conducted from TMD to Echo Bridge (**RM 26.3**) because of poor conditions. Below TMD, redds were not enumerated because of poor water clarity. Twenty-five fall chinook and 19 **coho** salmon carcasses were sampled (Table G-9).

Enumerating adult fall chinook and **coho** salmon redds and carcasses does not provide a good indicator of spawning distribution or success because survey conditions were too poor during late fall. Radio telemetry may be a better tool to determine spawning distribution of fall chinook and **coho** salmon.

Task 6.4: Calculate fecundity of fish found on spawning grounds. Estimate the number of eggs/redd and total eggs deposited by stream reach, stream and drainage.

The potential egg deposition of spring chinook salmon in the Umatilla River (above RM 51) during 1995 was approximately 90 redds x 4,376 (average fecundity, Table **G-10**), minus 3,607 (eggs retained) = 390,233. Based on previous surveys, we assume few spring chinook salmon successfully spawn below the mouth of McKay Creek. Few spring chinook salmon carcasses have been found below RM 51. Furthermore, the potential for natural production of spring chinook salmon in this reach is minimal because of high water temperatures.

Estimates of egg deposition by summer steelhead, fall chinook and **coho** salmon were difficult to calculate because of poor survey conditions during spawning season. However, previous surveys indicated that prespawning mortality for these species has been minimal (CTUIR research records). During the fall of 1994, survival to spawning above TMD was estimated from carcasses at 95.7% for fall chinook and 94.3 % for **coho** salmon. Egg deposition by fall chinook females would be about **1,076,000**, assuming 95.7% spawning success, 301 females above TMD and a mean fecundity of about 3,735 eggs/female. Egg deposition by **coho** would be approximately 884,000 based on 94.3% spawning success, 398 females and a mean fecundity of 2,356 eggs/female.

Steelhead egg deposition of approximately **4,887,000** was derived from 862 females (887 released above the TMD minus 51 adults harvested, with a 50-50 sex ratio) with a mean fecundity/female of 5,669, and assuming survival through spawning near 100%. While this provides an estimate of potential egg deposition, a better measure of reproductive success may be derived from estimating fry abundance the following summer.

Task 6.5: Compare Umatilla Basin spawning survey findings with other salmonid populations in the region if available.

In the Umatilla River redd index area (**RM 78.9 to 89.9**), we observed an average of 5.8 (3.9 to 8.7) spring chinook salmon **redds/mile** during the last five years. In Catherine Creek during the same period, spring chinook redds averaged **8.6/mile** and ranged from 2.0 and 16.5 **redds/mile**. The Upper Grande Ronde index area **redd** counts averaged 3.5 **redds/mile** and varied between 0.4 and 8.6 **redds/mile** from 1991 to 1995. The Imnaha redd index ranged from 2.5 to 27.5 **redds/mile** and averaged 10.8 during the same period. Only spring chinook salmon redd counts could be compared because of inconstant methods and variable survey conditions associated with spawning surveys for fall chinook salmon, **coho** salmon and summer steelhead.

OBJECTIVE 7: Smolt Trapping

Task 7.1: Install and operate rotary screw traps in Umatilla River below the mouth of Squaw Creek (RM 76) and below the mouth of Birch Creek (RM 48).

The rotary screw trap in the Umatilla River at Tumla (**RM 76**) operated 63 of 113 days from September 21, 1994 through January 13, 1995. High flows, ice buildup and damage to the trap prevented continuous operation of the trap at this site. The trap captured 596 juvenile steelhead. Mean trap efficiency rate was 9.9% for juvenile steelhead (51 recaptured from 516 marked and released). A total of 1,368 juvenile chinook salmon were captured. Mean trap efficiency rate was 28.8% for juvenile chinook (347 recaptured out of 1,207 marked and released; Table H-1, Figures H-1 Through H-4). On January 14, 1994, the trap and mooring systems were damaged during high flows and the river channel changed making the Tumla site unsuitable.

The rotary screw trap at the Imeqes C-mem-ini-kern site (**RM 79.5**) operated 43 out of 43 days from May 5 to June 16, 1995, and captured 304 juvenile steelhead. Mean trap efficiency rate was 6.6% for juvenile steelhead (18 recaptured from 273 marked and released). A total of 102 juvenile chinook salmon were captured. Mean trap efficiency rate was 10.5% for juvenile chinook (11 recaptured out of 95 marked and released; Tables H-1). Peak catches of juvenile steelhead and chinook salmon occurred in October, April and May.

The rotary screw trap at the **Barnhart** site (**RM 42.2**) operated 87 out of 125 days from March 3 to June 1, 1995. The trap captured 105 natural juvenile steelhead, 247 natural juvenile chinook salmon, five natural **coho**, 6,265 hatchery juvenile chinook salmon, 467 hatchery steelhead and 16,844 hatchery **coho**. Mean trap efficiency rates for salmonids ranged from 2.3% to 5.7% (Table H-1).

Several uncertainties affect the evaluation of trap data regarding naturally produced smolts emigrating from the basin. These uncertainties include large day to day variation in trap catch rates, lack of recaptures, low catch, winter mortality of fish moving past the trap in the fall before they leave the basin in the spring, the unknown number of salmonids passing the trap during the days the traps were not operated and the unknown proportion of the steelhead captured that were resident trout.

Nineteen bull trout were captured in the traps from October 4, 1994 to June 5, 1995 (Table I-5). In comparison, 139 bull trout were trapped during the previous season (fall of 1993 and the spring of 1994). This was likely because of trapping at RM 76 during the fall of 1994 as apposed to RM 79.5 during the fall of 1993 (Table I-5). The 15 bull trout trapped in October and November, 1994, averaged 279 mm (fork length; SD 50.3 n= 15) in contrast to the four trapped in

May and June, 1995, which averaged 152 mm (SD 12.9). The trend of larger fish being captured in the fall was similar during the previous two years.

Task 7.2: Install and operate modified pipe traps in Birch Creek.

The pipe traps were not installed or operated in Birch Creek.

Task 7.3: Estimate trap efficiencies.

See Task 7.1.

Task 7.4: Freeze brand fish for interrogation in the lower Umatilla And Columbia Rivers in coordination and cooperation with ODFW and the Fish Passage Center.

Freeze branding was postponed until the fall of 1995.

Task 7.5: Reconstruct emigration timing and minimum survival rates.

Emigration from the headwaters (past RM 79.5) by juvenile steelhead and chinook salmon during the last two years peaked in October and again during April and May (Figures H-5 through H-10, CTUIR 1994, Contor et al. 1995). Fish continue to move downstream throughout late fall and winter at lower rates. Apparently, portions of the population move out of the headwaters in the fall to utilize habitat made available as water temperatures drop below **20°C** (68°F). Considerably more juveniles (11,035 to 1,093) were estimated to have emigrated past Tumla in the fall than past Imeques C-mem-ini-kern in the spring. This disparity was only partly explained by the difference in trapping duration in the fall and the exclusion of Meacham Creek migrants in the spring. Peak migration during the fall from the headwaters was consistent with the previous trapping season in the Umatilla River (Contor et al. 1995) and in Lookingglass Creek (**Lofy and McLean 1995a, 1995b**). Chinook captured in the fall at Tumla (**RM 76**) averaged 20 mm longer than those captured in the spring at Imeques C-mem-ini-kern (**RM 79.5**; Figure H-1). During the fall, chinook lengths at Tumla were similar to those captured at **Barnhart (RM 42.2)** in the spring. Survival rates were not estimated because Task 7.4 was postponed.

Task 7.6: Design and conduct an eight month mark retention study.

The mark retention study was postponed until 1995-96.

OBJECTIVE 8: Tribal Harvest

Tasks 8.1 and 8.2: Design and implement creel and phone surveys to estimate tribal harvest of adult anadromous salmon.

Tribal steelhead angling in the Umatilla River was monitored 550 hours during 44 days from December, 1994 through April, 1995. Thirty-five tribal anglers were interviewed one or more times either while fishing or during telephone interviews. Thirty adult steelhead were estimated to have been harvested (25 hatchery and **five** natural) by tribal anglers. They reported

catching and releasing another 12 steelhead. Reported catch rates for tribal anglers ranged from 80 hours/fish to 7.5 hours/fish. Mike Hayes (ODFW, personal communication) estimated **non-tribal** anglers harvest an additional 21 steelhead (below the reservation boundary). There was no tribal season on spring chinook salmon during 1995. Harvest of fall chinook and **coho** salmon was minimal as very little angling effort was observed as a result of poor returns.

OBJECTIVE 9: Age and Growth

Tasks 9.1 and 9.2: Age analysis of adult and juvenile salmonids.

Based on scale analysis, 46.4% of Umatilla River natural adult summer steelhead returning to spawn in 1995 were from the 1990 brood year, 33.9% were from the 1991 brood year, and 19.6% were from the 1989 brood (Tables I-1 and I-2). Sixty-four percent of the steelhead sampled reared for two years in fresh water before emigrating while 36% reared three years (Table I-3).

During 1995, we collected and **aged** scales from 448 natural juvenile steelhead from Coonskin, Moonshine, Cottonwood, and Mission Creeks, and the Umatilla River (**RM** 81.8-89.6). An additional 303 scale samples were collected during index surveys.

Juvenile steelhead were the most abundant **salmonid** captured during biological surveys. From 87.7 to 96.2% of steelhead sampled were 0+ or **1+** while 3.8% to 12.3% were age **2+** or **3+**. Only one **4+** fish was sampled. Age structure of steelhead sampled in 1995 was similar to 1993 and 1994 findings (CTUIR 1994, Contor et al. 1995). Mean length, range and standard deviation by age class of sampled juvenile steelhead, and an expansion of age classes (by length) for all steelhead are presented in Table I-4. Age structure of 272 steelhead collected from index sites was 26.6% 0+, 48.5% **1+**, 22.8% **2+**, 1.5% **3+** and 0.7% **4+**. Scales from spring chinook carcasses indicated that 91.4% of adults returning in 1995 were from the 1991 brood and 8.6% were from the 1990 brood.

Attempts were made to separate hatchery and natural spring chinook salmon adults by examination of freshwater growth, circuli counts to the first (freshwater) **annulus**. A total of 20 scale samples of adipose clipped and coded wire tagged adult spring chinook salmon were compared with 20 scale samples of unmarked adult returners.

Most freshwater circuli counts from hatchery spring chinook salmon ranged from **20-40** while most unmarked salmon ranged below 16. However, 40% of the freshwater circuli counts from CWT spring chinook salmon released during November in 1992 (1991 Bonneville brood) overlapped with circuli counts from unmarked salmon. Since 100% of salmon from the 1991 Bonneville brood were not marked, we could not use circuli counts to determine the origin of the unmarked salmon.

Limited scale analysis indicated that most bull trout were age three and four years old (**2+** and **3+**, Table I-5). Ten bull trout (165 to 290 mm) were age three and six were age four (225 and 320 mm). Scales patterns indicated that growth was slow during the first two years and then increased rapidly. Most of the bull trout captured in the rotary trap at RM 79.5 have been captured in late October and November. Many had crooked but **healed** lower **caudal** fin rays, indicating that they apparently spawned at least once. None of the bull trout observed or sampled during the fall at the rotary screw trap were sexually mature.

OBJECTIVE 10: Genetic and Ecological Effects of Supplementation

Task 10.1: Establish a genetic baseline database from native steelhead.

This work was conducted and reported by Currens and Schreck (1993, 1995). Their efforts provided a genetic baseline for future comparisons.

Task 10.2: Review literature on effects of hatchery-reared salmonids on naturally produced salmonids.

The primary goal of “supplementation” as applied to steelhead in the Umatilla River Basin Restoration Project was to increase natural production and produce surplus adults for harvest (CTUIR 1984, ODFW 1986). The effects of releasing hatchery reared salmonids sympatric to wild and natural **salmonid** populations has been explored from a variety of perspectives. Strategies to examine this topic have ranged from monitoring genetic heterozygosity and the persistence of unique alleles to evaluating the performance of hatchery and wild salmonids spawning naturally. Some researchers have suggested that hatchery programs may decrease the production of natural salmonids (Nickelson et al. 1986, Vincent 1987, Leider et al. 1990, Flemming and Gross 1991). Others have advised using supplementation to restore and enhance natural populations (CTUIR 1984, ODFW 1986, Bowles and Leitzinger 1991).

The effects of supplementation on the genetics of natural populations has been of prime concern in the fisheries literature (Reisenbichler and Phelps 1989, Meffe 1992, Steward and Bjornn 1990). Research in stock genetics has demonstrated that hatchery spawning practices can have a variety of effects on population genetics. Allendorf and Phelps (1980) found hatchery cutthroat trout (*Oncorhynchus clarki*) had lost genetic variation over time. Reisenbichler and Phelps (1989) found significant genetic differences between hatchery and wild steelhead in northwest Washington. They attributed these genetic differences to hatchery broodstock selection and spawning practices. Ferguson et al. (1991) found ancestral and descendent rainbow trout had no significantly different allelic frequencies when modern breeding techniques were practiced. Byrne et. al (1992) modeled the genetics of steelhead supplementation strategies using an equally fit broodstock with different alleles. He demonstrated that often “supplementation of native stocks with hatchery fish caused replacement, not enhancement of native fish.” Byrne’s et. al (1992) and Meffe (1992) both emphasized that to enhance natural steelhead, carrying capacity of the rearing and migratory habitat must be restored and maintained.

The Umatilla hatchery program minimizes genetic risks by breeding primarily endemic, naturally produced steelhead with modern techniques (matrix spawning). Currently, we estimate there are few risks to the genetic integrity of the natural steelhead population.

Supplementation may impact survival, growth and behavior of natural salmonids through predation, competition, disease transmission, and behavior modification. Predation on natural salmonids by hatchery juveniles occurs when larger sized hatchery smolts are introduced in systems with natural **salmonid** fry and parr. Predation by hatchery fish on wild fry has been documented, however researchers report that hatchery steelhead smolts prey primarily on macroinvertebrates (Parkinson et al. 1989, **Hillman** and **Mullan** 1989, Steward and Bjornn 1990, Cannamela 1992). However, Horner (1978) found some hatchery steelhead became highly piscivorous with salmonids comprising 50% of their diets. Cannamela (1993) examined the stomachs of 6,700 hatchery steelhead smolts for predation on naturally produced chinook fry. Cannamela estimated hatchery smolts preyed on chinook fry at low rates (0.00148 **fry/smolt**).

However even at the low rates, 24,000 fry were estimated to have been eaten in 1992 by 744,000 hatchery steelhead smolts released into Idaho's upper Salmon River.

Competition and displacement occurs when individuals compete for limited resources (Chapman 1966, Everest and Chapman 1972). Evidence for increased competition of food and space was minimal in the Umatilla Basin. Hatchery releases generally occur during moderately high flows when space and food do not appear limiting. Furthermore, hatchery salmonids released into the Umatilla River begin their down stream migration directly after release. During electrofishing surveys (1993-95), few residual hatchery fish have been captured. Boston Canyon Creek, near the Bonifer Acclimation Facility was an exception. We estimated 1,100 hatchery steelhead residualized there in 1993. Natural steelhead over 75 mm appeared to have been displaced by hatchery steelhead. Researchers report that most residuals remain near the point of release (Cannamela 1992, 1993, Hillman and Mullan 1989). Hatchery residuals in the Umatilla Basin exhibit the same behavior. We estimated that approximately 4,000 hatchery steelhead residualize each year in Boston Canyon Creek, Meacham Creek, Minthorn Springs Creek and in the mainstem Umatilla River (Appendix E, CTUIR 1994, Contor et. al 1995). This was a residualization rate of 2.7% and represents 0.6% of the total juvenile steelhead in the basin. Residualization rates in the Umatilla were similar to Viola and Schuck's (1991) findings in southeast Washington (9.9% in early summer to 0.8% in October).

Hillman and Mullan (1989) observed altered behavior of natural chinook fry in the presence of hatchery reared chinook. Natural chinook fry not subject to the hatchery releases showed no change in behavior. However, natural chinook fry behavior did not change when hatchery steelhead were released. Vincent (1987) demonstrated dramatic increases of natural brown trout (*Salmo trutta*) and rainbow trout populations once stocking hatchery rainbow trout ceased. Vincent reported that stocking increased the natural mortality rates of wild trout. Bachman (1984) observed frequent and long antagonistic encounters between hatchery reared trout and wild trout which often resulted in exhaustion of the wild trout and disruption of the stable social structure. Poor survival, excessive activity and energy expenditure for "unnecessary aggressive behavior" by hatchery trout was also reported by Mesa (1991). Except for limited effects at the highest stocking rates, Petrosky and Bjornn (1988) found that stocking rainbow trout did not change the abundance, survival and growth of wild rainbow and cutthroat trout. Competition, predation and behavioral affects on natural salmonids from hatchery releases were estimated to be low in the Umatilla Basin. We estimated that effects were low because management limited the duration of temporal and spacial overlap of hatchery and naturally produced salmonids. Furthermore, the overlap does not appear to occur during summer low flow periods when food and space appear most limiting.

Task 10.3: Identify acceptable levels of impact from steelhead supplementation on natural steelhead and native trout.

Preliminary levels of acceptable impact from supplementation were determined and include the following: 1) small genetic changes are acceptable if they are near the scale of background genetic drift; acceptable levels would be near Nei's genetic differences of 0.02 (Nei and Roychoudhury 1974) and nucleotide diversity of 0.0003 as these levels would be impossible to differentiate from background noise currently found during two years of sampling (Currens and Schreck 1995); 2) residualization rate of five percent or less, and 3) a 10% decline in the number of natural spawners. Approximately 100 natural adults (5-10% of the run) are currently taken for artificial production each year. During poor return years, we supplement the natural brood stock

with hatchery adults (Rowan 1995). Management has defined the acceptable reduction of natural adults, by practice, at approximately 5-10% of the run. To date, no evidence exists that shows supplementation has significantly changed the number of returning natural adults. The relationship between adult returns and flows two years earlier has remained consistent since substantial supplementation efforts began in the mid 1980s (Figure B-1 and B-2). Supplementation was expected to increase the natural returns. While an increase in natural adult steelhead was not evident, neither was there a marked decrease. Our findings in the Umatilla Basin appear to concur with carrying capacity theory and with Byrne's (et al, 1992) and Bowles and Leitzinger's (1991) suggestions that natural rearing and migrational habitat must be restored and maintained to increase natural production.

Tasks 10.4 and 10.5: Examine the utility and feasibility of observing behavior and densities of naturally produced salmonids in treatment and control areas before and after releases of hatchery smolts, and the extent of residualization of hatchery smolts and the effects on naturally produced salmonids.

The options of conducting residualization studies and monitoring behavioral responses of naturally produced salmonids to hatchery releases were examined and found to be feasible but of lower priority. Electrofishing data indicate that most hatchery fish move out of the summer rearing areas soon after release (Appendix E, CTUIR 1994, Contor et al. 1995). Based on the research findings and as discussed above in Tasks 10.1-10.3, managers and researchers on the UMEOC did not recommend conducting steelhead behavior or residualization studies at this time.

OBJECTIVE 11: Supplementation Effects on Natural Steelhead

Task 11.1: Combine, examine and summarize data gathered in objectives 1-10 that would indicate enhancement of natural steelhead through hatchery supplementation.

Production and release of hatchery steelhead in the Umatilla River Basin from 1981 to 1991 has returned 3,306 adult hatchery steelhead to TMD (as of June, 1995). From 1981 to 1990, 1,174 naturally produced adult steelhead were taken for hatchery broodstock. We estimate that 2,844 natural steelhead would have been produced from those adults. To date, supplementation has returned approximately 462 additional adult steelhead to TMD (Table H-2). Assuming hatchery steelhead spawn and produce natural progeny equally as well as natural steelhead, the supplementation project would be considered marginally successful. There was some doubt that hatchery steelhead can naturally reproduce at the same rate as natural steelhead. Chilcote et al. (1986) and Campton et al. (1991) concluded that hatchery steelhead reproduced at 28% and 15% the rate of natural steelhead, respectively. Leider et al. (1990) found that the progeny of hatchery steelhead did not survive as well as progeny from natural steelhead. Nickelson et al. (1986) found that supplementing hatchery coho salmon reduced the number of wild coho juveniles but did not increase the number of adult returns. We speculate that Umatilla River hatchery adults reproduce at higher rates than Campton's et al. (1991) estimates because Umatilla steelhead are progeny of natural steelhead bred with modern techniques. However, we have no data to confirm this supposition.

The benefits to natural steelhead from supplementation appear to be limited at this time, probably because hatchery steelhead have not returned favorably. Smolt to adult survival estimates of hatchery steelhead (1987 to 1991 brood) ranged from 0.02 to 0.94% with a mean of 0.39%

(Rowan, CTUIR, personal communication). Since 1991, smolt quality and down stream passage has greatly improved and subsequent adult returns are expected to reflect these advancements. However, there remains a distinct probability that at least as many natural adult steelhead would have been produced without supplementation efforts. As Byrne (et al. 1992) suggests, supplementation may replace natural steelhead with hatchery steelhead. This would be expected if Chilcote's et al. (1986) and Campton's et al. (1991) findings hold true for Umatilla River hatchery steelhead spawning success.

We also explored carrying capacity theory in relation to the effects of supplementation on the natural production of steelhead. Adult steelhead taken from the natural spawning population for broodstock may have been surplus. Under this scenario, their loss did not affect natural production because carrying capacity in the Umatilla Basin had already been reached (under current habitat conditions). Some evidence of a carrying capacity has been found and was summarized in Appendix E and reported in previous progress reports (CTUIR 1994, Contor et al. 1995). Densities of juvenile steelhead were often as high as 100 **fish/100 m²** and have been as high as 222 **fish/100 m²**. Areas surveyed with few or no steelhead had poor environmental conditions. Additional steelhead produced through supplementation efforts would probably not have survived in the poor habitat any better than existing steelhead. Therefore, no net increase in natural production would be expected. Furthermore, the flow/steelhead relationships plotted in Figures B-1 and B-2 indicate that additional spawners may not produce more adults unless rearing and passage conditions improve. The fact that high steelhead densities exist in even moderately suitable habitat throughout the Umatilla Basin suggests that habitat may already be fully seeded. Under a fully seeded scenario, supplementation designed to increase natural production would have marginal success and would simply replace natural steelhead with steelhead of hatchery origin (Byrne et al. 1992). Supplementation has produced hatchery steelhead for harvest and allowed natural fish to become protected under catch and release regulations. Aggressive habitat improvement projects (past, present and future) are expected to increase suitable habitat throughout the Umatilla River Basin. In summary, available data (through 1995) does not indicate that steelhead supplementation has reduced the number of natural adult steelhead spawning in the Umatilla Basin.

Task 11.2: Examine potential tests to better evaluate supplementation.

Managers expect positive results from supplementation efforts and would like to document results for effective evaluation. Identifying levels of acceptable risk and negative impacts requires adequate measurement. However, researchers and managers concur that it is difficult to develop reliable methods to measure supplementation effects. Setting up replicate tests with effective experimental controls in the field is challenging. Furthermore, moderate effects of supplementation may be difficult to separate from effects of environmental stochasticity.

A management paradox may evolve if natural populations begin to decline. Increased supplementation would probably be implemented to "rescue" the natural runs. However, without a good measurement of supplementation effects, there remains a probability that supplementation replaces natural steelhead with hatchery steelhead as predicted by Byrne (et al. 1992). Increased supplementation could either solve the problem or magnify it.

Managers need reliable measurements of supplementation's effect on natural steelhead. Several strategies were examined that would assist in monitoring and evaluating the effects of supplementation on natural steelhead. Several of these strategies are being implemented and include monitoring genetic and phenotypic variation, adult returns, smolt production and smolt to

adult survival. However, the complicated effects of multiple environmental factors could mask effects of supplementation.

Additional strategies include tests with controls and treatments. Weirs could be used to control the number and type of adults allowed to attempt spawning in Meacham Creek (supplementation) and Birch Creek (natural). However, weirs are expensive, sometimes ineffective at high flows, and may impede or prevent beneficial (natural) movements of salmonids between subpopulations.

A new technique to mark steelhead progeny may be available soon. Unique, benign, biologically compatible compounds would be used as artificial markers of female spawner's progeny. The process would be similar to Rieman's work (Bruce Rieman, USFS, personal communication) with natural levels of selenium. Based on selenium concentrations in otoliths, he was able to determine if juvenile sockeye salmon in **Redfish** Lake, Idaho, were progeny from resident or anadromous female parents. For supplementation evaluations, a compound would be injected into adult hatchery females collected at TMD. The compound would bio-transfer to the gametes before the female spawned naturally in the wild. The indicator would be permanently incorporated into the progeny's otolith. Each progeny would retain the mark throughout life. The proportion of the naturally produced steelhead with this mark would indicate the level of success from supplementation efforts (adjusted by on marking and retention rates). Approximately **200** adults could be sampled each year from brood stock, from carcasses found during spawning surveys and from spawned out adults collected at TMD and **Westland** Dam. Juveniles collected at downstream migrant traps could also be sampled. While the technique has been met with optimistic expectations when discussed with researchers throughout the region, no compound or delivery technique has been developed and tested. CTUIR and UMEOC will continue to discuss and coordinate various approaches and techniques to evaluate supplementation.

ACKNOWLEDGEMENTS

This project was funded by Bonneville Power Administration to mitigate, in part, for salmon and steelhead losses associated with the construction and operation of Columbia and Snake River Dams. The Confederated Tribes of the Umatilla Indian Reservation thank Jay Marcotte, Jerry Bauer and other BPA personnel for their contributions.

The authors thank the following CTUIR employees for their contributions to the project: Aaron Jackson for deployment and recovery of thermographs, compilation of temperature, habitat and biological data and production of many of the tables and figures; Brent Spencer for assisting in all field related aspects of the radio telemetry project; Dave Wolf for creel surveys; Larry Cowapoo, Brian Conner, Vern Spencer and Louis Case for assistance at TMD; Gerald **Rowan**, Brian Zimmerman, Peter Lofy, Mike McLean and Todd Shaw for project experimental design and manuscript review, and Billy Quaempts for the designing and producing the Umatilla River Basin Map. We would also like to thank Gary James, fisheries program manager, for project management, assistance and manuscript review; Joe Richards and Michelle Thompson for contract administration; Julie Burke for managing the Fisheries Office, and Celeste Reves for secretarial assistance and help in report preparation. We specifically wish to thank the following for operating the traps and collecting biological, habitat, and spawning survey data: Melvin **Farrow**, Gene Shippentower, Darryl Thompson, James Marsh, Troy Rodriguez, Brian Thompson, Aaron Jackson, Brent Spencer, and Brian Conner.

We appreciate the help and cooperation of Ron Morris, Bureau of Reclamation, Josephine Miller of the U.S. Geological Survey and the following Oregon Department of Fish and Wildlife personnel: Tim Bailey; John Germond; Suzanne Knapp; William Cameron; Mike Hayes, **Shannon Fochoer**; Kim Jones; Chris Kern; Brad Lovatt; Kelley Moore, and **Stacey** Shapleigh.

The authors appreciate the efforts of all USBR employees involved with the operation and maintenance of Three Mile Falls Dam, **Westland** Dam, Feed Canal Dam and Stanfield Dam.

LITERATURE CITED

- Allendorf, F. W., Phelps, S. R. 1980. Loss of genetic variation in a hatchery stock of cutthroat trout. *Transactions of the American Fisheries Society* **109:537-543**.
- Angradi, T., Contor, C. R. 1989. Henry's Fork fisheries investigations. Idaho Department of Fish and Game, Idaho State University, University of Idaho, The Henry's Fork Foundation and The Idaho Cooperative Fish and Wildlife Research Unit. 96 pp.
- Bachman, R. A.** 1984. Foraging behavior of free ranging wild and hatchery brown trout, *Salmo trutta*, in a stream. *Transactions of the American Fisheries Society* 113: 1-32.
- Black, E. C. 1953. Upper lethal temperatures of some British Columbia freshwater fishes, J. Fish. Res. Board of Canada **10(4):196-210**.
- Bowles, Ed; Leitzinger, Eric. 1991. Salmon supplementation studies in Idaho rivers. Report Submitted to U.S. Department of Energy, Bonneville Power Administration. DOW/BP-01466-1. 167 pp.
- Brett, J. R. 1952. Temperature tolerance in young Pacific salmon, genus *Oncorhynchus*. *Journal of the Fisheries Research Board of Canada* **9(6):265-323**.
- Bugert, R., Busack, C., Mendel, G., Ross, L., Petersen, K., **Marbach, D.**, Dedloff, **J.** 1990. Lower Snake River Compensation Plan, Tucannon River spring chinook salmon hatchery evaluation program. 1990 annual report. U.S. Fish and Wildlife Service, Lower Snake River Compensation Plan. 81 pp.
- Bureau of Reclamation. 1988. Umatilla Basin Project, Oregon, planning report-final environmental impact statement. Pacific Northwest Region. Bureau of Reclamation, Department of Interior, Boise, Idaho.
- Byrne, Alan; **Bjornn, T. C.**, McIntyre, J. D. 1992. Modeling the response of native steelhead to hatchery supplementation programs in an Idaho River. *North American Journal of Fisheries Management* **12:62-78**.
- Campton, D. E.**, Allendorf, F. W., Behnke, R. J., Utter, F.M. 1991. Reproductive success of hatchery and wild steelhead. *Transactions of the American Fisheries Society* **120:816-822**.
- Cannamela, David A. 1992. Potential impacts of release of hatchery steelhead trout "smolts" on wild and natural juvenile chinook and sockeye salmon. Idaho Department of Fish and *Game*. 23 pp.
- Cannamela, David A. 1993. Hatchery steelhead smolt predation of wild and natural juvenile chinook salmon fry in the upper Salmon River, Idaho. Idaho Department of Fish and *Game*. 36 pp.


- Chapman, D. W. 1966. Food and space as regulators of **salmonid** populations in streams. *American Naturalist*. **100:345-357.** .
- Chilcote, Mark W., Leider, Steven A., Loch, John J. 1986. Differential reproductive success of hatchery and wild summer-run steelhead under natural conditions. *Transactions of the American Fisheries Society* **115:726-735.**
- Confederated Tribes of the Umatilla Indian Reservation (CTUIR). 1994. Umatilla Basin natural production monitoring and evaluation **annual** progress report, 1992-93. Report submitted to Bonneville Power Administration, Project No. 90-005-01. 49 pp.
- Contor, Craig R., and Griffith, John S. 1995. Nocturnal emergence of juvenile rainbow trout from winter concealment relative to light intensity. *Hydrobiologia* **299:179-183.**
- Contor, Craig R., Hoverson, Eric., Kissner, Paul., Volkman, Jed. 1995. Umatilla Basin natural production monitoring and evaluation, **Annual** Progress Report 1993-1994. Report submitted to Bonneville Power Administration, Project No. **90-005-01**. 43 pp.
- Currens**, Kenneth, P., Schreck, Carl B. 1993. Genetic analysis of Umatilla River rainbow trout. Oregon Cooperative Fishery Research Unit, Department of Fisheries and Wildlife, Oregon State University, Corvallis, Oregon 97331-3803.
- Currens, Kenneth, P., Schreck, Carl B. 1995. Final Report, genetic analysis of Umatilla River rainbow trout. Oregon Cooperative Fishery Research Unit, Department of Fisheries and Wildlife, Oregon State University, Corvallis, Oregon 97331-3803.
- Everest, F. H., Chapman, D. W. 1972. Habitat selection and spatial interaction by juvenile chinook salmon and steelhead trout in two Idaho streams. *Journal of the Fisheries Research Board of Canada* **29:91-100.**
- Ferguson, M. M., Ihssen, P. E., Hynes, J. D. 1991. Are cultured stocks of brown trout (*Salmo trutta*) and rainbow trout (*Oncorhynchus mykiss*) genetically similar to their source populations? *Canadian Journal of Fisheries and Aquatic Sciences* 48: 118-123.
- Flemming, Ian A., Gross, Mart R. 1991. Reproductive behavior of hatchery and wild **coho** salmon (*Oncorhynchus kisutch*): does it differ? *Aquaculture*, 103: 101-121.
- Hillman** T. W., **Mullan**, J. W. 1989. The effect of hatchery releases on the abundance and behavior of the wild juvenile salmonids. in *Summer and Winter Ecology of Juvenile Chinook Salmon and Steelhead Trout in the Wenatchee River, Washington*. Final Report to Chelan Public Utility District, Don Chapman Consultants, Inc. Boise, Idaho. 379 pp.
- Homer, N. J. 1978. Survival, densities, and behavior of **salmonid** fry in streams in relation to fish predation. Masters Thesis, University of Idaho, Moscow.

- Hubbard L. E., Herrett, T. A., Kraus, R. L., Ruppert, G. P., Courts, M. L. 1995. Water resources data, Oregon, water year 1994. US Geological Survey, Water-Data Report **OR-94-1**, U.S. Department of the Interior. 473 pp.
- Kissner, Paul D. 1992. Evaluation of adult **salmonid** passage in the lower Umatilla River. pp. 49-97 in Knapp, M. S., editor. Evaluation of juvenile fish bypass and adult fish passage facilities at water diversions in the Umatilla River. Report submitted to U.S. Department of Energy, Bonneville Power Administration, **DOW/BO-01385-3**. 107 pp.
- Kucera, Paul A., Banach, Michael J., **Blenden**, Michael L. 1991. Nez Perce Tribe LSRCP evaluation studies. **Annual** Project Report 1989. Nez Perce Tribe, Lapwai, Idaho 57 pp.
- Leider, S. A., Hulett, P. L., Loch, J. J., Chilcote, M. W. 1990. Electrophoretic comparison of the reproductive success of naturally spawning transplanted and wild steelhead through the returning adult stage. *Aquaculture* **88:239-252** (from Byrne et al. 1992).
- Lofy**, Peter T., McLean, Michael L. 1995a. Evaluation of reestablishing natural production of spring chinook salmon in Lookingglass Creek, Oregon, using a non-endemic hatchery stock. Lower Snake River Compensation Plan (1993 Progress Report). U.S. Fish and Wildlife Service Report, **AFF1/LSR 95-02**.
- Lofy, Peter T., McLean, Michael L. 1995b. Evaluation of reestablishing natural production of spring chinook salmon in Lookingglass Creek, Oregon, using a non-endemic hatchery stock. Lower Snake River Compensation Plan (1994 Progress Report). U.S. Fish and Wildlife Service Report, **AFF1/LSR 95-02**.
- Malvestuto, Stephen P., Davies, William D., Shelton, William L. 1978. An evaluation of the roving creel survey with non-uniform probability sampling. *Transactions of the American Fisheries Society*. **107(2):255-262**.
- Malvestuto, Stephen P. 1983. Sampling the recreational fishery. in Larry A. Nielson and David L. Johnson. Editors. *Fisheries Techniques*. American Fisheries Society, Bethesda, Maryland. 468 pp.
- Meffe, Gary K. 1992. Techno-arrogance and halfway technologies: salmon hatcheries on the pacific coast of north **america**. *Conservation Biology*, Volume 6, **3:350-354**.
- Mesa, Matthew G. 1991. Variation in feeding, aggression, and position choice between hatchery and wild cutthroat trout in an artificial stream. *Transactions of the American Fisheries Society* **120:723-727**.
- Moore, Kelly M. S., Jones, Kim K., Dambacher, Jeffrey M. 1993. Methods for stream habitat surveys: Oregon Department of Fish and Wildlife, Aquatic Inventory Project. Corvallis OR, 97330.

- Mullan**, James W., Williams, Kenneth, R., Rhodus, G., **Hillman**, Tracy W., McIntyre, John D. 1992. Production and habitat of salmonids in mid-Columbia River tributary streams. U.S. Fish and Wildlife Service, Monograph I. U.S. Government Printing Office 1993-695-600. 489 pp.
- Nei, M., Roychoudhury, A. 1974. Sampling variances of heterozygosity and genetic distance. *Genetics* **76:379-390**. (from Allendorf and Phelps 1980)
- Nickelson, T. E., Solazzi, M. F, Johnson, S. L. 1986. Use of hatchery **coho** salmon (*Oncorhynchus kisutch*) presmolts to rebuild wild populations in Oregon coastal streams. *Canadian Journal of Fisheries and Aquatic Sciences* **43:2443-2449**. (from Byrne et al. 1992).
- Nielson, Reed S. 1950. Survey of the Columbia River and its tributaries. Report prepared for the United States Department of the Interior, Report No. 38.
- Northwest Power Planning Council (NPPC). 1987. Columbia river basin fish and wildlife program. Northwest Power Planning Council, 850 S.W. Broadway, Suite 1100, Portland, Oregon 97205.
- Northwest Power Planning Council (NPPC). 1990. Umatilla River **subbasin** salmon and steelhead production plan. Northwest Power Planning Council and the Agencies of the Indian Tribes of the Columbia Basin Fish and Wildlife Authority.
- Oregon Department of Fish and Wildlife (ODFW). 1986. A comprehensive plan for rehabilitation of anadromous fish stocks in the Umatilla River **Basin**. Report Submitted to Bonneville Power Administration, Project No. **84-10**.
- Oregon Department of Fish and Wildlife (ODFW). 1987. United States vs. Oregon **subbasin** production reports. Portland, Oregon.
- Petrosky, C. E., Bjorn, T. C. 1988. Response of wild rainbow trout (*Salmo gairdneri*) and cutthroat trout (*S. clarki*) to stocked rainbow trout in fertile and infertile streams. *Canadian Journal of Fisheries and Aquatic Sciences* **45:2087-2105**.
- Petrosky, C. E., Holubetz, T. B. 1987. Evaluation and monitoring of **idaho** habitat enhancement and anadromous fish natural production. **Annual** Report Submitted to U.S. Department of Energy, Bonneville Power Administration, Report DOE/BP-1 338 1-3. 172 pp.
- Reisenbichler, R. R., Phelps, S. R. 1989. Genetic variation in steelhead (*Salmo gairdneri*) from the north coast of Washington. *Canadian Journal of Fisheries and Aquatic Sciences* 46:66-73.
- Rowan**, Gerald D. 1991. Minthom Springs Creek summer juvenile release and adult collecting facility, annual report 1990. Confederated Tribes of the Umatilla Indian Reservation. Report Submitted to U.S. Department of Energy, Bonneville Power Administration. **DOE/BP-17622-6**. 72 pp.

- Rowan**, Gerald D. 1994. Minthom Springs Creek summer juvenile release and adult collecting facility, annual report 1993. Confederated Tribes of the Umatilla Indian Reservation. Report Submitted to U.S. Department of Energy; Bonneville Power Administration. DOE/BP-17622-8. 49 pp.
- Rowan**, Gerald D. 1995. Minthom Springs Creek summer juvenile release and adult collecting facility, annual report 1993. Confederated Tribes of the Umatilla Indian Reservation. Report Submitted to U.S. Department of Energy, Bonneville Power Administration. DOE/BP-17622-9. 49 pp.
- Steward, C. R., Bjornn, T. C. 1990. Supplementation of salmon and steelhead stocks with hatchery fish: a synthesis of published literature. 126 pp. in Analysis of salmon and steelhead supplementation. Miller, W. H. editor. U.S. Department of Energy, Bonneville Power Administration. DOE/BP-92663-1.
- SYSTAT. 1992. SYSTAT Inc. Evanston, IL.
- Taylor, George H. 1993. Normal annual precipitation, state of Oregon, Period 1961-1990. Map. Oregon Climate Service, 326 Strand Ag. Hall, Oregon State University, Corvallis, Oregon.
- U.S. Geological Survey (USGS). 1989. Hydrological unit map, State of Oregon. U.S. Geological Survey, **Reston**, Virginia.
- Van Deventer, John S. Platts, William S. 1989. Microcomputer software system for generating population statistics from electrofishing data- User's Guide for **MicroFish** 3.0. USDA. U.S. Forest Service, Intermountain Research Station. General Technical Report INT-254 **29 pp.**
- Vincent, E. Richard. 1987. Effects of stocking catchable-size hatchery rainbow trout on two wild trout species in the Madison River and O'Dell Creek, Montana. North American Journal of Fisheries Management **7:91-105.**
- Viola, Arthur E., **Schuck**, Mark L. 1991. Estimates of **residualism** of hatchery reared summer steelhead and population size of yearling rainbow trout (*Oncorhynchus mykiss*) in southeast Washington, 1991. Washington Department of Wildlife. Report No. AFF 1/**LSR-92-02. 16 pp.**
- Volkman, Jed W. 1993. Evaluation of juvenile fish bypass and adult fish passage facilities at water diversions in the Umatilla River. Volume 2. Bonneville Power Administration. Project No. **89-024-01.**

- Volkman, **Jed W.** 1994. Evaluation of adult **salmonid** passage at water diversions on the Umatilla River and their movement following upriver transport and assessment of factors for homing into the Umatilla River pp. 101-143 in Knapp, S. M. editor. Evaluation of juvenile fish bypass and adult fish passage facilities at water diversions in the Umatilla River. Report submitted to U.S. Department of Energy, Bonneville Power Administration, **DOE/BP-01385-5.**
- Wagner, Paul G. 1990. Evaluation of the use of the McNary bypass system to divert adult **fall-**backs away from turbine intakes. Contract No. **DACW68-82-C-0077.**
- Wagner, Paul G., **Hillson**, Todd. 1991. Evaluation of adult fall-back through the McNary Dam juvenile bypass system. Report prepared for U.S. Army Corps of Engineers, Contract No. **DACW68-82-C-0077.**
- Walker, George W., **MacLeod**, Norman **S.** 1991. Geologic Map of Oregon, U.S. Geological Survey, U.S Department of the Interior. U.S. Geological Survey Map Distribution, Box 25286, Federal Center, Denver, CO 80225.
- Zimmerman, Brian C., Duke, Bill. 1995. Trapping and transportation of adult and juvenile salmon in the lower Umatilla River in northeast Oregon. Report submitted to Bonneville Power Administration, DOE/BP No. 98636-2.



Map compiled from various sources by the C.T.U.I.R. Mapping Dept. Locations are approximate.

Figure A-1. Map of Reservation and Ceded Lands of the Umatilla Indian Reservation in Northeast Oregon and

Washington
Oregon





Figure A-2. Map of the Umatilla River Basin with River Miles Denoted.




Figure A-3. Map of Index Site Locations in the Umatilla River Basin.




Figure A-4. Map of Spawning Areas.

APPENDIX B

Table B-1. Estimated Natural Populations of Summer Steelhead and Spring Chinook Salmon in the Umatilla River Basin.

| Umatilla River Basin Tributaries in the | Miles | Suitable Miles (STS) | Mile/ | Total STS | Suitable Miles (CHS) | CH/ Mile | Total CH |
|---|--------|----------------------|---------|-----------|----------------------|----------|----------|
| Umatilla River: RM 205-27.2 | 28.9 | 0.5 | 1'000 | 500 | 0.5 | | 25 |
| Umatilla River: 55.3-60.8 | 5.5 | 0.5 | 1'000 | 500 | 0.5 | 50 | 25 |
| Umatilla River: 60.8-64.2 | 3.4 | 0.1 | 0 | 0 | 0 | 0 | 0 |
| Umatilla River: 64.2-81.8 | 7 | *1.6 | *22 | 150 | *0 | *0 | 100 |
| Umatilla River: 81.8-89.6 | 89.6 | 17.6 | *1,650 | *29,040 | *7 | 1,250 | 10,000 |
| Subtotal | 134.4 | 28.2 | 98,392 | *58,744 | 25.6 | 1,441 | *10,087 |
| Butter Creek | 95 | 0 | 0 | 0 | 0 | 0 | 0 |
| Alkali Canyon | 20 | 8 | 0 | 8 | 0 | 0 | 8 |
| Spear canyon | 12 | 0 | 0 | 0 | 0 | 0 | 0 |
| Coombs Canyon | 18 | 0 | 0 | 0 | 0 | 8 | 8 |
| McKay Creek | 12 | 0 | 0 | 0 | 0 | 0 | 0 |
| Tutuilla Creek | 16 | 8 | 8 | 0 | 0 | 0 | 8 |
| Patawa Creek | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Wildhorse Creek | 90 | *0 | *0 | *0 | *0 | 10 | *0 |
| Subtotal | 343 | 0 | 0 | 0 | 0 | 0 | 0 |
| Birch Creek | 16 | 0 | 0 | 0 | 0 | 0 | 0 |
| Stewart Creek | 12 | 0 | 8 | 0 | 0 | 8 | 8 |
| West Birch Creek | 20 | 16 | 1,509 | *24,144 | 10 | *0 | 100 |
| Bridge Creek | 9 | 3 | 100 | 300 | 0 | 0 | 0 |
| Bear Creek | 13 | 10 | 500 | 5,000 | 8 | 8 | 8 |
| Stanley Creek | 6 | 4 | 100 | 400 | 0 | 0 | 0 |
| Willow Spring Can. | 7 | 15 | 500 | 2,000 | 0 | 0 | 0 |
| East Birch Creek | 18 | 0 | 4,916 | *73,740 | 10 | *0 | 100 |
| Wagner Creek | 7 | 0 | 0 | 0 | 0 | 0 | 0 |
| Spring Hollow | 6 | 0 | 0 | 0 | 0 | 0 | 0 |
| California | 12 | 0 | 0 | 0 | 0 | 0 | 0 |
| Creek Pearson Creek | 11 | 11 | 4,500 | 49,500 | 0 | 0 | 0 |
| sooth Canyon Creek | 5 | 4 | 1,000 | 4,000 | 0 | 0 | 0 |
| Westgate Canyon | 2 | 2 | 5,500 | 11,000 | 0 | 0 | 0 |
| Subtotal | 141 | 69 | 170,084 | 170,084 | 0 | 0 | 0 |
| Mission Creek | 7 | 1 | 279 | *837 | *0 | *0 | *0 |
| Cottonwood Creek | 5 | *2 | *292 | *584 | 1 | *22 | *22 |
| Moonshine Creek | 4 | *2 | *267 | *1,134 | 1 | 10 | *0 |
| Coonskin Creek | 4 | 2 | *712 | *1,424 | 1 | *9 | *9 |
| Buckaroo Creek | 6 | *3.3 | *1,200 | 3,600 | 10 | 10 | *0 |
| Subtotal | 26 | 12.3 | 7940 | 7940 | 0 | 0 | 31 |
| Squaw Creek | 13 | 8.75 | 4,367 | 38,211 | *8.75 | 50 | 1,102 |
| Batchelor Creek | 4 | 1 | 1,000 | 1,000 | 1 | 50 | 50 |
| Little squaw Creek | 1 | 1.5 | 1,000 | 1,500 | 10.7 | 0 | 50 |
| Subtotal | 17 | 11.25 | 40,711 | 40,711 | 0 | 0 | 1,202 |
| Meacham Creek, Lower 15 miles | 14 | 12.9 | *5,576 | 71,930 | *12.9 | *500 | 6,450 |
| Boston Canyon Creek | 3 | *2 | 1,650 | 3,300 | *0 | 10 | 10 |
| Line Creek | 3 | *2.4 | *1,931 | 4,634 | *0 | *0 | 10 |
| Camp Creek below falls | 3.1 | 3.1 | 2,144 | *6,646 | 10 | 10 | *0 |
| North Fork Meacham Creek tributary | 0.2 | 0 | *0 | *0 | 10 | 10 | 10 |
| North Fork Meacham Creek | 2 | *0 | *0 | *0 | *0 | *0 | *0 |
| Bear Creek | 10 | 3 | 4,500 | 36,000 | 0 | 1000 | 4,000 |
| Subtotal | 46 | 35.4 | 1,000 | 174,000 | 16.9 | 0 | 10,450 |
| Meacham Creek, Upper 21 miles | 21 | 17 | 4,500 | 76,500 | 0 | 0 | 0 |
| East Meacham Creek | 7 | 3 | 9,000 | 9,000 | 0 | 0 | 0 |
| Butcher Creek | 4 | 4 | 1,000 | 4,000 | 8 | 0 | 8 |
| Beaver Creek | 4 | 2 | 1,000 | 3,000 | 0 | 0 | 0 |
| Subtotal | 9 | 3 | 1,000 | 94,500 | 0 | 0 | 8 |
| Ryan Creek | 3 | 5 | 4,500 | 22,500 | 3 | 100 | 300 |
| Bobsled Creek | 1 | 1 | 1,000 | 1,000 | 0 | 0 | 0 |
| Bear Creek | 1 | 7 | 1,000 | 24,500 | 3 | 8 | 0 |
| Subtotal | 5 | 9 | 0 | 0 | 0 | 0 | 300 |
| North Fork Umatilla | 15 | 3 | 5,500 | 49,500 | 1 | 1,500 | 4,500 |
| Coyote Creek | 2 | 1 | 1,500 | 1,500 | 0 | 50 | 50 |
| Woodward Creek | 2 | 1 | 1,500 | 1,500 | 0 | 0 | 0 |
| Johnson Creek | 19 | 1 | 1,500 | 1,500 | 0 | 0 | 0 |
| Subtotal | 38 | 6 | 3,500 | 57,000 | 0 | 0 | 4,550 |
| South Fork Umatilla | 11 | 5 | 2,500 | 31,500 | 4 | 500 | 2,000 |
| Buck Creek | 6 | 5 | 2,000 | 12,500 | 2 | 500 | 1,000 |
| Thomas Creek | 5 | 4 | 2,000 | 10,000 | 2 | 500 | 1,000 |
| Shimshorn Creek | 5 | 4 | 2,000 | 8,000 | 1 | 50 | 50 |
| Subtotal | 31 | 27 | 70,000 | 70,000 | 10 | 50 | 4,100 |
| TOTAL | 769.90 | 233.15 | | 724,773 | 64.25 | | 52,770 |

* Estimated from empirical data




Figure B-1. Adult **Steelhead** Returns Compared to the Mean **Annual** Flows (cfs) at Umatilla Gage (**RM 1.2**) Two Years Prior to the Adult Return from 1982/3 to 1996/7, (1995/6 and 1996/7 adult returns approximated; STSFLWB 1. CH3)




Figure B-2. Adult **Steelhead** Returns and the Average of February, March and April Mean Monthly Flows (cfs) at Umatilla Gage (**RM 1.2**) Two Years Prior to the Adult Return from 1982/3 to 1996/7 (1995/6 and 1996/7 adult returns approximated; STSFLWB2.CH3).

APPENDIX c
Thermograph Locations and Recorded -Temperatures

Table C-1. **Thermographs in the Umatilla River.**

| LOCATION | AGENCY | RIVER MILE | DEPLOYMENT PERIOD | THERMOGRAPH TYPE |
|---|---------------|-------------------|--------------------------|-------------------------|
| Umatilla River (at Three Mile Falls Dam) | CTUIR | 3.7 | All Year | Temp-Mentor |
| Umatilla River (at Three Mile Falls Dam) | USBR | 3.7 | All Year | Hydromet |
| Umatilla River (at Maxwell Canal @ new gage) | USBR | 15 | All Year | Hydromet |
| Umatilla River (near Dillon Canal, at gage 03 10) | USBR | 24 | All Year | Hydromet |
| Umatilla River (near Feed Canal, at gage 0290) | USBR | 28 | All Year | Hydromet |
| Umatilla River (near Yoakum, at gage 0260) | USBR | 37 | All Year | Hydromet |
| Umatilla River (Near Rieth) | CTUIR | 49 | Moved to 42.5 | RTM2000 |
| Umatilla River (Near Barnhart) | CTUIR | 42.5 | All Year | RTM2000 |
| Umatilla River (Near Pendleton, at gage 0210) | USBR | 55.2 | All Year | Hydromet |
| Umatilla River (Near ODFW Office) | CTUIR | 56 | All Year | Temp-Mentor |
| Umatilla River | CTUIR | 78.5 | All Year | Temp-Mentor |
| Umatilla River | CTUIR | 79 | All Year | Temp-Mentor |
| Umatilla River (at USGS Gage) | CTUIR | 81.7 | All Year | Temp-Mentor |
| Umatilla River (Below mouth of N. and S. Forks) | USFS | 89.5 | Feb.-Dec. | Temp-Mentor |
| Minthorn Springs (Near Umatilla RM 65) | CTUIR | In springs | All Year | Temp-Mentor |
| Mission Creek | CTUIR | 3 | All Year | RTM2000 |
| Buckaroo Creek | CTUIR | 2 | All Year | Temp-Mentor |
| Squaw Creek | CTUIR | 2 | All Year | Temp-Mentor |
| Little Squaw Creek | CTUIR | 0.1 | All Year | Temp-Mentor |
| N.Fork Umatilla River | USFS | 0.1 | June-Oct. | Temp-Mentor |
| S.Fork Umatilla River | USFS | 0.1 | Feb.-Dec. | Temp-Mentor |
| S.Fork Umatilla River | USFS | 6 | June-Oct. | Temp-Mentor |
| Shhnmiehom | USFS | 0.1 | June-Oct. | Temp-Mentor |

Table C-2. Thermographs in Meacham Creek Drainage.

| LOCATION | AGENCY | RIVER MILE | DEPLOYMENT PERIOD | THERMOGRAPH TYPE |
|--|---------------|-------------------|--------------------------|-------------------------|
| Meacham Creek | CTUIR | 2 | All Year | Temp-Mentor |
| Meacham Creek | CTUIR | 5.25 | All Year | Temp-Mentor |
| Meacham Creek | CTUIR | 13 | Discontinued (lost) | RTM2000 |
| Meacham Creek | ODFW | 31.5 | All Year | Temp-Mentor |
| Meacham Creek | ODPW | 32.5 | All Year | Temp-Mentor |
| Bonifer Pond (near Meacham C. RM 2.5) | CTUIR | In Pond | All Year | Temp-Mentor |
| Camp Creek | CTUIR | 0.6 | All Year | RTM2000 |
| N .F. Meacham | ODFW | 0.1 | April to October | Hobo |
| N .F. Meacham | USFS | 2 | June-Oct. | Temp-Mentor |
| East Meacham | CTUIR | 0.1 | All Year | RTM2000 |
| Butcher Creek | CTUIR | 1 | All Year | RTM2000 |

Table C-3. Thermographs in Wildhorse Creek Drainage

| LOCATION | AGENCY | RIVER MILE | DEPLOYMENT PERIOD | THERMOGRAPH TYPE |
|-------------------------------------|--------|------------|-------------------|------------------|
| Wildhorse Creek (Mouth) | CTUIR | 0 | All Year | Temp-Mentor |
| Wildhorse Creek (Below new project) | CTUIR | 9.5 | All Year | Temp-Mentor |
| Wildhorse Creek (Above new project) | CTUIR | 11 | All Year | Temp-Mentor |
| Wildhorse Creek (Near Adams) | ODFW | 13 | All Year | Temp-Mentor |
| Wildhorse Creek (Headwaters) | CTUIR | 26 | All Year | Temp-Mentor |

Table C-4. Thermographs in the Walla Walla River Basin

| LOCATION | AGENCY | RIVER MILE | DEPLOYMENT PERIOD | THERMOGRAPH TYPE |
|--------------------------------------|--------|------------|-------------------|------------------|
| Walla Walla River | CTUIR | 8 | All Year | Temp-Mentor |
| Walla Walla River | CTUIR | 47 | All Year | Temp-Mentor |
| S.F. Walla Walla | CTUIR | 0.5 | All Year | RTM2000 |
| S.F. Walla Walla | CTUIR | 7. | All Year | Temp-Mentor |
| S.F. Walla Walla | CTUIR | 20 | All Year | RTM2000 |
| Elbow Creek (S.F. Walla Walla) | ODFW | 0.1 | April-Dec | Hobo |
| Burnt Cabin Creek (S.F. Walla Walla) | CTUIR | 0.1 | Discontinued | RTM2000 |
| Reser Creek (S.F. Walla Walla) | CTUIR | 0.1 | All Year | RTM2000 |
| N.F. Walla Walla | CTUIR | 0.1 | All Year | Temp-Mentor |
| N.F. Walla Walla | ODFW | 6 | April-Dec | Hobo |
| N.F. Walla Walla | ODFW | 12 | April-Dec | Hobo |
| Pine Creek | ODFW | 20.5 | All Year | Temp-Mentor |
| Pine Creek | ODFW | 29 | All Year | Temp-Mentor |

Table C-5 Thermographs in Birch Creek, Butter Creek, and Willow Creek Drainages.

| LOCATION | AGENCY | RIVER MILE | DEPLOYMENT PERIOD | THERMOGRAPH TYPE |
|------------------------------------|--------|------------|-------------------|------------------|
| Birch Creek | ODFW | 3.5 | All Year | Temp-Mentor |
| Birch Creek (near Sparks) | ODFW | 6.5 | All Year | Temp-Mentor |
| East Birch Creek | ODFW | 8.5 | All Year | Temp-Mentor |
| Westgate Canyon (East Birch Creek) | ODFW | 0.75 | All Year | Temp-Mentor |
| Pearson Creek | ODFW | 4 | April-Oct. | Hobo |
| West Birch Creek | ODFW | 2 | All Year | Hobo |
| West Birch Creek | ODFW | 15 | All Year | Hobo |
| Butter Creek | ODFW | 51 | April-Oct. | Hobo |
| Little Butter Creek (Near Gurdane) | ODFW | 7 | April-Oct. | Hobo |
| Little Butter Creek (Near Lena) | ODFW | 19.5 | April-Oct. | Hobo |
| Willow Creek | ODFW | 61 | April-Oct. | Hobo |
| Willow Creek | ODFW | 77.5 | April-Oct. | Hobo |
| Rhea Creek | ODFW | 16.7 | April-Oct. | Hobo |
| Rhea Creek | ODFW | 35 | April-Oct. | Hobo |




Figure C-1. Maximum and Minimum Temperatures Recorded in the Umatilla River, Near Rieth, RM 49.5, December 94 through May 1995 (TGUR9412.CH3).




Figure C-2. Maximum and Minimum Temperatures Recorded in the Umatilla River, Barnhart, RM 42.5, February Through June, 1995 (TCUB9502.CH3).




Figure C-3. The Maximum and Minimum Temperatures Recorded the Umatilla River, **near Barnhart** RM 42.5, June into December, 1995 (TCUB9506.CH3).




Figure C-4. Maximum **and** Minimum Temperatures Recorded in Butcher Creek, RM 1.5, May, 1995 to July, 1995 (TGBT9505.CH3).




Figure C-5. Maximum and Minimum Temperatures Recorded in Wildhorse Creek, RM 1.5, May, 1995 to January, 1996 (TGWD9505.CH3).




Figure C-6. Maximum and Minimum Temperatures Recorded in Mission Creek, RM 3, February through November, 1995 (TCMC9502.CH3).




Figure C-7. Maximum and Minimum Temperatures Recorded in Camp Creek, RM 0.5, December, 1994 to May, 1995 (TCCP9412.CH3).




Figure C-8. Maximum and Minimum Temperatures Recorded in Camp Creek, RM 0.5, August, 1994 to January, 1995 (TGCP9508.CH3).




Figure C-9. Maximum and Minimum Temperatures Recorded in East **Meacham** Creek, RM 0.125, April Through July, 1995 (TGME9504.CH3).

APPENDIX D
Physical Habitat Survey Data Summary Tables.

Table D-1. The Stream, **RM Range**, **RM Surveyed**, **Total Area**, Range of Elevation, Number of Habitat Units and Date of Habitat Surveys.

| STREAM | SECTION (RM) | RM SURVEYED | TOTAL AREA/m² | ELEVATION RANGE | # HABITAT UNITS | DATE (1995) |
|------------------|---------------------|--------------------|---------------------------------|------------------------|------------------------|--------------------|
| Umatilla River | 81.8-89.6 | 7.8 | 151,949 | 1,880-2,320 | 639 | 7/18-8/7 |
| Moonshine Creek | 0.0-4.4 | 4.4 | 11,213 | 1,400-2,590 | 594 | 8/28-9/5 |
| Mission Creek | 0.0-4.3 | 4.3 | 9,994 | 1,270-2,200 | 872 | 8/15-9/11 |
| Cottonwood Creek | 0.0-4.1 | 4.1 | 15,431 | 1,330-2,200 | 912 | 6/20-8/1 |
| Coonskin Creek | 0.0-2.0 | 2.0 | 5,860 | 1,420-1,890 | 626 | 6/20-7/17 |
| TOTAL | - | 22.6 | 194,447 | 1,270-2,590 | 3,643 | 6/20-9/11 |

Table D-2. Summary of Habitat Quality Rankings from Habitat Survey Data, 1995 (AC = Active Channel).

| HABITAT FEATURE | NUMERICAL VALUE OF HABITAT FEATURE | | | | |
|--|---|------------------------|----------------------|-------------------------|-----------------------|
| | Umatilla River | Moonshine Creek | Mission Creek | Cottonwood Creek | Coonskin Creek |
| Min Stream Temperature (C) | 10.0 | 10.0 | 6.0 | 10.5 | 11.0 |
| Max Stream Temperature (C) | 32.0 | 23.0 | 14.0 | 27.0 | 29.0 |
| Pool Area (%) | 29.4 | 18.5 | 10.0 | 24.9 | 29.5 |
| Mean Depth (m) | 0.45 | 0.15 | 0.09 | 0.12 | 0.18 |
| AC Width:Depth-All Units | 22.6 | 8.9 | 9.3 | 8.9 | 7.6 |
| AC Width:Depth-Riffles | 35.4 | 20.8 | 32.9 | 20.8 | 19.2 |
| Dry Channel (5%) | 0.3 | 58.6 | 76.3 | 49.2 | 0.2 |
| Undercut Bank (%) | 8.6 | 6.0 | 8.2 | 10.9 | 11.2 |
| Boulder Count | 4,772 | 1,158 | 35 | 522 | 307 |
| Wood Pieces (#/100m) | 1.5 | 1.2 | 6.6 | 3.4 | 1.6 |
| Wood Volume (m³/100m) | 2.1 | 0.6 | 1.6 | 0.9 | 1.2 |
| Mean Wood Complexity (#/unit) | 1.3 | 1.2 | 1.6 | 1.5 | 1.5 |
| Gravel (% of Wetted Area) | 35 | 36 | 44 | 37 | 34 |
| Silt-Sand-Organics (% Area) | 16 | 21 | 24 | 32 | 31 |
| # of Artificial Fish Passage Barriers | 0 | 1 | 2 | 3 | 1 |
| Mean Slope of all Habit Units | 1.4 | 2.7 | 2.8 | 3.3 | 3.1 |
| Eroding Bank (I) | 7.1 | 6.0 | 21.3 | 12.1 | 13.2 |
| Mean Surface Slope of Riparian (%) | 36 | 23 | 20 | 18 | 23 |
| Mean Open Sky of All Units (%) | 49 | 44 | 38 | 47 | 41 |
| Mean Riparian Canopy Closure (%) | 29 | 16 | 12 | 25 | 23 |
| Valley Width Index (VWI) | 5.0 | 10.0 | 31.1 | 19.6 | 11.5 |

Table D-3. Habitat Unit Summary for the Umatilla River, RM 81.8 to 89.6, July 18-August 7, 1995.

| REACH 0 | | | | | | | | | | | | REACH 0 | | | | | | | | | | | | | |
|---------------------------|--------------|------------------|---------------|---------------|------------------------------|-------------------------|-----------|-----------|------|------|------|---------------------|-----|------|------|------|---|---|---|---|---|---|---|---|---|
| HABITAT DETAIL | | | | | | | | | | | | | | | | | | | | | | | | | |
| Habitat Type | Number Units | Total Length (m) | Avg Width (m) | Avg Depth (m) | Total Area (m ²) | Large Boulders (#>0.5m) | S/O | Substrate | | | | Percent Uetted Area | | | | | | | | | | | | | |
| | | | | | | | | Snd | Crvl | cbbl | Bldr | S/O | Snd | Crvl | cbbl | Bldr | | | | | | | | | |
| DRY UNITS | 1 | 6 | 4.4 | 0.00 | 24 | 0 | 0 | 10 | 40 | 40 | 10 | 10 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| GLIDE | 63 | 1,321 | 7.6 | 0.47 | 13,871 | 558 | 10 | 13 | 33 | 28 | 13 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| POOL-BACKWATER | 42 | 316 | 2.3 | 0.30 | 755 | 62 | 16 | 20 | 31 | 22 | 9 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| POOL-BEAVER DAM | 1 | 67 | 7.8 | 2.00 | 519 | 0 | 30 | 20 | 20 | 10 | 10 | 10 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| POOL-DAMMED | 5 | 92 | 6.7 | 0.56 | 680 | 22 | 12 | 18 | 32 | 26 | 12 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| POOL-ISOLATED | 24 | 1,369 | 2.4 | 0.41 | 4,640 | 116 | 13 | 15 | 33 | 26 | 10 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| POOL-LATERAL SCOUR | 108 | 2,204 | 8.7 | 0.88 | 23,629 | 493 | 6 | 12 | 33 | 29 | 13 | 7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| POOL-PLUNGE | 3 | 28 | 6.7 | 1.02 | 250 | 13 | 13 | 17 | 33 | 20 | 13 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| POOL-STRAIGHT SCWR | 63 | 1,271 | 9.1 | 0.70 | 14,201 | 459 | 5 | 9 | 3 | 4 | 34 | 16 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| PUDDLED CHANNEL | 6 | 224 | 1.9 | 0.23 | 461 | 4 | 5 | 12 | 35 | 37 | 12 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RAPID/BEDROCK | 3 | 21 | 5.5 | 0.33 | 131 | 10 | 0 | 0 | 13 | 23 | 20 | 43 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RAPID/BOULDERS | 63 | 1,021 | 8.7 | 0.29 | 9,614 | 492 | 0 | 1 | 35 | 40 | 22 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RIFFLE | 206 | 5,525 | 8.9 | 0.26 | 60,403 | 1249 | 3 | 9 | 38 | 36 | 13 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| RIFFLE W/ POCKETS | 47 | 1,849 | 10.9 | 0.35 | 22,653 | 1282 | 4 | 10 | 32 | 34 | 19 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| STEP/BOULDERS | 1 | 2 | 11.1 | 0.30 | 24 | 10 | 10 | 10 | 20 | 40 | 20 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| STEP/LOG | 1 | 0 | 2.8 | 0.15 | 1 | 0 | 10 | 20 | 40 | 20 | 10 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| STEP/STRUCTURE | 2 | 6 | 11.1 | 0.15 | 95 | 2 | 10 | 10 | 35 | 25 | 15 | 5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total: | 639 | 15,322 | 8.1 | 0.45 | 151,949 | 4772 | Avg: 6 | 10 | 35 | 32 | 14 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| HABITAT SUMMARY | | | | | | | | | |
|------------------------------|-----------|------------------|---------------|---------------|-------------------------------|-----------------------|-----------------------------|------------|-------|
| Habitat Group | No. Units | Total Length (m) | Avg Width (m) | Avg Depth (m) | Wetted Area (m ²) | Large Boulders Number | Percent #/100m ² | Wood Class | |
| | | | | | | | | Number | Class |
| Dammed & BW Pools | 72 | 1,843 | 2.7 | 0.38 | 6593 | 4.34 | 200 | 3.03 | 1.4 |
| Scour Pools | 174 | 3,503 | 8.8 | 0.82 | 38080 | 25.06 | 965 | 2.53 | 1.4 |
| Glides | 63 | 1,321 | 7.6 | 0.47 | 13871 | 9.13 | 558 | 4.02 | 1.3 |
| Riffles | 253 | 7,374 | 9.3 | 0.27 | 83056 | 54.66 | 2531 | 3.05 | 1.1 |
| Rapids | 66 | 1,043 | 8.6 | 0.29 | 9745 | 6.41 | 502 | 5.15 | 1.0 |
| Cascades | 0 | 0 | . | . | 0 | 0.00 | 0 | 0.00 | . |
| Step/Falls | 4 | 9 | 9.0 | 0.19 | 120 | 0.08 | 12 | 10.02 | 1.0 |
| Small Streams (SS) | 0 | 0 | . | . | 0 | 0.00 | 0 | 0.00 | . |
| Dry | 7 | 230 | 2.3 | 0.19 | 485 | 0.32 | 4 | 0.82 | 1.1 |

Table D-4. Stream Summary for the Umatilla River, RM 81.8 to 89.6, July 18-August 7, 1995.

| STREAM SUMMARY | | | | UMATILLA RIVER | | | | | | | | |
|-----------------|------------------------|---------------------|---------------------|------------------------------------|----------------------------------|------|------|------|------|---------------------------|-------|---|
| Number Units | Total Length (m) | Avg Yidth (m) | Avg Depth (m) | Total Area (m ²) | Substrate Percent Uetted Area | | | | | Total Large Boulder | | |
| | | | | | S/O | Sand | Grvl | Cbbl | Bldr | Bdrk | | |
| - | - | - | - | - | - | - | - | - | - | - | - | - |
| 639 | 15,322 | 8.1 | 0.45 | 151,949 | 6 | 10 | 35 | 32 | 14 | 3 | 4,772 | |

| Wetted Area | | |
|------------------------|-------------------|---------|
| Habitat Group | (m ²) | Percent |
| Scour Pool | 38,080 | 25.1 |
| Backwater Pools | 6,593 | 4.3 |
| Glide | 13,871 | 9.1 |
| Riffle | 83,056 | 54.7 |
| Rapid | 9,745 | 6.4 |
| Cascade | 0 | 0.0 |
| Step | 120 | 0.1 |
| Dry | 485 | 0.3 |

Table D-5. Valley, Channel, Bank and Wood Summary for the Umatilla River. RM 81.8 to 89.6, July 18-August 7, 1995.

Valley and Channel Summary

Valley Characteristics (Percent Reach Length)

| Narrow Valley Floor | | Broad Valley Floor | |
|---------------------|---|-----------------------|-----|
| Steep V-shape | 0 | Constraining Terraces | 100 |
| Moderate V-shape | 0 | Multiple Terraces | 0 |
| Open V-shape | 0 | Wide Floodplain | 0 |

Valley Width Index avg: 5.0 range: 5.0-5.0

Channel Morphology (Percent Reach Length)

| Constrained | | Unconstrained | |
|-------------------|-----|------------------|---|
| Hillslope | 0 | Single Channel | 0 |
| Bedrock | 0 | Multiple Channel | 0 |
| Terrace | 0 | Braided Channel | 0 |
| Alt. Terrace/Hill | 100 | | |
| Landuse | 0 | | |

Channel Characteristics

| Type | Length | Area | Dry Units |
|-----------|--------|---------|-----------|
| Primary | 10,525 | 132,443 | 0 |
| Secondary | 4,797 | 19,505 | 7 |

Channel Dimensions

| Wetted Surface | | Active Channel | | First Terrace | |
|----------------|------|----------------|------|---------------|------|
| Width | 8.1 | Width | 16.3 | Width | 1a.9 |
| Depth | 0.45 | Height | 0.4 | Height | 0.8 |
| W:D | 35.4 | | | | |

Stream Flow Type: MF Water Temp: 11.0-11.0
 Avg. Unit Gradient: 1.4 Habitat Units/100m: 4.2

Riparian, Bank, and Wood Summary

Land Use: ST,TT Riparian Veg.: C 30-50 D 1

Bank Stability

| Bank Class | Percent Reach Length | Undercut Banks |
|-----------------------|----------------------|---------------------|
| Non-Erodible | 7.8 | Unit Average: 8.64% |
| Vegetation Stabilized | 74.6 | Open Sky (% of 180) |
| Boulder-cobble | 10.4 | Unit Average: 49 |
| Actively Eroding | 7.1 | Range: 3-69 |

Large Woody Debris

| | | | |
|---------------------------|-----|-------------------------|-----|
| Average Complexity Score: | 1.3 | | |
| Pieces | 163 | Volume(m ³) | 221 |
| Pieces/100m | 1.5 | Volume/100m | 2.1 |

Table D-6. Riparian Summary for the Umatilla River, RM 81.8 to 89.6, July 18-August 7, 1995.

REACH 0 REACH 0

RIPARIAN ZONE VEGETATION SUMMARY

Reach 0 is represented by 22 transects

Predominant **landform** in each zone

| | Zone 1 0-10 meters | Zone 2 10-20 meters | Zone 3 20-30 meters |
|-----------------------|-----------------------|------------------------|------------------------|
| Hillslope | 9 | 1a | 30 |
| High terrace | 27 | 23 | 16 |
| Low terrace | 45 | 41 | 43 |
| Floodplain | 0 | 0 | 0 |
| Wetland/meadow | 0 | 0 | 0 |
| Stream channel | 11 | 14 | 9 |
| Roadbed/Railroad | 0 | 0 | 0 |
| Riprap | 0 | 0 | 0 |
| Surface slope (%) | 41 | 33 | 35 |

Canopy closure and ground cover

| | Zone 1 0-10 meters (%) | Zone 2 10-20 meters (%) | Zone 3 20-30 meters (%) |
|-------------------------|------------------------------|-------------------------------|-------------------------------|
| Canopy closure | 31 | 29 | 28 |
| Shrub cover | 39 | 37 | 42 |
| Grass/forb cover | 30 | 37 | 38 |

Average number of trees in a **5-meter wide** band

| Diameter class (cm) | Zone 1 0-10 meters | | Zone 2 10-20 meters | | Zone 3 20-30 meters | | Zones 1-3 0-30 meters | |
|-------------------------------|-----------------------|----------|------------------------|----------|------------------------|----------|--------------------------|----------|
| | Conifer | Hardwood | Conifer | Hardwood | Conifer | Hardwood | Conifer | Hardwood |
| 3-15cm | 0.6 | 4.4 | 0.4 | 1.8 | 0.3 | 1.6 | 1.3 | 7.8 |
| 15-30cm | 0.1 | 0.5 | 0.5 | 0.1 | 0.3 | 0.3 | 0.9 | 1.0 |
| 30-50cm | 0.2 | 0.5 | 0.7 | 0.3 | 0.6 | 0.2 | 1.6 | 1.0 |
| 50-90cm | **.* | 0.2 | 0.1 | 0.1 | 0.0 | **.* | 0.1 | 0.4 |
| >90cm | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Total/100m² | 1.0 | 5.7 | 1.7 | 2.3 | 1.3 | 2.1 | 1.3 | 3.4 |

Table D-7. Water Diversions in the Umatilla River, RM 81.8-89.6, Habitat Survey 7/18-8/7, 1995.

| UNIT NUMBER | RIVER MILE | DESIGNATED USE | TYPE | SIZE |
|-------------|------------|----------------|---------------------------------|-----------------------------------|
| 5 | 81.9 | Private Pond | Partially Screened Ditch | 1m wide x .22m deep |
| 24 | 82.0 | Private | screened PVC Pipe | 2" |
| 94 | 82.7 | private | Screened PVC pipe | 1.5" |
| 95 | 82.7 | Private | Screened Metal pipe | 2' |
| 391 | 87.6 | private | Screened Metal pipe | 1.5" |

Table D-8. Surface Springs identified in the Habitat Survey, Umatilla River, Survey Dates 7/18-8/7, 1995.

| RIVER MILE | UNIT TYPE | BANK SIDE | | AREA (m ²) |
|--------------|-----------|----------------|-----------------|------------------------|
| 82.0 | BW | LEFT | | 21 |
| 83.1 | LP | RIGHT | | 108 |
| 83.3 | RI | RIGHT | | 221 |
| 83.7 | IP | RIGHT | | 195 |
| 83.7 | IP | LEFT | | 60 |
| 84.2 | IP | LEFT | | 10 |
| 84.5 | IP | RIGHT | | 21 |
| 84.7 | IP | RIGHT | | 150 |
| 84.8 | IP | RIGHT | | 980 |
| 85.0 | IP | LEFT | | 750 |
| 85.5 | GL | LEFT | | 140 |
| 85.5 | IP | LEFT | | 210 |
| 85.6 | IP | LEFT | | 320 |
| 85.8 | IP | RIGHT | | 1,050 |
| 86.0 | GL | LEFT | | 90 |
| 86.0 | IP | LEFT | | 45 |
| 86.3 | IP | LEFT | | 400 |
| 86.3 | IP | LEFT | | 24 |
| 86.4 | GL | RIGHT | | 22 |
| 86.6 | IP | RIGHT | | 35 |
| 87.8 | IP | RIGHT | | 60 |
| 87.8 | LP | RIGHT | | 132 |
| 89.1 | BW | LEFT | | 70 |
| 89.1 | IP | RIGHT | | 50 |
| 89.2 | IP | RIGHT | | 15 |
| 89.2 | IP | LEFT | | 130 |
| 89.4 | IP | RIGHT | | 180 |
| TOTAL | 27 | 13 LEFT | 14 RIGHT | 5,489 |

Table D-9. Habitat Unit Summary for Moonshine! Creek, RM 0.0 to 4.4, August 28-September 5, 1995.

| REACH 0 | | | | | | | | | | | | REACH 0 | |
|---------------------|-----------------|------------------------|---------------------|---------------------|------------------------------------|-------------------------------|----------------|---------------|------|-----------|------|----------|----|
| HABITAT DETAIL | | | | | | | | | | | | | |
| Habitat Type | Number Units | Total Length (m) | Avg Width (m) | Avg Depth (m) | Total Area (m ²) | Large Boulders (#>0.5m) | Substrate | | | | | | |
| | | | | | | | Percent Uetted | | Area | | | | |
| | | | | | | | S/O | Snd | Grvl | Cbbl | Bldr | Bdrk | |
| CASCADE/BEDROCK | 3 | 19 | 1.2 | 0.20 | 25 | 3 | 7 | 13 | 20 | 17 | 7 | 37 | |
| CULVERT CROSSING | 2 | 18 | 1.5 | 0.05 | 31 | 0 | 5 | 0 | 0 | 0 | 0 | 45 | |
| DRY CHANNEL | 43 | 1,981 | 2.8 | 0.12 | 5,494 | 655 | 0 | 10 | 36 | 39 | 13 | 2 | |
| DRY UNITS | 12 | 306 | 2.2 | 0.00 | 702 | 35 | 0 | 9 | 45 | 30 | 16 | 0 | |
| GLIDE | 48 | 332 | 1.4 | 0.17 | 523 | 25 | 15 | 11 | 37 | 28 | a | 1 | |
| POOL-BACKWATER | 11 | 9 | 1.2 | 0.21 | 11 | 3 | 11 | 11 | 35 | 25 | 5 | 14 | |
| POOL-BEAVER DAR | 3 | a2 | 5.0 | 0.68 | 612 | 0 | 45 | 32 | 19 | 3 | 0 | 0 | |
| POOL-ISOLATED | 10 | 145 | 0.8 | 0.22 | 170 | 0 | 23 | 31 | 29 | 17 | 0 | 0 | |
| POOL-LATERAL SCOUR | 110 | 487 | 1.4 | 0.26 | 729 | 53 | 10 | 11 | 37 | 31 | 9 | 2 | |
| POOL-PLUNGE | 9 | 22 | 3.0 | 0.49 | 75 | 5 | 13 | 13 | 34 | 26 | 11 | 2 | |
| POOL-STRAIGHT SCOUR | 68 | 273 | 1.5 | 0.22 | 467 | 51 | 11 | 8 | 36 | 30 | 10 | 5 | |
| POOL-TRENCH | 2 | 7 | 1.0 | 0.45 | a | 1 | 10 | 10 | 35 | 25 | 10 | 10 | |
| PUDDLED CHANNEL | 13 | 298 | 1.1 | 0.18 | 376 | 100 | 10 | 10 | 32 | 31 | 15 | 2 | |
| RAPID/BEDROCK | 9 | 45 | 1.2 | 0.05 | 58 | 2 | 15 | 7 | 13 | 10 | 4 | 50 | |
| RAPID/BOULDERS | 48 | 220 | 1.4 | 0.05 | 306 | 65 | 10 | 7 | 35 | 33 | 15 | 1 | |
| RIFFLE | 158 | 977 | 1.2 | 0.06 | 1,172 | 78 | 11 | a | 40 | 31 | a | 1 | |
| RIFFLE W/ POCKETS | 34 | 341 | 1.3 | 0.10 | 438 | ao | 11 | 9 | 32 | 32 | 15 | 1 | |
| STEP/BEDROCK | 1 | 1 | 2.0 | 0.05 | 2 | 1 | 10 | 10 | 30 | 30 | 10 | 10 | |
| STEP/BOULDERS | 1 | 1 | 1.5 | 0.05 | 2 | 1 | 10 | 10 | 40 | 30 | 10 | 0 | |
| STEP/COBBLE | 2 | 1 | 1.3 | 0.05 | 10 | 10 | 15 | 10 | 40 | 25 | 10 | 0 | |
| STEP/LOG | 4 | 2 | 1.5 | 0.05 | 4 | 0 | 18 | 18 | 33 | 1a | 13 | 0 | |
| STEP/STRUCTURE | 3 | 4 | 2.7 | 0.02 | 9 | 0 | 4 | 0 | 2 | 3 | 7 | 0 | 30 |
| Total: | | 594 | 5,571 | 1.5 | 0.15 | 11,213 | 1158 | Avg:11 | 10 | 36 | 30 | 10 | 3 |

| HABITAT SUMMARY | | | | | | | | | |
|--------------------------|--------------|------------------------|---------------------|---------------------|----------------------------------|---------------------------|--|-------------|-----|
| Habitat Group | No. Units | Total Length (m) | Avg Width (m) | Avg Depth (m) | Uetted Area (m ²) | Large Boulders Percent | Large Boulders Number #/100m ² | Wood | |
| | | | | | | | | Class | |
| Dammed 8 BW Pools | 24 | 236 | 1.5 | 0.27 | 792 | 7.07 | 3 | 0.38 | 1.2 |
| Scour Pools | 189 | 789 | 1.5 | 0.26 | 1280 | 11.41 | 110 | 8.60 | 1.5 |
| Glides | 48 | 332 | 1.4 | 0.17 | 523 | 4.66 | 25 | 4.78 | 1.1 |
| Riffles | 192 | 1,318 | 1.2 | 0.07 | 1610 | 14.36 | 158 | 9.81 | 1.1 |
| Rapids | 57 | 265 | 1.4 | 0.05 | 363 | 3.24 | 67 | 1a.44 | 1.2 |
| Cascades | 3 | 19 | 1.2 | 0.20 | 25 | 0.22 | 3 | 12.10 | 1.3 |
| Step/Falls | 11 | 9 | 1.8 | 0.04 | 18 | 0.16 | 2 | 11.17 | 1.2 |
| Small Streams (SS) | 0 | 0 | . | . | 0 | 0.00 | 0 | 0.00 | . |
| Dry | 68 | 2,585 | 2.4 | 0.11 | 6572 | 58.61 | 790 | 12.02 | 1.1 |

Table D-10. Stream Summary for Moonshine Creek, RM 0.0 to 4.4, August 28-September 5, 1995.

| STREAM SUMMARY | | | | MOONSHINE CREEK | | | | | | | |
|-----------------|------------------------|---------------------|---------------------|------------------------------------|----------------------------------|---------|------|------|------|------|---------------------------|
| Number Units | Total Length (m) | Avg Width (m) | Avg Depth (m) | Total Area (m ²) | Substrate Percent Uetted Area | | | | | | Total Large Boulder |
| | | | | | S/O | Sand | Grvl | Cbbl | Bldr | Bdrk | |
| 594 | 5,571 | 1.5 | 0.15 | 11,213 | 11 | 10 | 36 | 30 | 10 | 3 | 1,158 |
| Uetted Area | | | | | | | | | | | |
| | | | | Habitat Group | (m ²) | Percent | | | | | |
| | | | | Scour Pool | 1,280 | 11.4 | | | | | |
| | | | | Backwater Pools | 792 | 7.1 | | | | | |
| | | | | Glide | 523 | 4.7 | | | | | |
| | | | | Riffle | 1,610 | 14.4 | | | | | |
| | | | | Rapid | 363 | 3.2 | | | | | |
| | | | | Cascade | 25 | 0.2 | | | | | |
| | | | | Step | 1a | 0.2 | | | | | |
| | | | | Dry | 6,572 | 58.6 | | | | | |

Table D-11. Valley, Channel, Bank and Wood **Summary** for Moonshine Creek, RM 0.0 to 4.4, August 28-September 5, 1995.

Valley and Channel Summary

Valley Characteristics (Percent Reach Length)

| <u>Narrow Valley Floor</u> | | <u>Broad Valley Floor</u> | |
|----------------------------|---|---------------------------|----|
| Steep V-shape | 0 | Constraining Terraces | 94 |
| Moderate V-shape | 0 | Multiple Terraces | 0 |
| Open V-shape | 0 | Wide Floodplain | 6 |

Valley Width **Index** avg: 10.0 range: 10.0-10.0

Channel Morphology (Percent Reach Length)

| <u>Constrained</u> | | <u>Unconstrained</u> | |
|--------------------|----|-------------------------|-----------|
| Hillslope | 0 | Single Channel | 48 |
| Bedrock | 0 | Multiple Channel | 0 |
| Terrace | 0 | Braided Channel | 0 |
| Alt. Terrace/Hill | 52 | | |
| Landuse | 0 | | |

Channel Characteristics

| <u>Type</u> | <u>Length</u> | <u>A r e a</u> | <u>Dry Units</u> |
|-------------|---------------|----------------|------------------|
| Primary | 5,351 | 10,980 | 68 |
| Secondary | 220 | 233 | 0 |

Channel Dimensions

| <u>Wetted Surface</u> | | <u>Active Channel</u> | | <u>First Terrace</u> | |
|-----------------------|-------------|-----------------------|-----|----------------------|-----|
| Width | 1.5 | Width | 5.1 | Width | 5.9 |
| Depth | 0.15 | Height | 0.5 | Height | 0.8 |
| W:D | 20.8 | | | | |

Stream **Flow** Type: LF **Water Temp:** 0.0-19.5
 Avg. Unit Gradient: 2.7 Habitat **Units/100m:** 10.7

Riparian, Bank, and Wood Summary

Land Use: AG,RR Riparian Veg.: D,S

Bank Stability

| <u>Bank Class</u> | <u>Percent Reach Length</u> | <u>Undercut Banks</u> |
|-----------------------|-----------------------------|----------------------------|
| Non-Erodible | 2.1 | Unit Average: 6.02% |
| Vegetation Stabilized | 91.5 | Open sky (% of 180) |
| Boulder-cobble | 0.3 | Unit Average: 44 |
| Actively Eroding | 6.0 | Range: 0-94 |

Large Woody Debris

| | | | |
|----------------------------------|-----|------------------------------|-----|
| Average Complexity Score: | 1.2 | | |
| Pieces | 63 | Volume(m³) | 34 |
| Pieces/100m | 1.2 | Volume/100m | 0.6 |

Table D-12. Riparian Summary for Moonshine Creek, RM 0.0 to 4.4, August 28-September 5, 1995.

REACH 0

REACH 0

RIPARIAN ZONE VEGETATION SUMMARY

Reach 0 is represented by 20 transects

Predominant **landform** in each zone

| | Zone 1 <u>0-10 meters</u> | Zone 2 <u>10-20 meters</u> | Zone 3 <u>20-30 meters</u> |
|-----------------------|-------------------------------------|--------------------------------------|--------------------------------------|
| Hillslope | 10 | 15 | 1a |
| High terrace | 53 | 50 | 60 |
| Lou terrace | 38 | 35 | 23 |
| Floodplain | 0 | 0 | 0 |
| Wetland/meadow | 0 | 0 | 0 |
| Stream channel | 0 | 0 | 0 |
| Roadbed/Railroad | 0 | 0 | 0 |
| Riprap | 0 | 0 | 0 |
| Surface slope (%) | 34 | 17 | 19 |

Canopy closure and ground cover

| | Zone 1 <u>0-10 meters</u> (%) | Zone 2 <u>10-20 meters</u> (%) | Zone 3 <u>20-30 meters</u> (%) |
|-------------------------|---|--|--|
| Canopy closure | 27 | 14 | 6 |
| Shrub cover | 48 | 43 | 40 |
| Grass/forb cover | 46 | 52 | 55 |

Average **number** of trees in a **5-meter** wide band

| Diameter class (cm) | Zone 1 <u>0-10 meters</u> | | Zone 2 <u>10-20 meters</u> | | Zone 3 <u>20-30 meters</u> | | Zones 1-3 <u>0-30 meters</u> | |
|-------------------------------|-------------------------------------|----------|--------------------------------------|----------|--------------------------------------|----------|--|----------|
| | Conifer | Hardwood | Conifer | Hardwood | Conifer | Hardwood | Conifer | Hardwood |
| 3-15cm | 0.0 | 4.0 | 0.1 | 2.1 | 0.0 | 0.6 | 0.1 | 6.7 |
| 15-30cm | 0.0 | 0.9 | 0.0 | 0.4 | 0.0 | 0.2 | 0.0 | 1.4 |
| 30-50cm | 0.0 | 1.2 | 0.0 | 0.2 | 0.0 | 0.1 | 0.0 | 1.4 |
| 50-90cm | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| >90cm | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Total/100m² | 0.0 | 6.1 | 0.1 | 2.7 | 0.0 | 0.8 | ***. * | 3.2 |

Table D-13. Surface Springs identified in the Habitat Survey, Moonshine Creek, **RM 0.0-4.4, 8/28-9/5, 1995.**

| UNIT NUMBER | UNIT TYPE | BANK SIDE | UNIT AREA (m ²) |
|--------------|-----------|----------------------------------|-----------------------------|
| 1 | GL | LEFT | 60 |
| 5 | SP | RIGHT | 14 |
| 11 | GL | LEFT | 40 |
| 13 | IP | RIGHT | 70 |
| 69 | LP | RIGHT | 3 |
| 100 | IP | RIGHT | 1 |
| 140 | LP | LEFT | 6 |
| 149 | SP | RIGHT | 3 |
| 159 | RI | RIGHT | 6 |
| 188 | IP | LEFT | 1 |
| 211 | LP | RIGHT | 7 |
| 214 | RI | RIGHT | 6 |
| 220 | RI | LEFT | 11 |
| 231 | IP | LEFT | 1 |
| 255 | RI | LEFT | 6 |
| 269 | LP | RIGHT | 11 |
| 277 | LP | RIGHT | 9 |
| 439 | IP | RIGHT | 70 |
| 449 | RE | RIGHT | 6 |
| 460 | PP | RIGHT | 7 |
| 476 | RP | LEFT | 8 |
| 510 | BP | RIGHT | 7 |
| 520 | RR | LEFT | 10 |
| 530 | GL | RIGHT | 4 |
| 553 | PD | LEFT | 30 |
| 580 | PD | RIGHT | 18 |
| 584 | PD | LEFT | 25 |
| TOTAL | 27 | 11 LEFT 16 RIGHT | 440 |

Table D-14. Habitat Unit Summary for Mission Creek, RM 0.0 to 4.3, August 15-September 11, 1995.

| REACH 0 | | | | | | | | | | | | |
|-----------------------|--------------|------------------|---------------|---------------|------------------------------|-------------------------|---------------|-----------|-----------|-----------|-----------|----------|
| HABITAT DETAIL | | | | | | | | | | | | |
| Habitat Type | Number Units | Total Length (m) | Avg Width (m) | Avg Depth (m) | Total Area (m ²) | Large Boulders (#>0.5m) | Substrate | | | | | |
| | | | | | | | Percent S/O | Uettd Snd | Grvl | Cbbl | Area Bldr | Bdrk |
| CULVERT CROSSING | 3 | 53 | 1.2 | 0.14 | 59 | 0 | 7 | 27 | 3 | 3 | 0 | 60 |
| DRY CHANNEL | 166 | 2,745 | 2.3 | 0.00 | 6,243 | 5 | 0 | 19 | 30 | 40 | 11 | 0 |
| DRY UNITS | 44 | 486 | 2.4 | 0.00 | 1,209 | 4 | 0 | 11 | 44 | 34 | 10 | 1 |
| GLIDE | 35 | 150 | 1.1 | 0.10 | 176 | 0 | a | 20 | 49 | 21 | 2 | 0 |
| POOL-BACKWATER | 20 | 29 | 0.4 | 0.08 | 16 | 0 | 14 | 49 | 31 | 4 | 0 | 2 |
| POOL-DAMMED | 6 | 22 | 0.8 | 0.17 | 19 | 0 | 10 | 35 | 43 | 12 | 0 | 0 |
| PWL-ISOLATED | 14 | 40 | 0.8 | 0.12 | 40 | 1 | 17 | 27 | 31 | 16 | 4 | 4 |
| POOL-LATERAL SCOUR | 148 | 515 | 1.0 | 0.19 | 552 | 3 | a | 22 | 47 | 21 | 2 | 1 |
| POOL-PLUNGE | 9 | 25 | 2.2 | 0.42 | 52 | 6 | 6 | 14 | 39 | 31 | a | 2 |
| POOL-STRAIGHT SCOUR | 78 | 248 | 1.0 | 0.18 | 260 | 7 | a | 22 | 47 | 19 | 4 | 1 |
| POOL-TRENCH | 10 | 51 | 1.0 | 0.40 | 54 | 0 | 7 | 20 | 33 | 1a | 3 | 20 |
| PUDDLED CHANNEL | 1a | 253 | 0.7 | 0.06 | 167 | 1 | 14 | 20 | 34 | 21 | 7 | 4 |
| RAPID/BEDROCK | 9 | 28 | 0.7 | 0.05 | 21 | 0 | a | 14 | 10 | 1 | 0 | 67 |
| RAPID/BOULDERS | 49 | 190 | 0.7 | 0.06 | 139 | 7 | 1 | 10 | 35 | 40 | 13 | 0 |
| RIFFLE | 232 | 945 | 1.2 | 0.05 | 852 | 1 | 3 | 16 | 57 | 21 | 2 | 0 |
| RIFFLE W/ POCKETS | 13 | 110 | 0.9 | 0.07 | 101 | 0 | 5 | 12 | 41 | 31 | 11 | 0 |
| STEP/BEDROCK | 1 | 2 | 0.4 | 0.03 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 100 |
| STEP/BOULDERS | 1 | 0 | 0.7 | 0.05 | 0 | 0 | 0 | 0 | 30 | 20 | 50 | 0 |
| STEP/COBBLE | 3 | 1 | 0.6 | 0.02 | 1 | 0 | 0 | 13 | 50 | 37 | 0 | 0 |
| STEP/LOG | 3 | 5 | 1.3 | 0.01 | 4 | 0 | 0 | 17 | 63 | 10 | 10 | 0 |
| STEP/STRUCTURE | 10 | 40 | 0.8 | 0.02 | 20 | 0 | 10 | 27 | 39 | 6 | 0 | 18 |
| Total: | 872 | 5,937 | 1.3 | 0.09 | 9,986 | 35 | Avg: 5 | 19 | 44 | 25 | 5 | 2 |

| HABITAT SUMMARY | | | | | | | | | |
|------------------------------|-----------|------------------|---------------|---------------|-------------------------------|------------------------|---|------------|-----|
| Habitat Group | No. Units | Total Length (m) | Avg Width (m) | Avg Depth (m) | Wetted Area (m ²) | Large Boulders Percent | Large Boulders Number #/100m ² | Wood Class | |
| | | | | | | | | | |
| Dammed & BW Pools | 40 | 90 | 0.6 | 0.11 | 75 | 0.75 | 1 | 1.33 | 1.7 |
| Scour Pools | 245 | a39 | 1.1 | 0.20 | 918 | 9.19 | 16 | 1.74 | 1.8 |
| Glides | 35 | 150 | 1.1 | 0.10 | 176 | 1.76 | 0 | 0.00 | 1.7 |
| Riffles | 245 | 1,055 | 1.2 | 0.05 | 953 | 9.55 | 1 | 0.10 | 1.3 |
| Rapids | 58 | 218 | 0.7 | 0.06 | 160 | 1.61 | 7 | 4.37 | 1.3 |
| Cascades | 0 | 0 | . | . | 0 | 0.00 | 0 | 0.00 | . |
| Step/Falls | 1a | 48 | 0.8 | 0.02 | 25 | 0.25 | 0 | 0.00 | 2.1 |
| Small Streams (SS) | 0 | 0 | . | . | 0 | 0.00 | 0 | 0.00 | . |
| Dry | 228 | 3,484 | 2.2 | *,** | 7619 | 76.29 | 10 | 0.13 | 1.6 |

Table D-15. Stream Summary for Mission Creek, RM 0.0 to 4.3, August 15-September 11, 1995.

| STREAM SUMMARY | | | | MISSION CREEK | | | | | | | |
|-----------------|------------------------|---------------------|---------------------|------------------------------------|----------------------------------|---------|------|------|------|---------------------------|----|
| Number Units | Total Length (m) | Avg Width (m) | Avg Depth (m) | Total Area (m ²) | Substrate Percent Uetted Area | | | | | Total Large Boulder | |
| | | | | | S/O | Sand | Grvl | Cbbl | Bldr | Bdrk | |
| a72 | 5,937 | 1.3 | 0.09 | 9,986 | 5 | 19 | 44 | 25 | 5 | 2 | 35 |
| Uetted Area | | | | | | | | | | | |
| | | | | Habitat Group | (m ²) | Percent | | | | | |
| | | | | Scour Pool | 918 | 9.2 | | | | | |
| | | | | Backwater Pools | 75 | 0.8 | | | | | |
| | | | | Glide | 176 | 1.8 | | | | | |
| | | | | R i f f l e | 953 | 9.5 | | | | | |
| | | | | Rapid | 160 | 1.6 | | | | | |
| | | | | Cascade | 0 | 0.0 | | | | | |
| | | | | Step | 25 | 0.3 | | | | | |
| | | | | Dry | 7,619 | 76.3 | | | | | |

Table D16. Valley, Channel, Bank and Wood Summary for Mission Creek, RM 0.0 to 4.3, August 15-September 11, 1995.

Valley and Channel **Summary**

Valley Characteristics (Percent Reach Length)

| <u>Narrow Valley Floor</u> | | <u>Broad Valley Floor</u> | |
|----------------------------|---|---------------------------|-----|
| Steep V-shape | 0 | Constraining Terraces | 100 |
| Moderate V-shape | 0 | Multiple Terraces | 0 |
| Open V-shape | 0 | Wide Floodplain | 0 |

Valley Width Index avg: 31.1 range: 1.0-100.0

Channel Morphology (Percent Reach Length)

| <u>Constrained</u> | | <u>Unconstrained</u> | |
|--------------------------|----|-------------------------|---|
| High slope | 0 | Single Channel | 0 |
| Bedrock | 0 | Multiple Channel | 0 |
| Terrace | 9 | Braided Channel | 0 |
| Alt. Terrace/Hill | 11 | | |
| Landuse | 0 | | |

Channel Characteristics

| <u>Type</u> | <u>Length</u> | <u>Area</u> | <u>Dry Units</u> |
|-------------|---------------|-------------|------------------|
| Primary | 5,757 | 9,759 | 228 |
| Secondary | 181 | 227 | 0 |

Channel Dimensions

| <u>Wetted Surface</u> | | <u>Active Channel</u> | | <u>First Terrace</u> | |
|-----------------------|------|-----------------------|-----|----------------------|-----|
| Width | 1.3 | Width | 3.2 | Width | 5.3 |
| Depth | 0.09 | Height | 0.4 | Height | 1.1 |
| W:D | 32.9 | | | | |

Stream **Flow** Type: LF Water **Temp**: 0.0-54.0
 Avg. Unit Gradient: 2.8 Habitat **Units/100m**: 14.7

Riparian, Bank, and Wood **Summary**

Land Use: **HG/RR** Riparian Veg.: D **30-50/S**

| <u>Bank Stability</u> | | | <u>Undercut Banks</u> | |
|-----------------------|----------------------|---------------|----------------------------|--------------|
| <u>Bank Class</u> | <u>Percent Reach</u> | <u>Length</u> | Unit Average: | 8.17% |
| Non-Erodible | | 1.7 | | |
| Vegetation Stabilized | 72.4 | | Open Sky (% of 180) | |
| Boulder-cobble | | 4.6 | Unit Average: | 38 |
| Actively Eroding | | 21.3 | Range: | 0-98 |

Large Woody Debris

| | | | |
|----------------------------------|------------|------------------------------|-----|
| Average Complexity Score: | 1.6 | | |
| Pieces | 378 | Volume(m³) | 93 |
| Pieces/100m | 6.6 | Volume/100m | 1.6 |

Table D17. Riparian Summary for Mission Creek, RM 0.0 to 4.3, August 15-September 11, 1995.

REACH 0

REACH 0

RIPARIAN ZONE VEGETATION SUMMARY

Reach 0 is represented by 36 transects

Predominant **landform** in each zone

| | zone1 0-10 meters | Zone 2 10-20 meters | zone3 20-30 meters |
|-------------------|----------------------|------------------------|-----------------------|
| Hillslope | 11 | 15 | 19 |
| High terrace | a9 | 85 | a1 |
| Lou terrace | 0 | 0 | 0 |
| Floodplain | 0 | 0 | 0 |
| Wetland/meadow | 0 | 0 | 0 |
| Stream channel | 0 | 0 | 0 |
| Roadbed/Railroad | 0 | 0 | 0 |
| Riprap | 0 | 0 | 0 |
| Surface slope (%) | 39 | 12 | 9 |

Canopy closure and ground cover

| | Zone 1 0-10 meters (%) | Zone 2 10-20 meters (%) | Zone 3 20-30 meters (%) |
|-------------------------|------------------------------|-------------------------------|-------------------------------|
| Canopy closure | 23 | a | 4 |
| Shrub cover | 33 | 15 | 7 |
| Grass/forb cover | 44 | 60 | 69 |

Average **number** of trees in a **5-meter** wide band

| Diameter class (cm) | Zone 1 0-10 meters | | Zone 2 10-20 meters | | Zone 3 20-30 meters | | zones 1-3 0-30 meters | |
|-------------------------------|-----------------------|----------|------------------------|----------|------------------------|--------|--------------------------|-----|
| | Conifer | Hardwood | Conifer | Hardwood | Boaiferd w o o d | | Hardwood | |
| 3-15cm | 0.1 | 6.7 | 0.1 | 0.1 | 0.0 | 0.0 | 0.2 | 6.8 |
| 15-30cm | 0.1 | 0.9 | 0.2 | 0.1 | 0.0 | ** . * | 0.2 | 1.0 |
| 30-50cm | 0.0 | 0.5 | 0.0 | 0.1 | 0.0 | 0.1 | 0.0 | 0.7 |
| 50-90cm | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | ** . * | 0.0 | 0.2 |
| >90cm | 0.0 | ** . * | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Total/100m² | 0.2 | 8.3 | 0.3 | 0.3 | 0.0 | 0.2 | 0.1 | 2.9 |

Table D-18. Surface Springs identified in the Habitat Survey, Mission Creek, RM 0.0-4.3, 8/15-9/11, 1995

| UNIT NUMBER | UNIT TYPE | BANK SIDE | UNIT AREA (m ²) |
|--------------|-----------|---------------------------------|-----------------------------|
| 29 | DP | RIGHT | 5 |
| 87 | IP | RIGHT | 1 |
| 247 | LP | RIGHT | 4 |
| 251 | BW | LEFT | 1 |
| 392 | PD | LEFT | 8 |
| 497 | LP | RIGHT | 9 |
| 559 | RB | LEFT | 6 |
| 578 | c c | RIGHT | 7 |
| 611 | LP | LEFT | 3 |
| 711 | RB | LEFT | 5 |
| 714 | LP | RIGHT | 4 |
| 742 | LP | LEFT | 6 |
| 748 | LP | RIGHT | 7 |
| 766 | LP | RIGHT | 4 |
| 774 | PP | RIGHT | 7 |
| 786 | LP | RIGHT | 5 |
| 796 | LP | RIGHT | 7 |
| 826 | SP | RIGHT | 1 |
| 849 | RP | RIGHT | 14 |
| 859 | SP | LEFT | 5 |
| 862 | PP | RIGHT | 10 |
| TOTAL | 21 | 7 LEFT 14 RIGHT | 119 |

Table D-19. Habitat Unit Summary for Cottonwood Creek, RM 0.0 to 4.1, June 20-August 1, 1995.

| REACH 0 | | | | | | | | | | | | REACH 0 | | | | | | | | | | | |
|---------------------|--------------|------------------|---------------|---------------|------------------------------|-------------------------|---------------|-------------|--------------|--------------|--------------|--------------|----|--|--|--|--|--|--|--|--|--|--|
| HABITAT DETAIL | | | | | | | | | | | | | | | | | | | | | | | |
| Habitat Type | Number Units | Total Length (m) | Avg Width (m) | Avg Depth (m) | Total Area (m ²) | Large Boulders (#>0.5m) | Substrate | | | | | | | | | | | | | | | | |
| | | | | | | | Percent S/O | Percent Snd | Percent Grvl | Percent Cbbl | Percent Bldr | Percent Bdrk | | | | | | | | | | | |
| CULVERT CROSSING | 4 | 26 | 1.0 | 0.24 | 26 | 100 | 1a | 38 | 5 | 0 | 0 | 0 | 15 | | | | | | | | | | |
| DRY UNITS | 113 | 2,205 | 3.1 | 0.00 | 6,759 | 282 | 1 | 10 | 26 | 38 | 23 | 2 | | | | | | | | | | | |
| GLIDE | 61 | 398 | 1.3 | 0.17 | 620 | 2 | 21 | 32 | 34 | 12 | 1 | 0 | | | | | | | | | | | |
| POOL-BACKWATER | 27 | 44 | 0.6 | 0.13 | 35 | 1 | 23 | 40 | 30 | 7 | 0 | 0 | | | | | | | | | | | |
| POOL-BEAVER DAM | 12 | 186 | 3.0 | 0.44 | 1,011 | 0 | 33 | 54 | 13 | 0 | 0 | 0 | | | | | | | | | | | |
| POOL-DAMMED | 16 | 100 | 1.7 | 0.25 | 198 | 0 | 17 | 49 | 29 | 4 | 0 | 0 | | | | | | | | | | | |
| POOL-ISOLATED | 23 | 357 | 1.6 | 0.20 | 1,346 | 5 | 26 | 27 | 30 | 13 | 2 | 3 | | | | | | | | | | | |
| POOL-LATERAL SCWR | 145 | 630 | 1.3 | 0.23 | 908 | 15 | 13 | 23 | 41 | 16 | 3 | 4 | | | | | | | | | | | |
| POOL-PLUNGE | 11 | 31 | 1.6 | 0.45 | 58 | 5 | 14 | 22 | 40 | 20 | 5 | 1 | | | | | | | | | | | |
| POOL-STRAIGHT SCOUR | 65 | 222 | 1.2 | 0.19 | 274 | 15 | 12 | 22 | 41 | 20 | 3 | 2 | | | | | | | | | | | |
| POOL-TRENCH | 4 | 10 | 1.3 | 0.29 | 12 | 0 | 13 | 13 | 5 | 0 | 0 | 70 | | | | | | | | | | | |
| PUDDLED CHANNEL | 36 | 537 | 1.1 | 0.06 | 826 | 19 | 21 | 16 | 31 | 21 | 10 | 1 | | | | | | | | | | | |
| RAPID/BEDROCK | 15 | 81 | 0.7 | 0.07 | 53 | 3 | 9 | 5 | a | 5 | 0 | 72 | | | | | | | | | | | |
| RAPID/BOULDERS | 34 | 176 | 1.0 | 0.07 | 198 | 32 | 1 | 9 | 26 | 42 | 20 | 1 | | | | | | | | | | | |
| RIFFLE | 304 | 2,344 | 1.1 | 0.08 | 2,846 | 30 | 7 | 21 | 49 | 19 | 4 | 1 | | | | | | | | | | | |
| RIFFLE W/ POCKETS | 16 | 189 | 1.2 | 0.10 | 232 | 13 | 10 | 16 | 38 | 25 | 10 | 1 | | | | | | | | | | | |
| STEP/BEDROCK | 2 | 2 | 0.9 | 0.06 | 2 | 0 | 10 | 0 | 0 | 0 | 0 | 90 | | | | | | | | | | | |
| STEP/BOULDERS | 3 | 1 | 0.8 | 0.04 | 1 | 0 | 7 | 13 | 10 | 23 | 47 | 0 | | | | | | | | | | | |
| STEP/COBBLE | 3 | 1 | 0.5 | 0.05 | 10 | 0 | 0 | 3 | 27 | 63 | 7 | 0 | | | | | | | | | | | |
| STEP/LOG | 3 | 1 | 1.1 | 0.03 | 1 | 0 | 27 | 40 | 33 | 0 | 0 | 0 | | | | | | | | | | | |
| STEP/STRUCTURE | 15 | 9 | 2.7 | 0.03 | 24 | 0 | 63 | 17 | a | 4 | 1 | a | | | | | | | | | | | |
| Total: | 912 | 7,547 | 1.4 | 0.12 | 15,431 | 522 | Avg:11 | 21 | 37 | 20 | 7 | 3 | | | | | | | | | | | |

| HABITAT SUMMARY | | | | | | | | | |
|--------------------|-----------|------------------|---------------|---------------|-------------------------------|-----------------------|------------------------------|------------|-----|
| Habitat Group | No. Units | Total Length (m) | Avg Width (m) | Avg Depth (m) | Uetted Area (m ²) | Large Boulders Number | Boulders #/100m ² | Uood Class | |
| Damned & BU Pools | 78 | 686 | 1.5 | 0.23 | 2590 | 16.79 | 6 | 0.23 | 1.9 |
| Scour Pools | 225 | 92 | 1.3 | 0.23 | 1252 | 8.11 | 35 | 2.80 | 1.9 |
| Glides | 61 | 398 | 1.3 | 0.17 | 620 | 4.02 | 2 | 0.32 | 1.5 |
| Riffles | 320 | 2,534 | 1.1 | 0.08 | 3078 | 19.95 | 43 | 1.40 | 1.4 |
| Rapids | 49 | 256 | 0.9 | 0.07 | 251 | 1.62 | 35 | 13.97 | 1.2 |
| Cascades | 0 | 0 | . | . | 0 | 0.00 | 0 | 0.00 | . |
| Step/Falls | 26 | 14 | 1.9 | 0.03 | 29 | 0.19 | 0 | 0.00 | 1.7 |
| Small Streams (SS) | 0 | 0 | . | . | 0 | 0.00 | 0 | 0.00 | . |
| Dry | 149 | 2,742 | 2.6 | 0.01 | 7585 | 49.16 | 301 | 3.97 | 1.1 |

Table D-20. Stream Summary for Cottonwood Creek, RM 0.0 to 4.1, June 20-August 1, 1995.

| STREAM SUMMARY | | | | COTTONWOOD CREEK | | | | | | | | |
|-----------------|--------------|-----------|-----------|-------------------|-------------------|---------|------|-----------|------|-------------|---------|--|
| Number Units | Total Length | Avg Width | Avg Depth | Total Area | Substrate | | | | | Total Large | | |
| | (m) | (m) | (m) | (m ²) | S/O | Sand | Grvl | Cbbl | Bldr | Bdrk | Boulder | |
| 912 | 7,547 | 1.4 | 0.12 | 15,431 | 11 | 21 | 37 | 20 | 7 | 3 | 522 | |
| Utted Area | | | | | | | | | | | | |
| | | | | Habitat Group | (m ²) | Percent | | | | | | |
| | | | | Scour Pool | 1,252 | a.1 | | | | | | |
| | | | | Backwater Pools | 2,590 | 16.8 | | | | | | |
| | | | | Glide | 620 | 4.0 | | | | | | |
| | | | | Riffle | 3,078 | 19.9 | | | | | | |
| | | | | Rapid | 251 | 1.6 | | | | | | |
| | | | | Cascade | 0 | 0.0 | | | | | | |
| | | | | Step | 29 | 0.2 | | | | | | |
| | | | | Dry | 7,585 | 49.2 | | | | | | |

Table D-21. Valley, Channel, Bank and Wood Summary for Cottonwood Creek, RM 0.0 to 4.1, June 20-August 1, 1995.

Valley and Channel **Summary**

Valley Characteristics (Percent Reach Length)

| <u>Narrow Valley Floor</u> | | <u>Broad Valley Floor</u> | |
|----------------------------|---|---------------------------|-----------|
| Steep V-shape | 0 | Constraining Terraces | 75 |
| Moderate V-shape | 0 | Multiple Terraces | 25 |
| Open V-shape | 0 | Wide Floodplain | 0 |

Valley Width Index avg: 19.6 range: 2.0-50.0

Channel Morphology (Percent Reach Length)

| <u>Constrained</u> | | <u>Unconstrained</u> | |
|--------------------------|-----------|----------------------|---|
| Hillslope | 0 | Single Channel | 0 |
| Bedrock | 0 | Multiple Channel | 0 |
| Terrace | 75 | Braided Channel | 0 |
| Alt. Terrace/Hill | 25 | | |
| Landuse | 0 | | |

Channel Characteristics

| <u>Type</u> | <u>Length</u> | <u>Area</u> | <u>Dry Units</u> |
|-------------|---------------|-------------|------------------|
| Primary | 7,018 | 13,999 | 149 |
| Secondary | 529 | 1,432 | 0 |

Channel Dimensions

| <u>Wetted Surface</u> | | <u>Active Channel</u> | | <u>First Terrace</u> | |
|-----------------------|--------------|-----------------------|-----|----------------------|-----|
| Width | 1.4 | Width | 3.6 | Width | 6.3 |
| Depth | 0.12 | Height | 0.3 | Height | 0.7 |
| U:D | ***.* | | | | |

Stream Flow Type: LF **Water Temp:** 12.0-21.0
 Avg. Unit Gradient: 3.3 Habitat **Units/100m:** 12.1

Riparian, Bank, and **Wood Summary**

Land Use: **HG,HG** Riparian Veg.: D **30-50,D 1**

Bank Stability

| <u>Bank Class</u> | <u>Percent Reach Length</u> | <u>Undercut Banks</u> |
|-----------------------|-----------------------------|----------------------------|
| Non-Erodible | 4.0 | Unit Average: 10.94% |
| Vegetation Stabilized | 76.4 | Open Sky (% of 180) |
| Boulder-cobble | 7.5 | Unit Average: 47 |
| Actively Eroding | 12.1 | Range: **-96 |

Large Woody Debris

| | | | |
|----------------------------------|------------|------------------------------|-----|
| Average Complexity Score: | 1.5 | | |
| Pieces | 236 | Volume(m³) | 61 |
| Pieces/100m | 3.4 | Volume/100m | 0.9 |

Table D-22. Riparian Summary for Cottonwood Creek, RM 0.0 to 4.1, June 20-August 1, 1995.

REACH 0 REACH 0
 RIPARIAN **ZONE** VEGETATION **SUMMARY**

Reach 0 is represented by 32 transects

Predominant **landform** in each zone

| | <u>Zone 1</u> 0-10 meters | <u>Zone 2</u> 10-20 meters | <u>Zone 3</u> 20-30 meters |
|-----------------------|------------------------------|-------------------------------|-------------------------------|
| Hillslope | 13 | 25 | 31 |
| High terrace | 72 | 70 | 66 |
| Lou terrace | 14 | 3 | 3 |
| Floodplain | 0 | 0 | 0 |
| Wetland/meadow | 0 | 0 | 0 |
| Stream channel | 0 | 0 | 0 |
| Roadbed/Railroad | 0 | 0 | 0 |
| Riprap | 0 | 0 | 0 |
| Surface slope (%) | 28 | 12 | 14 |

Canopy closure and **ground** cover

| | <u>Zone 1</u> 0-10 meters (%) | <u>Zone 2</u> 10-20 meters (%) | <u>Zone 3</u> 20-30 meters (%) |
|-------------------------|-------------------------------------|--------------------------------------|--------------------------------------|
| Canopy closure | 41 | 21 | 14 |
| Shrub cover | 33 | 29 | 21 |
| Grass/forb cover | 47 | 53 | 60 |

Average **number** of trees in a **5-meter** wide band

| Diameter class (cm) | <u>Zone 1</u> 0-10 meters | | <u>Zone 2</u> 10-20 meters | | <u>Zone 3</u> 20-30 meters | | <u>zones 1-3</u> 0-30 meters | |
|-------------------------------|------------------------------|----------|-------------------------------|----------|-------------------------------|----------|---------------------------------|------------|
| | Conifer | Hardwood | Conifer | Hardwood | Conifer | Hardwood | Conifer | Hardwood |
| 3-15cm | 3.0 | 13.1 | 0.4 | 4.3 | 0.3 | 1.6 | 3.7 | 19.0 |
| 15-30cm | 0.1 | 1.1 | ** | 0.6 | 0.1 | 0.2 | 0.2 | 1.8 |
| 30-50cm | 0.0 | 0.3 | 0.0 | 0.2 | 0.0 | 0.2 | 0.0 | 0.6 |
| 50-90cm | 0.0 | 0.1 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.3 |
| >90cm | 0.0 | 0.1 | 0.0 | ** | 0.0 | 0.0 | 0.0 | 0.1 |
| Total/100m² | 3.0 | 14.7 | 0.5 | 5.2 | 0.4 | 2.0 | 1.3 | 7.3 |

Table D-23. Surface Springs identified in the Habitat Survey, Cottonwood Creek, RM 0.0-4.1, 6/20-8/1, 1995

| UNIT NUMBER | UNIT TYPE | BANK SIDE | | UNIT AREA (m ²) |
|--------------|-----------|----------------|----------------|-----------------------------|
| 1 | GL | LEFT | | 14 |
| 7 | BP | LEFT | | 825 |
| 8 | IP | LEFT | | 1,200 |
| 9 | IP | LEFT | | 150 |
| 204 | RI | LEFT | | 129 |
| 246 | LP | LEFT | | 13 |
| 299 | LP | RIGHT | | 4 |
| 311 | RR | LEFT | | 4 |
| 316 | LP | LEFT | | 16 |
| 322 | RI | RIGHT | | 15 |
| 337 | PD | RIGHT | | 1 |
| 649 | LP | LEFT | | 2 |
| 662 | IP | LEFT | | 5 |
| 694 | LP | LEFT | | 1 |
| 724 | RI | LEFT | | 4 |
| 741 | PD | LEFT | | 17 |
| 773 | RB | BIGHT | | 2 |
| 776 | RI | LEFT | | 41 |
| 783 | RB | BIGHT | | 7 |
| 795 | PD | BIGHT | | 13 |
| 810 | IP | LEFT | | 3 |
| 843 | IP | BIGHT | | 1 |
| 886 | TP | LEFT | | 5 |
| TOTAL | 23 | 16 LEFT | 7 RIGHT | 2,472 |

Table D-24. Habitat Unit Summary for Coonskin Creek, RM 0.0 to 2.0, June 20-July 17, 1995.

| REACH 0 | | | | | | | | | | | | REACH 0 | | | | | | | | | | | |
|---------------------------|--------------|------------------|---------------|---------------|------------------------------|-------------------------|-------------|---------------|------|------|------|-------------|---|--|--|--|--|--|--|--|--|--|--|
| HABITAT DETAIL | | | | | | | | | | | | | | | | | | | | | | | |
| Habitat Type | Number Units | Total Length (m) | Avg Width (m) | Avg Depth (m) | Total Area (m ²) | Large Boulders (#>0.5m) | Substrate | | | | | | | | | | | | | | | | |
| | | | | | | | Percent S/O | Grvl | Cbbl | Bldr | Bdrk | Wetted Area | | | | | | | | | | | |
| CULVERT CROSSING | 1 | 23 | 0.6 | 0.05 | 14 | 0 | 10 | 10 | 30 | 30 | 20 | 0 | | | | | | | | | | | |
| DRY UNITS | 2 | a | 1.7 | 0.00 | 11 | 0 | 20 | 30 | 30 | 15 | 5 | 0 | | | | | | | | | | | |
| GLIDE | 14 | 133 | 2.3 | 0.23 | 385 | 14 | 21 | 26 | 34 | 14 | 3 | 1 | | | | | | | | | | | |
| POOL-ALCOVE | 1 | 76 | 1.7 | 0.35 | 130 | 0 | 30 | 60 | 10 | 0 | 0 | 0 | | | | | | | | | | | |
| POOL-BACKWATER | 14 | 30 | 0.9 | 0.15 | 33 | 3 | 20 | 39 | 29 | 9 | 2 | 0 | | | | | | | | | | | |
| POOL-DAMMED | 4 | 16 | 1.3 | 0.20 | 19 | 0 | 1a | 38 | 33 | 13 | 0 | 0 | | | | | | | | | | | |
| POOL-ISOLATED | 2 | 19 | 1.4 | 0.38 | 22 | 0 | 20 | 45 | 25 | 10 | 0 | 0 | | | | | | | | | | | |
| POOL-LATERAL SCOUR | 126 | 531 | 1.3 | 0.26 | 776 | 19 | 12 | 21 | 34 | 24 | 7 | 1 | | | | | | | | | | | |
| POOL-PLUNGE | 27 | 65 | 2.0 | 0.39 | 134 | 14 | 14 | 23 | 27 | 20 | 10 | 4 | | | | | | | | | | | |
| POOL-STRAIGHT SCWR | 109 | 393 | 1.4 | 0.25 | 587 | 47 | 13 | 21 | 33 | 23 | a | 1 | | | | | | | | | | | |
| POOL-TRENCH | 7 | 23 | 1.2 | 0.55 | 29 | 1 | 10 | 23 | 16 | 11 | 6 | 34 | | | | | | | | | | | |
| RAPID/BEDROCK | 7 | 47 | 1.3 | 0.09 | 57 | 1 | 10 | 13 | 13 | 6 | 1 | 57 | | | | | | | | | | | |
| RAPID/BOULDERS | 48 | 264 | 1.5 | 0.08 | 422 | 55 | 11 | 13 | 33 | 27 | 16 | 0 | | | | | | | | | | | |
| RIFFLE | 171 | 1,629 | 1.4 | 0.08 | 2,240 | 55 | 11 | 16 | 41 | 24 | a | 1 | | | | | | | | | | | |
| RIFFLE W/ POCKETS | 62 | 726 | 1.3 | 0.13 | 977 | a7 | 11 | 16 | 31 | 28 | 13 | 1 | | | | | | | | | | | |
| STEP/BEDROCK | 11 | 9 | 1.4 | 0.05 | 12 | 1 | 19 | 13 | a | 6 | a | 45 | | | | | | | | | | | |
| STEP/BOULDERS | 6 | 2 | 1.2 | 0.05 | 2 | 10 | 10 | 17 | 32 | 25 | 17 | 0 | | | | | | | | | | | |
| STEP/COBBLE | 1 | 0 | 0.5 | 0.05 | 0 | 0 | 20 | 20 | 10 | 20 | 10 | 20 | | | | | | | | | | | |
| STEP/LOG | a | 4 | 0.9 | 0.09 | 3 | 0 | 18 | 24 | 36 | 21 | 1 | 0 | | | | | | | | | | | |
| STEP/STRUCTURE | 5 | 3 | 2.1 | 0.06 | 7 | 0 | 26 | 24 | 28 | 16 | 4 | 2 | | | | | | | | | | | |
| Total: | | 626 | 4,001 | 1.4 | 0.18 | 5,860 | 307 | Avg:12 | 19 | 34 | 23 | 9 | 3 | | | | | | | | | | |

| HABITAT SUMMARY | | | | | | | | | |
|------------------------------|-----------|------------------|---------------|---------------|-------------------------------|-----------------------|--------------------------|------------|--|
| Habitat Group | No. Units | Total Length (m) | Avg Width (m) | Avg Depth (m) | Wetted Area (m ²) | Large Boulders Number | Wood #/100m ² | Wood Class | |
| | | | | | | | | | |
| Dammed & BU Pools | 21 | 141 | 1.0 | 0.19 | 204 | 3 | 1.47 | 1.7 | |
| Scour Pools | 269 | 1,012 | 1.4 | 0.28 | 1526 | a1 | 5.31 | 1.7 | |
| Glides | 14 | 133 | 2.3 | 0.23 | 385 | 14 | 3.64 | 1.5 | |
| Riffles | 233 | 2,354 | 1.4 | 0.10 | 3217 | 142 | 4.41 | 1.4 | |
| Rapids | 55 | 311 | 1.5 | 0.08 | 480 | 56 | 11.68 | 1.3 | |
| Cascades | 0 | 0 | . | . | 0 | 0.00 | 0.00 | . | |
| Step/Falls | 31 | 19 | 1.3 | 0.06 | 25 | 0.43 | 43.65 | 1.2 | |
| Small Streams (SS) | 0 | 0 | . | . | 0 | 0.00 | 0.00 | . | |
| Dry | 2 | a | 1.7 | 0.00 | 11 | 0.19 | 0.00 | 1.0 | |

Table D-25. Stream Summary for Coonskin Creek, RM 0.0 to 2.0, June 20-July 17, 1995.

| STREAM SUMMARY | | | | COONSKIN CREEK | | | | | | | |
|-----------------|------------------------|------------------------|---------------------|------------------------------------|----------------------------------|------|------|------|------|---------------------------|-----|
| Number Units | Total Length (m) | Avg Width (m) | Avg Depth (m) | Total Area (m ²) | Substrate Percent Vetted Area | | | | | Total Large Boulder | |
| | | | | | S/O | Sand | Grvl | Cbbl | Bldr | Bdrk | |
| 626 | 4,001 | 1.4 | 0.18 | 5,860 | 12 | 19 | 34 | 23 | 9 | 3 | 307 |
| Vetted Area | | | | | | | | | | | |
| | | Habitat Group | (m ²) | Percent | | | | | | | |
| | | Scour Pool | 1,526 | 26.0 | | | | | | | |
| | | Backwater Pools | 204 | 3.5 | | | | | | | |
| | | Glide | 385 | 6.6 | | | | | | | |
| | | Riffle | 3,217 | 54.9 | | | | | | | |
| | | Rapid | 480 | a.2 | | | | | | | |
| | | Cascade | 0 | 0.0 | | | | | | | |
| | | Step | 25 | 0.4 | | | | | | | |
| | | Dry | 11 | 0.2 | | | | | | | |

Table D-26. Valley, Channel, Bank and Wood Summary for Coonskin Creek, RM 0.0 to 2.0, June 20-July 17, 1995.

Valley and Channel Summary

Valley Characteristics (Percent Reach Length)

| Narrow Valley Floor | | Broad Valley Floor | |
|---------------------|---|-----------------------|-----|
| Steep V-shape | 0 | Constraining Terraces | 100 |
| Moderate V-shape | 0 | Multiple Terraces | 0 |
| Open V-shape | 0 | Wide Floodplain | 0 |

Valley Width Index avg: 11.5 range: 10.0-50.0

Channel Morphology (Percent Reach Length)

| Constrained | | Unconstrained | |
|-------------------|-----|------------------|---|
| Hillslope | 0 | Single Channel | 0 |
| Bedrock | 0 | Multiple Channel | 0 |
| Terrace | 100 | Braided Channel | 0 |
| Alt. Terrace/Hill | 0 | | |
| Landuse | 0 | | |

Channel Characteristics

| Type | Length | Area | Dry Units |
|-----------|--------|-------|-----------|
| Primary | 3,496 | 5,299 | 1 |
| Secondary | 505 | 561 | 1 |

Channel Dimensions

| Wetted Surface | | Active Channel | | First Terrace | |
|----------------|------|----------------|-----|---------------|-----|
| Width | 1.4 | Width | 3.5 | Width | 5.7 |
| Depth | 0.18 | Height | 0.4 | Height | 0.8 |
| W:D | 19.2 | | | | |

Stream Flow Type: MF Water Temp: 12.5-21.0
 Avg. Unit Gradient: 3.1 Habitat Units/100m: 15.6

Riparian, Bank, and Wood Summary

Land Use: AG, LG Riparian Veg.: S, G

Bank Stability

| Bank Class | Percent Reach Length | Undercut Banks |
|-----------------------|----------------------|----------------------|
| Non-Erodible | 2.0 | Unit Average: 11.23% |
| Vegetation Stabilized | 83.8 | Open Sky (% of 180) |
| Boulder-cobble | 0.5 | Unit Average: 41 |
| Actively Eroding | 13.2 | Range: 0-92 |

Large Woody Debris

| Average Complexity Score: 1.5 | | | |
|-------------------------------|-----|-------------------------|-----|
| Pieces | 55 | Volume(m ³) | 43 |
| Pieces/100m | 1.6 | Volume/100m | 1.2 |

Table D-27. Riparian Summary for Coonskin Creek, RM 0.0 to 2.0, June 20-July 17, 1995.

REACH 0 REACH 0
 RIPARIAN **ZONE VEGETATION SUMMARY**

Reach 0 is represented by 23 transects

Predominant **landform** in each zone

| | zone 1 <u>0-10 meters</u> | zone2 <u>10-20 meters</u> | Zone3 <u>20-30 meters</u> |
|-----------------------|------------------------------|------------------------------|------------------------------|
| Hillslope | 2 | 4 | 9 |
| High terrace | 43 | 44 | 50 |
| Low terrace | 55 | 49 | 39 |
| Floodplain | 0 | 0 | 0 |
| Wetland/meadow | 0 | 0 | 0 |
| Stream channel | 0 | 2 | 2 |
| Roadbed/Railroad | 0 | 0 | 0 |
| Riprap | 0 | 0 | 0 |
| Surface slope (%) | 32 | 17 | 19 |

Canopy closure and ground cover

| | Zone 1 <u>0-10 meters</u> (%) | zone 2 <u>10-20 meters</u> (%) | Zone3 <u>20-30 meters</u> (%) |
|------------------|-------------------------------------|--------------------------------------|-------------------------------------|
| Canopy closure | 31 | 22 | 15 |
| Shrub cover | 51 | 42 | 35 |
| Grass/forb cover | 44 | 51 | 53 |

Average **number** of trees in a **5-meter** wide band

| Diameter class (cm) | Zone 1 <u>0-10 meters</u> | | zone 2 <u>10-20 meters</u> | | Zone3 <u>20-30 meters</u> | | zones 1-3 <u>0-30 meters</u> | |
|-------------------------------|------------------------------|----------|-------------------------------|----------|------------------------------|----------|---------------------------------|----------|
| | Conifer | Hardwood | Conifer | Hardwood | Conifer | Hardwood | Conifer | Hardwood |
| 3-15cm | 0.0 | 3.3 | **.* | 1.8 | 0.0 | 1.0 | **.* | 6.2 |
| 15-30cm | 0.0 | 0.7 | **.* | **.* | **.* | 0.2 | 0.1 | 0.9 |
| 30-50cm | 0.0 | 0.6 | **.* | 0.5 | 0.0 | 0.1 | 0.0 | 1.2 |
| 50-90cm | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| >90cm | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Total/100m² | 0.0 | 4.7 | 0.1 | 2.3 | ***.* | 1.3 | 0.1 | 2.8 |

Table D-28. Surface Springs identified in the Habitat Survey, Coonsk Creek, RM 0.0-2.0, 6/20-7/17, 1995.

| UNIT NUMBER | UNIT TYPE | BANK SIDE | | UNIT AREA (m ²) |
|--------------|-----------|----------------|----------------|-----------------------------|
| 11 | IP | LEFT | | 15 |
| 87 | LP | RIGHT | | 3 |
| 92 | RI | LEFT | | 4 |
| 137 | RJ | LEFT | | 15 |
| 179 | LP | LEFT | | 3 |
| 216 | LP | LEFT | | 4 |
| 221 | LP | RIGHT | | 5 |
| 263 | RP | LEFT | | 13 |
| 268 | SP | RIGHT | | 11 |
| 405 | LP | LEFT | | 5 |
| 487 | LP | RIGHT | | 3 |
| 498 | LP | RIGHT | | 2 |
| 531 | RI | RIGHT | | 2 |
| 548 | TP | LEFT | | 5 |
| 602 | SP | LEFT | | 5 |
| 621 | RP | RIGHT | | 19 |
| 625 | RB | LEFT | | 24 |
| TOTAL | 17 | 10 LEFT | 7 RIGHT | 138 |

APPENDIX E
Biological Survey Data Summary Tables and Figures

Table E-1. Mean Density and Population Estimate of Rainbow/Steelhead and Bull Trout, Chinook Salmon, and Mountain Whitefish. Umatilla River, RM 81.8-89.6, 8/8-8/25, 1995.

| HABITAT TYPE | # OF UNITS | # UNITS SAMPLED | % OF TOTAL SAMPLED | AREA-M ² | AREA-M ² SAMPLED | % OF AREA SAMPLED | MEAN SALMONID DENSITY | EST. # SALMONIDS |
|-----------------------------|-------------|-----------------|--------------------|---------------------|-----------------------------|-------------------|-----------------------|------------------|
| POOLS | | | | | | | | |
| Plunge Pool | 3 | 1 | 33.3 | 250 | 165 | 66.0 | 0.9515 | 238 |
| Scour Pool | 63 | 8 | 12.7 | 14,201 | 1,057 | 7.4 | 0.4541 | 6,449 |
| Lateral Pool | 108 | 11 | 10.2 | 23,629 | 364 | 1.5 | 0.8709 | 20,578 |
| Dammed Pool | 5 | 1 | 20.0 | 680 | 26 | 3.8 | 1.7692 | 1,203 |
| Beaver Dam Pool | 1 | 0 | 0.0 | 519 | 0 | 0.0 | | |
| SUBUNIT POOLS | | | | | | | | |
| Back Water Pool | 42 | 5 | 11.9 | 755 | 87 | 11.5 | 0.9080 | 606 |
| Isolated Pool | 14 | 1 | 7.1 | 1,657 | 43 | 2.6 | 0.0930 | 154 |
| Isolated Pool w/ss | 10 | 7 | 70.0 | 2,983 | 2,604 | 87.3 | 0.0545 | 163 |
| Unclass. Isolated Pool w/ss | 9* | 9 | 100.0 | 2,053 | 2,053 | 100.0 | 0.1495 | 307 |
| Puddled | 6 | 2 | 33.3 | 461 | 63 | 13.7 | 0.1111 | 51 |
| GLIDES | | | | | | | | |
| Glide | 63 | 8 | 12.7 | 13,871 | 1,178 | a.5 | 0.1469 | 2,037 |
| Subtotal | 324 | | 16.4 | 61,059 | 7,640 | 12.5 | 0.5219 | 31,866 |
| RIFFLES | | | | | | | | |
| Riffle | 206 | | 4.4 | 60,403 | 1,228 | 2.0 | 0.3461 | 20,905 |
| Riffle With Pockets | 47 | | 8.5 | 22,653 | 732 | 3.2 | 0.5137 | 11,636 |
| RAPIDS | | | | | | | | |
| Rapid-Boulder | 63 | | 9.5 | 9,614 | 635 | 6.6 | 0.4898 | 4,709 |
| Rapid-Bedrock | 3 | | 0.0 | 131 | 0 | 0.0 | | |
| Subtotal | 319 | | 6.0 | 92,801 | 2,595 | 2.8 | 0.4614 | 37,250 |
| SPECIAL CASES | | | | | | | | |
| Steps | 4 | | 0.0 | 120 | 0 | 0.0 | | |
| Dry | | 0 | 0.0 | | | 0.0 | | 0 |
| Subtotal | 5 | 0 | 0.0 | 144 | 0 | 0.0 | 0 | 0 |
| TOTALS | 648* | 72 | 11.1 | 154,004 | 10,235 | 6.7 | 0.4488 | 69,116 |

The physical properties of Steps, and Dry Units prevented sampling.

* Includes 9 units unclassified during the habitat survey, but identified during the biological survey.

† Was not sampled because the habitat type could not be sampled effectively or accurately.

Table E-2. Mean Density and Population Estimate of Natural **Rainbow/Steelhead** Trout, Chinook and Coho Salmon, Moonshine Creek, RM **0.0-4.4, 9/18-9/21, 1995.**

| HABITAT TYPE | # OF UNITS | # UNITS SAMPLED | % OF TOTAL SAMPLED | AREA-M ² | AREA-M ² SAMPLED | % OF AREA SAMPLED | MEAN SALMONID DENSITY | EST. # SALMONIDS |
|----------------------|------------|-----------------|--------------------|---------------------|-----------------------------|-------------------|-----------------------|------------------|
| POOLS | | | | | | | | |
| Plunge Pool | 9 | 6 | 66.7 | 75 | 64 | 80.0 | 2.2188 | 166 |
| Scour Pool | 68 | 13 | 19.1 | 467 | 171 | 36.6 | 0.4795 | 224 |
| Lateral Pool | 110 | 18 | 16.4 | 729 | 135 | 18.5 | 0.4667 | 340 |
| Trench Pool | 2 | 1 | 50.0 | 8 | 7 | 87.5 | 1.8571 | 15 |
| Beaver Dam Pool | 3 | 1 | 33.3 | 612 | 31 | 5.1 | 0.1290 | 79 |
| SUBUNIT POOLS | | | | | | | | |
| Back Water Pool | 11 | 2 | 18.2 | 11 | 9 | 81.8 | 0.1111 | 1 |
| Isolated Pool | 10 | 1 | 10.0 | 170 | 2 | 1.2 | 0.0000 | 0 |
| Puddled | 13 | 2 | 15.4 | 376 | 167 | 44.4 | 0.0599 | 23 |
| GLIDES | | | | | | | | |
| Glide | 48 | 10 | 20.8 | 528 | 157 | 29.7 | 0.0764 | 40 |
| Subtotal | 274 | 54 | 19.7 | 2,976 | 743 | 25.0 | 0.2984 | 888 |
| RIFFLES | | | | | | | | |
| Rime | 158 | 13 | 8.2 | 1,172 | 156 | 13.3 | 0.0385 | 45 |
| Riffle With Pockets | 34 | 8 | 23.5 | 438 | 100 | 22.8 | 0.4800 | 210 |
| RAPIDS | | | | | | | | |
| Rapid-Boulder | 48 | 7 | 14.6 | 3% | 46 | 15.0 | 0.0435 | 13 |
| Rapid-Bedrock | 9 | 4 | 44.4 | 58 | 36 | 62.1 | 0.0556 | 3 |
| CASCADES | | | | | | | | |
| Cascade-Bedrock | 3 | 3 | 100.0 | 25 | 23 | 92.0 | 0.3913 | 10 |
| Subtotal | 252 | 35 | 13.9 | 1,999 | 361 | 18.1 | 0.1406 | 281 |
| SPECIAL CASES | | | | | | | | |
| Steps | 11 | 0 | 0.0 | 18 | 0 | 0.0 | - | - |
| Culvert Crossing | 2 | 0 | 0.0 | 31 | 0 | 0.0 | | |
| Dry | 55 | 0 | 0.0 | 6.1% | 0 | 0.0 | 0 | 0 |
| Subtotal | 68 | 0 | 0.0 | 6.245 | 0 | 0.0 | 0 | 0 |
| TOTALS | 594 | 89 | 15.0 | 11,213 | 1,104 | 9.9 | 0.1042 | 1,169 |

The physical properties of Steps, Dry Units, and Culvert Crossings prevented sampling.

Table E-3. Mean Density and Population Estimate of Rainbow/Steelhead Trout and Coho Salmon, Mission Creek, RM 0.043, 9/5-9/13, 1995.

| HABITAT TYPE | # OF UNITS | # UNITS SAMPLED | % OF TOTAL SAMPLED | AREA-M ² | AREA-M ² SAMPLED | % OF AREA SAMPLED | MEAN SALMONID DENSITY | EST. # SALMONIDS |
|----------------------|------------|-----------------|--------------------|---------------------|-----------------------------|-------------------|-----------------------|------------------|
| POOLS | | | | | | | | |
| Plunge Pool | 9 | 6 | 66.7 | 52 | 47 | 90.4 | 1.6170 | 84 |
| scour Pool | 78 | 7 | 9.0 | 260 | 42 | 16.2 | 0.7857 | 204 |
| Lateral Pool | 148 | 18 | 12.2 | 552 | 73 | 13.2 | 0.8356 | 461 |
| Trench Pool | 10 | 4 | 40.0 | 54 | 27 | 50.0 | 1.0370 | 56 |
| Dammed Pool | 6 | 2 | 33.3 | 19 | 12 | 63.2 | 0.3333 | 6 |
| SUBUNIT POOLS | | | | | | | | |
| Back Water Pool | 20 | 3 | 15.0 | 16 | 4 | 25.0 | 0.2500 | 4 |
| Isolated Pool | 14 | 2 | 14.3 | 40 | 9 | 22.5 | 0.0000 | 0 |
| Puddled | 18 | 0 | 0.0 | 167 | 0 | 0.0 | | |
| GLIDES | | | | | | | | |
| Glide | 35 | 4 | 11.4 | 176 | 41 | 23.3 | 0.3659 | 64 |
| Subtotal | 338 | 46 | 13.6 | 1,336 | 255 | 19.1 | 0.6579 | 879 |
| RIFFLES | | | | | | | | |
| Riffle | 232 | 10 | 4.3 | 852 | 114 | 13.4 | 0.0351 | 30 |
| Riffle With Pockets | 13 | 4 | 30.8 | 101 | 48 | 47.5 | 0.2083 | 21 |
| RAPIDS | | | | | | | | |
| Rapid-Boulder | 49 | 5 | 10.2 | 139 | 22 | 15.8 | 0.0000 | 0 |
| Rapid-Bedrock | 9 | 0 [*] | 0.0 | 21 | 0 | | 0.0 | - |
| Subtotal | 303 | 19 | 6.3 | 1,113 | 184 | 16.5 | 0.0458 | 51 |
| SPECIAL CASES | | | | | | | | |
| Steps | 18 | 0 | 0.0 | 26 | 0 | 0.0 | | |
| Culvert Crossing | 3 | 0 [^] | 0.0 | 59 | 0 | 0.0 | | |
| Dry | 210 | 0 [^] | 0.0 | 7,452 | 0 | 0.0 | .0000 | 0 |
| Subtotal | 231 | 0 | 0.0 | 7,537 | 0 | 0.0 | .0000 | 0 |
| TOTALS | 872 | 65 | 7.5 | 9,986 | 440 | 4.4 | 0.0931 | 930 |

[^] The physical properties of Steps, Dry Umts, and Culvert Crossings prevented sampling,

^{*} Was not sampled because habitat was not suitable for salmonids.

^{*} Was not sampled because habitat type could not be sampled effectively or accurately.

Table E-4. Mean Density and Population Estimate of Natural Rainbow/Steelhead Trout, Chinook and Coho Salmon, Cottonwood Creek, IRM 0.041, 7/5-8/1, 1995.

| HABITAT TYPE | # OF UNITS | # UNITS SAMPLED | % OF TOTAL SAMPLED | AREA-M ² | AREA-M ² SAMPLED | % OF AREA SAMPLED | MEAN SALMONID DENSITY | EST. # SALMONIDS |
|----------------------|------------|-----------------|--------------------|---------------------|-----------------------------|-------------------|-----------------------|------------------|
| POOLS | | | | | | | | |
| Plung Pool | 11 | 6 | 54.5 | 58 | 48 | 82.8 | 2.5000 | 145 |
| scour Pool | 65 | 13 | 20.0 | 274 | 69 | 25.2 | 0.2319 | 64 |
| Lateral Pool | 145 | 14 | 9.7 | 908 | 118 | 13.0 | 0.1949 | 177 |
| Trench Pool | 4 | 1 | 25.0 | 12 | 4 | 33.4 | 1.0000 | 12 |
| Dammed Pool | 16 | 3 | 18.8 | 198 | 48 | 24.2 | 0.1250 | 15 |
| Beaver Dam Pool | 12 | 2 | 16.7 | 1,011 | 796 | 78.7 | 0.0000 | 0 |
| SUBUNIT POOLS | | | | | | | | |
| Back Water Pool | 27 | 1 | 3.7 | 35 | 1 | 2.9 | 0.0000 | 0 |
| Isolated Pool | 23 | 3 | 13.0 | 1,346 | 1,143 | 84.9 | 0.0367 | 49 |
| Puddled | 36 | 1 | 2.8 | 826 | 7 | 0.8 | 0.0000 | 0 |
| GLIDES | | | | | | | | |
| Glide | 61 | 8 | 13.1 | 620 | 141 | 22.7 | 0.0355 | 22 |
| Subtotal | 400 | 52 | 13.0 | 5,288 | 2,375 | 44.9 | 0.0915 | 484 |
| RIFFLES | | | | | | | | |
| Riffle | 304 | 12 | 3.9 | 2,846 | 417 | 14.7 | 0.0312 | 89 |
| Riffle With Pockets | 16 | 2 | 12.5 | 232 | 23 | 9.9 | 0.1304 | 30 |
| RAPIDS | | | | | | | | |
| Rapid-Boulder | 34 | 3 | 8.8 | 1% | 26 | 13.1 | 0.1154 | 23 |
| Rapid-Bedrock | 15 | 1 | 6.7 | 53 | 5 | 9.4 | 0.0000 | 0 |
| Subtotal | 369 | 18 | 4.9 | 3,329 | 471 | 14.1 | 0.0427 | 142 |
| SPECIAL CASES | | | | | | | | |
| Steps | 26 | 0 | 0.0 | 29 | 0 | 0.0 | | |
| Culvert Crossing | 4 | 0 | 0.0 | 26 | 0 | 0.0 | | |
| Dry | 113 | 0 | 0.0 | 6,759 | 0 | 0.0 | 0.0000 | 0 |
| Subtotal | 143 | 0 | 0.0 | 6,814 | 0 | 0.0 | 0.0000 | 0 |
| TOTALS | 912 | 70 | 7.7 | 15,431 | 2,824 | 18.3 | 0.0406 | 626 |

The physical properties of Steps, Dry Units, and Culvert Crossings prevented sampling.

Table E-5. Population Density Estimate of **Rainbow/Steelhead** Trout, Chinook and **Coho** Salmon, Coonskin Creek, RM 0.0-2.0, **6/29-7/18, 1995.**

| HABITAT TYPE | # OF UNITS | # UNITS SAMPLED | % OF TOTAL SAMPLED | AREA-M ² | AREA-M ² SAMPLED | % OF AREA SAMPLED | MEAN SALMONID DENSITY | EST. # SALMONIDS |
|----------------------|------------|-----------------|--------------------|---------------------|-----------------------------|-------------------|-----------------------|------------------|
| POOLS | | | | | | | | |
| Plunge Pool | 27 | 9 | 33.3 | 134 | 56 | 41.8 | 1.6964 | 227 |
| scow Pool | 109 | 16 | 14.7 | 587 | 94 | 16.0 | 1.1277 | 662 |
| Lateral Pool | 126 | 19 | 15.1 | 144 | | 18.6 | 0.5000 | 388 |
| Trench Dammed Pool | 7 | 5 | 71.4 | 29 | 20 | 69.0 | 4.8000 | 139 |
| | 4 | 2 | 50.0 | 19 | 8 | 42.1 | 0.0000 | 0 |
| SUBUNIT POOLS | | | | | | | | |
| Alcove | 1 | 0* | 0.0 | 130 | 0 | 0.0 | - | - |
| Back Water Pool | 14 | 1 | 7.1 | 33 | 1 | 0.1 | 0.0000 | 0 |
| Isolated Pool (IP) | 2 | 2 | 100.0 | 22 | 20 | 0.9 | 0.0000 | 0 |
| GLIDES | | | | | | | | |
| Glide | 14 | 5 | 35.7 | 385 | 171 | 44.4 | 0.0877 | 34 |
| Subtotal | 304 | 57 | 18.8 | 2,115 | 514 | 24.3 | 0.6856 | 1,450 |
| RIFFLES | | | | | | | | |
| Riffle | 171 | 12 | 7.0 | 2,240 | 130 | 0.1 | 0.0846 | 190 |
| Riffle With Pockets | 62 | 6 | 9.7 | 977 | 135 | 13.8 | 0.1555 | 152 |
| RAPIDS | | | | | | | | |
| Rapid-Boulder | 48 | 9 | 9.7 | 422 | 104 | 24.6 | 0.1731 | 73 |
| Rapid-Bedrock | 7 | 2 | 28.6 | 57 | 22 | 38.6 | 0.1818 | 10 |
| Subtotal | 288 | 29 | 10.1 | 3,696 | 391 | 10.6 | 0.1150 | 425 |
| SPECIAL CASES | | | | | | | | |
| Steps | 31 | 0 | 0.0 | 24 | 0 | 0.0 | | |
| Culvert Crossing | 1 | 0 | 0.0 | 14 | 0 | 0.0 | | |
| Dry | 2 | 0* | 0.0 | 11 | 0 | 0.0 | 0.0000 | 0 |
| subtotal | 34 | 0 | 0.0 | 49 | 0 | 0.0 | 0.0808 | 0 |
| TOTALS | 626 | 88 | 14.1 | 5,860 | 905 | 15.4 | 0.3200 | 1,875 |

* The physical properties of Steps, Dry Units, and Culvert Crossings prevented sampling.

" Was not sampled because the habitat was not suitable for salmonids.

Table E-6. Mean Density and Population Estimate of **Rainbow/Steelhead** and Bull Trout, Umatilla River, RM 81.8-89.6, **8/8-8/25**, 1995.

| HABITAT TYPE | TOTAL AREA SAMPLED/M ² | AREA/M ² W/SPP. PRESENT | MEAN DENSITY IN TOTAL AREA | EST# IN UNIT |
|--|-----------------------------------|------------------------------------|----------------------------|--------------|
| Natural Rainbow/Steelhead Trout | | | | |
| Plunge Pool | 165 | 165 | .9515 | 157 |
| Lateral Pool | 364 | 364 | .7967 | 290 |
| Backwater Pool | 87 | 78 | .7126 | 62 |
| Riffle With Pockets | 732 | 732 | .4481 | 328 |
| Rapid Over Boulders | 635 | 635 | .4000 | 254 |
| Dammed Pool | 26 | 26 | .3846 | 10 |
| scour Pool | 1,057 | 1,057 | .3349 | 354 |
| Riffle | 1,228 | 1,215 | .3119 | 383 |
| Puddled | 63 | 44 | .1111 | 7 |
| unclass. IP w/ss | 2,053 | 1,988 | .0974 | 200 |
| Isolated Pool | 43 | 43 | .0930 | 4 |
| Glide | 1,178 | 1,178 | .0925 | 109 |
| Isolated Pool w/ss | 2,604 | 2,604 | .0445 | 116 |
| Bull Trout | | | | |
| Plunge Pool | 165 | 165 | .0121 | 2 |
| Riffle With Pockets | 732 | 330 | .0027 | 2 |
| scour Pool | 1,057 | 66 | .0009 | 1 |
| Natural Juvenile Chinook Salmon | | | | |
| Dammed Pool | 26 | 26 | 1.3461 | 35 |
| Backwater Pool | 87 | 34 | 0.2759 | 24 |
| Plunge Pool | 165 | 165 | 0.1333 | 22 |
| Glide | 1,178 | 890 | 0.0993 | 117 |
| Lateral Pool | 364 | 265 | 0.0522 | 19 |
| Unclass. IP w/ss | 2,053 | 1,757 | 0.0502 | 103 |
| Scour Pool | 1,057 | 1,057 | 0.0435 | 46 |
| Riffle | 1,228 | 1,140 | 0.0269 | 33 |
| Isolated Pool w/ss | 2,604 | 1,242 | 0.0092 | 24 |
| Riffle With Pockets | 732 | 402 | 0.0068 | 5 |
| Rapid Over Boulders | 635 | 169 | 0.0063 | 4 |
| Adult Chinook Salmon | | | | |
| Plunge Pool | 165 | 165 | 0.0060 | 1 |
| Lateral Pool, | 364 | 53 | 0.0027 | 1 |
| Rapid Over Boulders | 635 | 169 | 0.0016 | 1 |
| Mountain Whitefish | | | | |
| Plunge Pool | 165 | 165 | 0.1273 | 21 |
| Rapid Over Boulders | 635 | 528 | 0.0760 | 48 |
| Scour Pool | 1,057 | 622 | 0.0757 | 80 |
| Riffle With Pockets | 732 | 557 | 0.0533 | 39 |
| Lateral Pool | 364 | 150 | 0.0247 | 9 |
| Riffle | 1,228 | 534 | 0.0060 | 7 |

Table E-7. Mean Density and Population Estimate per Habitat Type of **Rainbow/Steelhead Trout, Coho, and Chinook Salmon**, Moonshine Creek, RM **0.0-4.4, 9/18-9/21, 1995.**

| HABITAT TYPE | TOTAL AREA SAMPLED/M ² | AREA/M ² W/ SPP. PRESENT | MEAN DENSITY AND TOTAL AREA | EST. # IN UNIT |
|--|-----------------------------------|-------------------------------------|-----------------------------|----------------|
| Natural Rainbow/Steelhead Trout | | | | |
| Plunge Pool | 64 | 64 | 2.2186 | 142 |
| Trench Pool | 7 | 7 | 1.8571 | 13 |
| Riffle With Pockets | 100 | 87 | 0.4900 | 49 |
| Lateral Pool | 135 | 90 | 0.4667 | 63 |
| Scour Pool | 171 | 165 | 0.4269 | 73 |
| Cascade Over Bedrock | 23 | 15 | 0.3913 | 9 |
| Backwater Pool | 9 | 8 | 0.2222 | 2 |
| Beaver Dam Pool | 31 | 31 | 0.1290 | 4 |
| Glide | 157 | 111 | 0.0764 | 12 |
| Puddled | 167 | 26 | 0.0599 | 10 |
| Rapid Over Bedrock | 36 | 17 | 0.0556 | 2 |
| Rapid Over Boulder | 46 | 15 | 0.0435 | 2 |
| Riffle | 156 | 55 | 0.0385 | 6 |
| Natural Juvenile Coho Salmon | | | | |
| Scour Pool | 171 | 73 | 0.0526 | 9 |
| Natural Juvenile Chinook Salmon | | | | |
| Scour Pool | 171 | 73 | 0.0058 | 1 |

Table E-8. Mean Density and Population Estimate per Habitat Type of **Rainbow/Steelhead Trout, and Coho Salmon**, Mission Creek, RM **0.0-4.3, 9/5-9/13, 19%**.

| HABITAT TYPE | TOTAL AREA SAMPLED/M ² | AREA/M ² W/ SPP. PRESENT | MEAN DENSITY IN TOTAL AREA | EST. # IN UNIT |
|---|-----------------------------------|-------------------------------------|----------------------------|----------------|
| Natural Rainbow/Steelhead Trout | | | | |
| Plunge Pool | 47 | 39 | 1.2766 | 60 |
| Trench Pool | 27 | 22 | 1.0370 | 28 |
| Lateral Pool | 73 | 60 | 0.7945 | 58 |
| Scour Pool | 42 | 30 | 0.6905 | 29 |
| Glide | 41 | 32 | 0.3659 | 15 |
| Dammed Pool | 12 | 7 | 0.3333 | 4 |
| Backwater Pool | 4 | 2 | 0.2500 | 1 |
| Riffle With Pockets | 48 | 12 | 0.2083 | 10 |
| Riffle | 114 | 66 | 0.0351 | 4 |
| Hatchery Rainbow/Steelhead Trout | | | | |
| Plunge Pool | 47 | 7 | 0.0213 | 1 |
| Natural Juvenile Coho Salmon | | | | |
| Plunge Pool | 47 | 7 | 0.3191 | 15 |
| Scour Pool | 42 | 10 | 0.0952 | 4 |
| Lateral Pool | 73 | 5 | 0.0274 | 2 |

Table E-9. Mean Density and Population Estimate per Habitat Type of **Rainbow/Steelhead Trout**, Coho and Chinook Salmon, Cottonwood Creek, RM 0.0-4.1, 7/5-8/1, 1995.

| HABITAT TYPE | TOTAL AREA SAMPLED/M ² | AREA/M ² W/SPP. PRESENT | MEAN DENSITY IN TOTAL AREA | EST. # IN UNIT |
|--|-----------------------------------|------------------------------------|----------------------------|----------------|
| Natural Rainbow/Steelhead Trout | | | | |
| Plunge Pool | 48 | 30 | 1.9167 | 92 |
| Trench Pool | 4 | 4 | 1.0000 | 4 |
| Scour Pool | 69 | 29 | 0.2319 | 16 |
| Lateral Pool | 118 | 63 | 0.1441 | 17 |
| Riffle With Pockets | 23 | 10 | 0.1304 | 3 |
| Rapid Over Boulders | 26 | 15 | 0.1154 | 3 |
| Dammed Pool | 48 | 45 | 0.1042 | 5 |
| Glide | 141 | 44 | 0.0355 | 5 |
| Riffle | 417 | 87 | 0.0288 | 12 |
| Isolated Pool | 1,143 | 7 | 0.0201 | 23 |
| Natural Juvenile Coho Salmon | | | | |
| Plunge Pool | 48 | 43 | 0.5625 | 27 |
| Lateral Pool | 118 | 40 | 0.0424 | 5 |
| Isolated Pool | 1,143 | 1,076 | 0.0149 | 17 |
| Riffle | 417 | 23 | 0.0024 | 1 |
| Natural Juvenile Chinook Salmon | | | | |
| Isolated Pool | 1,143 | 1,076 | 0.0009 | 1 |

Table E-10. Mean Density and Population Estimate per Habitat Type of **Rainbow/Steelhead Trout**, Coho and Chinook Salmon, Coonskin Creek, RM 0.0-2.0, 6/29-7/18, 1995.

| HABITAT TYPE | TOTAL AREA SAMPLED/M ² | AREA/M ² W/SPP. PRESENT | MEAN DENSITY IN TOTAL AREA | EST # IN UNIT |
|--|-----------------------------------|------------------------------------|----------------------------|---------------|
| Natural Rainbow/Steelhead Trout | | | | |
| Trench Pool | 20 | 20 | 4.0000 | 80 |
| Plunge Pool | 56 | 37 | 0.7090 | 95 |
| Scour Pool | 94 | 61 | 0.6596 | 62 |
| Lateral Pool | 144 | 83 | 0.2430 | 35 |
| Riffle With Pockets | 135 | 53 | 0.1556 | 21 |
| Rapid Over Boulders | 104 | 42 | 0.1154 | 12 |
| Glide | 171 | 147 | 0.0877 | 15 |
| Riffle | 130 | 33 | 0.0462 | 6 |
| Natural Juvenile Coho Salmon | | | | |
| Trench Pool | 20 | 3 | 0.7000 | 14 |
| Scour Pool | 94 | 17 | 0.3617 | 34 |
| Lateral Pool | 144 | 56 | 0.2431 | 35 |
| Rapid Over Boulders | 104 | 20 | 0.0673 | 7 |
| Rapid Over Bedrock | 22 | 12 | 0.0454 | 1 |
| Riffle | 130 | 13 | 0.0385 | 5 |
| Natural Juvenile Chinook Salmon | | | | |
| Rapid Over Bedrock | 22 | 12 | 0.1364 | 3 |
| Scour Pool | 94 | 17 | 0.0851 | 8 |
| Trench Pool | 20 | 2 | 0.0500 | 1 |

Table E-1 1. Habitat of Mountain Whitefish, Umatilla River. RM 81.8-89.6. 8/8-8/25, 1995.

| HABITAT TYPE | AREA SAMPLED /m ² | # CAPTURED | % OF TOTAL CATCH | DENSITY* | EXPANDED POPULATION ESTIMATE* | RM RANGE | MEAN RM | |
|--------------------------------|------------------------------|--------------|------------------|---------------|-------------------------------|---------------|------------------|-------------|
| FAST WATER T TYPE | | | | | | | | |
| Rapid Over Boulders | 9,614 | 635 | 40 | 21.6 | 0.0630* | 606 | 88.3-88.7 | 88.3 |
| Riffle With Pockets | 22,653 | 732 | 35 | 18.9 | 0.0478' | 1,083 | 82.2-88.4 | 87.0 |
| Riffle | 60,403 | 1,228 | 7 | 3.8 | 0.0060* | 344 | 82.4-83.6 | 83.0 |
| Subtotal | 92,670 | 2,595 | 82 | 44.3 | 0.0220* | 2,033 | 82.2-88.7 | 87.3 |
| SLOW WATER HABITAT TYPE | | | | | | | | |
| Straight Scour Pool | 14,201 | 1,057 | 73 | 39.5 | 0.0691* | 981 | 82.3-88.5 | 87.8 |
| Plunge Pool | 250 | 165 | 21 | 11.4 | 0.1273* | 32 | 89.2 | 89.2 |
| Lateral Scour Pool | 23,629 | 364 | 9 | 4.9 | 0.0247' | 584 | 83.3-88.6 | 87.9 |
| Subtotal | 38,080 | 1,586 | 103 | 55.7 | 0.0649* | 1,597 | 82.3-88.6 | 88.1 |
| TOTAL | 130,750 | 4,181 | 185 | 100.00 | 0.0442* | 3,630* | 82.2-88.7 | 87.7 |

* Density was only estimated for units where mountain whitefish were captured.

- Mountain whitefish were not captured in other habitat types.

Table E-12. Actual, Estimated Number and Percentage with Minimum, Maximum and Mean Lengths, and RM Range of Salmonids captured in the Umatilla River, RM 81.8-89.6, 8/8-8/25, 1995.

| SPECIES | % SPECIES COMPOSITION | TOTAL # CAPTURED | EXPANDED ESTIMATED # OF EACH SPECIES | MIN,MEAN,MAX LENGTHS (mm) | RM RANGE |
|-----------------------------------|-----------------------|------------------|--------------------------------------|---------------------------|------------------|
| Rainbow/Steelhead Trout - Natural | 78.50 | 1,899 | 54,258 | 29,84,258 | 81.9-89.4 |
| Juvenile Chinook Salmon - Natural | 13.52 | 327 | 9,343 | 65,89,127 | 81.9-89.3 |
| Mountain Whitefish - Natural | 7.65 | 185 | 5,286 | 116,258,440 | 82.2-88.7 |
| Bull Trout - Natural | 0.21 | 5 | 152 | 170,223,265 | 87.7-89.2 |
| Adult Spring Chinook | 0.12 | 3 | 96 | 540,655,850 | 88.0-89.2 |
| TOTAL | 100.00% | 2,419 | 69,116 | 29,99,850 | 81.9-89.4 |

Table E-13. Actual, Estimated Number and Percentage with Minimum, Maximum and Mean Lengths, and RM Range of Salmonids captured in Moonshine Creek, RM 0-4.4, 9/18-9/21, 1995.

| SPECIES | % SPECIES COMPOSITION | TOTAL # CAPTURED | EXPANDED ESTIMATED # OF EACH SPECIES | MIN,MEAN,MAX LENGTHS (mm) | RM RANGE |
|-----------------------------------|-----------------------|------------------|--------------------------------------|---------------------------|----------------|
| Rainbow/Steelhead Trout - Natural | 97.36 | 369 | 1,138 | 48,107,240 | 0.0-4.2 |
| Coho Salmon - Natural | 2.38 | 9 | 28 | 88,91,95 | 0.2 |
| Chinook Salmon - Natural | 0.26 | 1 | 3 | 88 | 0.2 |
| TOTAL | 100.00% | 379 | 1,169 | 48,107,240 | 0.0-4.2 |

Table E-14. Actual, Estimated Number and Percentage with Minimum, Maximum and Mean Lengths, and RM Range of Salmonids captured in Mission Creek, RM 0-4.3, 9/5-9/13, 1995.

| SPECIES | % SPECIES COMPOSITION | TOTAL # CAPTURED | EXPANDED ESTIMATED # OF EACH SPECIES | MIN,MEAN,MAX LENGTHS (mm) | RM RANGE |
|------------------------------------|-----------------------|------------------|--------------------------------------|---------------------------|----------------|
| Rainbow/Steelhead Trout - Natural | 90.18 | 202 | 839 | 56,122,290 | 0.5-4.2 |
| Coho Salmon - Natural | 9.38 | 21 | 87 | 75,90,100 | 0.5 |
| Rainbow/Steelhead Trout - Hatchery | 0.44 | 1 | 4 | 230 | 0.5 |
| TOTAL | 100.00% | 224 | 930 | 56,120,290 | 0.5-4.2 |

Table E-15. Actual, Estimated Number and Percentage with Minimum, Maximum and Mean Lengths, and RM range of Salmonids captured in Cottonwood Creek, RM O-4.1, 7/5-8/1, 1995.

| SPECIES | % SPECIES COMPOSITION | TOTAL # CAPTURED | EXPANDED ESTIMATED # OF EACH SPECIES | MIN,MEAN,MAX LENGTHS (mm) | RM RANGE |
|-----------------------------------|-----------------------|------------------|--------------------------------------|---------------------------|----------------|
| Rainbow/Steelhead Trout - Natural | 78.18 | 172 | 489 | 37,111,340 | 0.0-3.1 |
| Coho Salmon - Natural | 21.36 | 47 | 134 | 69,84,103 | 0.1-1.1 |
| Chinook Salmon - Natural | 0.46 | 1 | 3 | 63 | 0.0-0.1 |
| TOTAL | 100.00% | 220 | 626 | 37,105,340 | 0.0-3.1 |

Table E-16. Actual, Estimated Number and Percentage with Minimum, Maximum and Mean Lengths, and RM Range of Salmonids captured in Coonskin Creek, RM O-2.0, 6/29-7/18, 1995.

| SPECIES | % SPECIES COMPOSITION | TOTAL # CAPTURED | EXPANDED ESTIMATED # OF EACH SPECIES | MIN,MEAN,MAX LENGTHS (mm) | RM RANGE |
|-----------------------------------|-----------------------|------------------|--------------------------------------|---------------------------|----------------|
| Rainbow/Steelhead Trout - Natural | 76.04 | 311 | 1,426 | 42,108,327 | 0.0-2.0 |
| Coho Salmon - Natural | 21.03 | 86 | 394 | 64,79,90 | 0.1-0.2 |
| Chinook Salmon - Natural | 2.93 | 12 | 55 | 74,83,90 | 0.1-0.2 |
| TOTAL | 100.00 | 409 | 1,875 | 42,101,327 | 0.0-2.0 |

Table E-17. Number of Non-Salmonids visually estimated or captured* from 74 of 648 units, Umatilla River, RM 81.8-89.6, 8/8-8/25, 1995.

| SPECIES | NUMBER VISUALLY ESTIMATED | % OF NUMBER VISUALLY ESTIMATED | DENSITY OF NON-SALMONIDS | EXPANDED NON-SALMONID ESTIMATE | NON-SALMONID TO SALMONID RATIO |
|--|---------------------------|--------------------------------|--------------------------|--------------------------------|--------------------------------|
| Speckled Dace (Rhinichthys osculus) | 5,411 | 53.71 | 0.5287 | 81,418 | 1.180:1 |
| Sculpin (Cottus spp.) | 4,550 | 45.16 | 0.4446 | 68,463 | 0.991:1 |
| Redside Shiner (Richardsonius balteatus) | 91 | 0.90 | 0.0089 | 1,369 | 0.020: 1 |
| Sucker (Catostomus spp.) | 17 | 0.17 | 0.0017 | 256 | 0.004:1 |
| Northern Squawfish^ (Ptychocheilus oregonesis) | 6 | 0.06 | 0.0006 | 6 | 0.001:1 |
| TOTAL | 10,075 | 100.00 | 0.9844 | 151,511 | 2.193:1 |

* Conservative estimate, see methods section for expansion methodology.

^ Northern Squawfish were the only non-salmonid captured.

Table E-18. Number of Non-Salmonids visually estimated from 90 of 594 units, Moonshine Creek, RM O-4.4, 9/18-9/21, 1995.

| SPECIES | NUMBER VISUALLY ESTIMATED | % OF NUMBER VISUALLY ESTIMATED | DENSITY OF NON-SALMONIDS | EXPANDED NON-SALMONID ESTIMATE | NON-SALMONID TO SALMONID RATIO |
|-------------------------------------|---------------------------|--------------------------------|--------------------------|--------------------------------|--------------------------------|
| Sucker (Catostomus spp.) | 455 | 44.70 | 0.4121 | 4,621 | 3.953:1 |
| Sculpin (Cottus spp.) | 368 | 36.15 | 0.3334 | 3,738 | 3.198:1 |
| Speckled Dace (Rhinichthys osculus) | 195 | 19.15 | 0.1767 | 1,981 | 1.695:1 |
| TOTAL | 1,018 | 100.00 | 0.9221 | 10,340 | 8.845:1 |

Table E-19. Number of Non-Salmonids visually estimated from 65 of 872 units, Mission Creek, RM o-4.3, 9/5-9/13, 1995.

| SPECIES | NUMBER VISUALLY ESTIMATED | % OF NUMBER VISUALLY ESTIMATED | DENSITY OF NON-SALMONIDS | EXPANDED NON-SALMONID ESTIMATE | NON-SALMONID TO SALMONID RATIO |
|---|---------------------------|--------------------------------|--------------------------|--------------------------------|--------------------------------|
| Speckled Dace (<i>Rhinichthys osculus</i>) | 350 | 76.92 | 0.7954 | 7,943 | 8.541:1 |
| Sculpin (<i>Cottus</i> spp.) | 85 | 18.68 | 0.1932 | 1,929 | 2.074:1 |
| Redside Shiner (<i>Richardsonius blateatus</i>) | 20 | 4.40 | 0.0455 | 454 | 0.488:1 |
| TOTAL | 455 | 100.00 | 1.0340 | 10,326 | 11.103:1 |

Table E-20. Number of Non-Salmonids visually estimated from 70 of 912 units, Cottonwood Creek, RM 04.1, 7/5-8/1, 1995.

| SPECIES | NUMBER VISUALLY ESTIMATED | % OF NUMBER VISUALLY ESTIMATED | DENSITY OF NON-SALMONIDS | EXPANDED NON-SALMONID ESTIMATE | NON-SALMONID TO SALMONID RATIO |
|---|---------------------------|--------------------------------|--------------------------|--------------------------------|--------------------------------|
| Speckled Dace (<i>Rhinichthys osculus</i>) | 1,401 | 85.06 | 0.4926 | 7,602 | 10.150:1 |
| Sculpin (<i>Cottus</i> spp.) | 106 | 6.44 | 0.0373 | 575 | 0.768:1 |
| Redside Shiner (<i>Richardsonius blateatus</i>) | 80 | 4.86 | 0.0281 | 434 | 0.579:1 |
| Sucker (<i>Catostomus</i>) | 60 | 3.64 | 0.0211 | 326 | 0.435:1 |
| TOTAL | 1,647 | 100.00 | 0.5792 | 8,937 | 11.932:1 |

Table E-21. Number of Non-Salmonids visually estimated from 87 of 626 units, Coonskin Creek, RM O-2.0, 6/29-7/18, 1995.

| SPECIES | NUMBER VISUALLY ESTIMATED | % OF NUMBER VISUALLY ESTIMATED | DENSITY OF NON-SALMONIDS | EXPANDED NON-SALMONID ESTIMATE | NON-SALMONID TO SALMONID RATIO |
|--|---------------------------|--------------------------------|--------------------------|--------------------------------|--------------------------------|
| Speckled Dace (<i>Rhinichthys osculus</i>) | 215 | 71.19 | 0.2375 | 1,392 | 0.742:1 |
| Sculpin (<i>Cottus</i> spp.) | 87 | 21.81 | 0.0961 | 563 | 0.300:1 |
| TOTAL | 302 | 100.00 | 0.3336 | 1,955 | 1.043:1 |

Table E-22. Index Site Summary; Site, Date Sampled, Site Composition, Discharge, Salmonid Catch Per Unit Effort (Fish Per Minute), and Mean Catch, 1995. (^ Juvenile Hatchery Coho).

| SITE # | STREAM | RM | DATE 1995 | SITE L(m)* | SLOW Water L(m)* | % | FAST Water L(m)* | % | DISCHG* | CPUE* (FPM) | MEAN CPUE |
|--------|----------------|------|-----------|------------|------------------|----|------------------|----|---------|-------------|-----------|
| 01 | Umatilla River | 1.5 | 4/10 | 213 | 110 | 52 | 103 | 48 | MF/HF | 0.8 | 0.4 |
| 01 | Umatilla River | 1.5 | 9/13 | 213 | 147 | 69 | 66 | 38 | LF | 0 | |
| 01 | Umatilla River | 1.5 | 1/28 | 213 | 133 | 62 | 80 | 38 | HF | 0.5 | |
| 02 | Umatilla River | 9.0 | 4/10 | 152 | 95 | 63 | 57 | 37 | MF/HF | 1.9^ | 0.6^ |
| 02 | Umatilla River | 9.0 | 9/18 | 152 | 100 | 66 | 52 | 34 | LF | 0 | |
| 02 | Umatilla River | 9.0 | 11/28 | 152 | 97 | 64 | 55 | 36 | MF/HF | 0 | |
| 03 | Umatilla River | 25.0 | 4/10 | 138 | 91 | 66 | 47 | 34 | MF/HF | 0.4 | 0.2 |
| 03 | Umatilla River | 25.0 | 9/13 | 138 | 85 | 62 | 63 | 38 | LF | 0 | |
| 03 | Umatilla River | 25.0 | 11/28 | 138 | 46 | 33 | 91 | 67 | MFIHF | 0.1 | |
| 04 | Umatilla River | 38.0 | 4/17 | 402 | 314 | 78 | 88 | 22 | MF/HF | 0 | 0 |
| 04 | Umatilla River | 38.0 | 9/20 | 402 | 324 | 81 | 78 | 19 | LF/MF | 0.1 | |
| 04 | Umatilla River | 38.0 | 11/21 | 402 | 337 | 84 | 65 | 16 | MF | 0 | |
| 05 | Umatilla River | 50.0 | 4/17 | 148 | 43 | 29 | 105 | 71 | MFIHF | 0 | 0.1 |
| 05 | Umatilla River | 50.0 | 9/14 | 148 | 95 | 64 | 53 | 36 | LF | 0.1 | |
| 05 | Umatilla River | 50.0 | 11/21 | 148 | 43 | 29 | 105 | 7 | MF | 0.1 | |
| 06 | Umatilla River | 60.0 | 4/6 | 127 | 29 | 23 | 98 | 77 | MF | 0.1 | 0.2 |
| 06 | Umatilla River | 60.0 | 9/14 | 127 | 28 | 22 | 99 | 78 | LF | 0.5 | |
| 06 | Umatilla River | 60.0 | 11/16 | 127 | 27 | 21 | 100 | 79 | MF | 0 | |
| 07 | Umatilla River | 67.5 | 4/5 | 234 | 70 | 30 | 164 | 70 | MF | 0 | 0.4 |
| 07 | Umatilla River | 67.5 | 9/19 | 234 | 106 | 45 | 128 | 55 | LF | 0.9 | |
| 07 | Umatilla River | 67.5 | 11/16 | 234 | 60 | 26 | 174 | 74 | MF | 0.4 | |
| 08 | Umatilla River | 74.0 | 4/6 | 168 | 78 | 46 | 130 | 54 | MF | 0.2 | 0.2 |
| 08 | Umatilla River | 74.0 | 9/20 | 168 | 63 | 38 | 105 | 62 | LF | 0.2 | |
| 08 | Umatilla River | 74.0 | 11/27 | 168 | 78 | 46 | 90 | 54 | MF/HF | 0.1 | |
| 09 | Umatilla River | 81.0 | 4/5 | 70 | 24 | 34 | 46 | 66 | MF | 0.8 | 0.7 |
| 09 | Umatilla River | 81.0 | 9/12 | 70 | 20 | 29 | 50 | 71 | LF | 1.0 | |
| 09 | Umatilla River | 81.0 | 11/27 | 70 | 25 | 36 | 45 | 64 | MF/HF | 0.3 | |
| 10 | Umatilla River | 88.0 | 4/5 | 92 | 53 | 58 | 39 | 42 | MF | 0.5 | 1.5 |
| 10 | Umatilla River | 88.0 | 9/12 | 92 | 54 | 59 | 38 | 41 | LF/MF | 1.6 | |
| 10 | Umatilla River | 88.0 | 11/30 | 92 | 56 | 61 | 36 | 39 | HF | 2.3 | |
| 11 | NF Umatilla R. | 1 | 3/24 | 37 | 13 | 35 | 24 | 65 | MFIHF | 0.5 | 1.8 |
| 11 | NF Umatilla R. | 1 | 9/27 | 37 | 16 | 43 | 26 | 57 | MF | 1.2 | |
| 11 | NF Umatilla R. | 1 | 11/20 | 37 | 13 | 35 | 24 | 65 | MF | 3.7 | |
| 12 | NF Umatilla R. | 3 | 3/24 | 41 | 9 | 22 | 32 | 78 | MF/HF | 0.4 | 1.0 |
| 12 | NF Umatilla R. | 3.0 | 9/27 | 41 | 16 | 39 | 25 | 61 | MF | 1.6 | |
| 12 | NF Umatilla R. | 3.0 | 11/20 | 41 | 13 | 32 | 28 | 68 | MF | 1.1 | |
| 13 | SF Umatilla R. | 1.0 | 3/27 | 76 | 33 | 43 | 43 | 57 | MF | 0.3 | 2.1 |
| 13 | SF Umatilla R. | 1.0 | 9/12 | 76 | 38 | 50 | 38 | 50 | LF | 3.9 | |
| 13 | SF Umatilla R. | 1.0 | | | | | | | | | |
| 14 | SF Umatilla R. | 4.0 | 3/27 | 47 | 3 | 28 | 34 | 72 | MF | 0.2 | 2.7 |
| 14 | SF Umatilla R. | 4.0 | 8/13 | 47 | 12 | 26 | 35 | 74 | LF | 3.8 | |
| 14 | SF Umatilla R. | 4.0 | 11/13 | 47 | 10 | 21 | 37 | 79 | HF | 4.0 | |

Table E-22. Continued.

| SITE # | STREAM | RM | DATE 1995 | SITE L(m) | SLOW Water L(m) | % | FAST Water L(m) | % | DISCHG | CPUE (FPM) | MEAN CPUE |
|--------|-----------------|------|-----------|-----------|-----------------|----|-----------------|-----|--------|------------|-----------|
| 15 | Birch Creek | 5.5 | 3/28 | 94 | 34 | 36 | 60 | 64 | MF/HF | 0.1 | 0.1 |
| 15 | Birch Creek | 5.5 | 9/18 | 94 | 58 | 62 | 36 | 38 | LF | 0 | |
| 15 | Birch Creek | 5.5 | 11/21 | 94 | 30 | 32 | 64 | 68 | MF | 0.1 | |
| 16 | Birch Creek | 10.0 | 3/28 | 77 | 16 | 21 | 61 | 79 | MF/HF | 0 | 0 |
| 16 | Birch Creek | 10.0 | 8/8 | 77 | 23 | 30 | 54 | 70 | LF | 0 | |
| 16 | Birch Creek | 10.0 | 11/14 | 77 | 23 | 30 | 54 | 70 | MF | 0 | |
| 17 | W. Birch Creek | 2.0 | 3/21 | 49 | 26 | 53 | 23 | 47 | HF | 0.2 | 0.2 |
| 17 | W. Birch Creek | 2.0 | 8/8 | 49 | 42 | 86 | 7 | 14 | LF | 0.2 | |
| 17 | W. Birch Creek | 2.0 | 11/14 | 49 | 38 | 76 | 11 | 24 | MF | 0.2 | |
| 18 | W. Birch Creek | 10.5 | 3/21 | 33 | 8 | 24 | 25 | 76 | MF/HF | 0.5 | 1.0 |
| 18 | W. Birch Creek | 10.5 | 8/8 | 33 | 0 | 0 | 33 | 100 | LF | 2.1 | |
| 18 | W. Birch Creek | 10.5 | 11/14 | 33 | 3 | 9 | 30 | 91 | MF | 0.3 | |
| 19 | E. Birch Creek | 4.5 | 3/21 | 45 | 15 | 33 | 30 | 67 | HF | 0.3 | 0.9 |
| 19 | E. Birch Creek | 4.5 | 8/8 | 45 | 0 | 0 | 45 | 100 | LF | 2.1 | |
| 19 | E. Birch Creek | 4.5 | 11/14 | 45 | 3 | 7 | 42 | 93 | MF | 0. | |
| 20 | E. Birch Creek | 13.0 | 3/21 | 18 | 9 | 50 | 9 | 50 | MF/HF | 1.9 | 3.4 |
| 20 | E. Birch Creek | 13.0 | 8/8 | 18 | 12 | 67 | 6 | 33 | LF | 3.5 | |
| 20 | E. Birch Creek | 13.0 | 11/14 | 18 | 13 | 72 | 5 | 28 | LF/MF | 4.9 | |
| 21 | Bear Creek | 1.0 | 4/12 | 29 | 29 | 10 | 0 | 0 | MF | 0.5 | 0.3 |
| 21 | Bear Creek | 1.0 | 9/22 | 29 | 23 | 09 | 6 | 21 | LF | 0.2 | |
| 21 | Bear Creek | 1.0 | 11/15 | 29 | 17 | 59 | 12 | 41 | LF/MF | 0.2 | |
| 22 | Bear Creek | 4.5 | 4/12 | 77 | 22 | 29 | 55 | 71 | MF | 1.5 | 2.8 |
| 22 | Bear Creek | 4.5 | 8/8 | 77 | 61 | 79 | 16 | 21 | MF | 1.9 | |
| 22 | Bear Creek | 4.5 | 11/15 | 77 | 34 | 44 | 43 | 56 | LF/MF | 5.0 | |
| 23 | Bridge Creek | 1.0 | 3/22 | 33 | 16 | 48 | 17 | 52 | MF/HF | 0.5 | 0.6 |
| 23 | Bridge Creek | 1.0 | 8/8 | 33 | 13 | 39 | 30 | 61 | LF | 0.5 | |
| 23 | Bridge Creek | 1.0 | 11/14 | 33 | 8 | 24 | 25 | 76 | LF/MF | 0.8 | |
| 24 | Pearson Creek | 2.0 | 3/21 | 21 | 12 | 57 | 9 | 43 | MF/HF | 0.9 | 2.9 |
| 24 | Pearson Creek | 2.0 | 8/8 | 21 | 4 | 19 | 19 | 81 | LF | 3.5 | |
| 24 | Pearson Creek | 2.0 | 11/14 | 21 | 9 | 43 | 12 | 57 | MF | 4.4 | |
| 25 | Buckaroo Creek | 1.0 | 3/17 | 17 | 10 | 59 | 8 | 41 | MF/HF | 0 | 0.9 |
| 25 | Buckaroo Creek | 1.0 | 8/4 | 17 | 8 | 41 | 10 | 59 | LF | 1.3 | |
| 25 | Buckaroo Creek | 1.0 | 11/8 | 17 | 8 | 47 | 9 | 53 | LF | 1.5 | |
| 26 | Squaw Creek | 2.5 | 3/23 | 57 | 17 | 30 | 40 | 70 | MF | 0.2 | 2.6 |
| 26 | Squaw Creek | 2.5 | 8/7 | 57 | 8 | 14 | 49 | 86 | LF | 3.5 | |
| 26 | Squaw Creek | 2.5 | 11/8 | 57 | 11 | 19 | 46 | 81 | LF | 4.2 | |
| 27 | Squaw Creek | 7.0 | 3/23 | 71 | 13 | 18 | 58 | 82 | MF | 0.1 | 3.1 |
| 27 | Squaw Creek | 7.0 | 8/7 | 71 | 13 | 18 | 58 | 82 | LF | 6.7 | |
| 27 | Squaw Creek | 7.0 | 11/30 | 71 | 9 | 13 | 62 | 87 | MF/HF | 2.3 | |
| 28 | Boston Can. Cr. | 0.6 | 3/20 | 27 | 7 | 26 | 20 | 74 | MF/HF | 2.7 | 3.2 |
| 28 | Boston Can. Cr. | 0.6 | 8/4 | 27 | 7 | 26 | 20 | 74 | LF | 3.6 | |
| 28 | Boston Can. Cr. | 0.6 | 11/13 | 27 | 7 | 26 | 20 | 74 | MF | 3.3 | |

Table E22. Continued.

| SITE # | STREAM | RM | DATE 1995 | SITE L(m) | SLOW Water L(m) | % | PAS T Water L(m) | % | DISCH G | CPUE (FPM) | MEAN CPUE |
|--------|-----------------|------|-----------|-----------|-----------------|----|------------------|-----|---------|------------|-----------|
| 29 | Line Creek | 0.5 | 3/17 | 14 | 5 | 36 | 9 | 64 | MF/HF | 3.3 | 2.8 |
| 29 | Line Creek | 0.5 | 8/4 | 14 | 4 | 29 | 10 | 71 | LF | 2.3 | |
| 29 | Line Creek | 0.5 | 11/13 | 14 | 4 | 29 | 10 | 71 | MF | 2.7 | |
| 30 | Meacham Creek | 9.0 | 4/6 | 76 | 0 | 0 | 76 | 100 | MF | 0.2 | 0.9 |
| 30 | Meacham Creek | 9.0 | 8/8 | 76 | 0 | 0 | 76 | 100 | LF | 3 | |
| 30 | Meacham Creek | 9.0 | 11/29 | 76 | 0 | 0 | 76 | 100 | HF | 0.4 | |
| 31 | Camp Creek | 0.6 | 3/17 | 46 | 11 | 24 | 35 | 76 | MF/HF | 1.1 | 2.4 |
| 31 | Camp Creek | 0.6 | 8/4 | 46 | 20 | 43 | 26 | 57 | LF | 3 | |
| 31 | Camp Creek | 0.6 | 11/13 | 46 | 15 | 33 | 31 | 67 | MF | 3.1 | |
| 32 | NF Meacham | 0.5 | 4/14 | 80 | 42 | 53 | 38 | 47 | MF/HF | 0.3 | 1.8 |
| 32 | NF Meacham | 0.5 | 8/9 | 80 | 54 | 68 | 26 | 32 | LF | 3.5 | |
| 32 | NF Meacham | 0.5 | 11/29 | 80 | 44 | 55 | 36 | 45 | HF | 1.7 | |
| 33 | NF Meacham | 1.2 | 4/13 | 64 | 31 | 48 | 33 | 52 | MF/HF | 0.1 | 2.0 |
| 33 | NF Meacham | 1.2 | 8/9 | 64 | 34 | 53 | 30 | 47 | LF | 3.8 | |
| 33 | NF Meacham | 1.2 | -- | -- | -- | -- | -- | -- | -- | -- | |
| 34 | Meacham Creek | 17.0 | 4/6 | 79 | 42 | 53 | 37 | 47 | MF | 0.4 | 2.1 |
| 34 | Meacham Creek | 17.0 | 8/4 | 79 | 45 | 57 | 34 | 43 | LF | 5.3 | |
| 34 | Meacham Creek | 17.0 | 11/29 | 79 | 22 | 28 | 57 | 72 | HF | 0.7 | |
| 35 | E. Meacham Cr. | 0.3 | 3/22 | 42 | 21 | 50 | 21 | 50 | MF/HF | 0.1 | 2.0 |
| 35 | E. Meacham Cr. | 0.3 | 8/9 | 42 | 23 | 55 | 19 | 45 | LF | 3.9 | |
| 35 | E. Meacham Cr. | 0.3 | -- | -- | -- | -- | -- | -- | -- | -- | |
| 36 | Meacham Creek | 28.5 | 3/29 | 38 | 16 | 42 | 22 | 58 | MF/HF | 0.1 | 1.4 |
| 36 | Meacham Creek | 28.5 | 8/9 | 38 | 16 | 42 | 22 | 58 | LF | 4.0 | |
| 36 | Meacham Creek | 28.5 | 11/29 | 38 | 16 | 42 | 22 | 58 | HF | 0 | |
| 37 | Ryan Creek | 1.0 | -- | -- | -- | -- | -- | -- | -- | -- | 5.1 |
| 37 | Ryan Creek | 1.0 | -- | -- | -- | -- | -- | -- | -- | -- | |
| 37 | Ryan Creek | 1.0 | 11/16 | 51 | 10 | 20 | 41 | 80 | MF | 5.1 | |
| 38 | Thomas Creek | 2.5 | 3/20 | 20 | 4 | 20 | 16 | 80 | MF | 0 | 0 |
| 38 | Thomas Creek | 2.5 | 8/2 | 20 | 4 | 20 | 16 | 80 | LF | 0 | |
| 38 | Thomas Creek | 2.5 | 11/8 | 20 | 4 | 20 | 16 | 80 | LF | 0 | |
| 39 | Spring Creek | 0.2 | 3/20 | 23 | 7 | 30 | 16 | 70 | MF/HF | 0.2 | 3.1 |
| 39 | Spring Creek | 0.2 | 8/3 | 23 | 10 | 43 | 13 | 57 | LF | 5.5 | |
| 39 | Spring Creek | 0.2 | 11/8 | 23 | 7 | 30 | 16 | 70 | LF | 3.5 | |
| 40 | Shiiehom Cr. | 0.5 | 5/5 | 42 | 7 | 17 | 35 | 83 | MF | 0 | 1.8 |
| 40 | Shimmiehorn Cr. | 0.5 | 8/3 | 42 | 5 | 12 | 37 | 88 | LF | 3.5 | |
| 40 | Shimmiehorn Cr. | 0.5 | -- | -- | -- | -- | -- | -- | -- | -- | |

L (m) = site length in meters; LF = low flow; MF = medium flow; HF = high flow; CPUE = catch per unit effort; FPM = salmonid/minute.

Table E-23. Fish Passage Barriers in the Umatilla River Basin, Surveyed 3/16-11/8, 1994.

| STREAM | RIVER MILE | BARRIER TYPE | COMPOSITION | STEP HEIGHT (m) | DEGREE | RECOMMENDED ACTION |
|--|-------------------|-------------------------------|-----------------------|------------------------|-----------------|---------------------------|
| Umatilla River | 1.5 | Channel Modification | concrete | 0.7 | Partial | Modify |
| Umatilla River | 2.4 | Irrigation Dam | Concrete | 1.0 | Partial | Modify |
| Umatilla River | 49.0 | Vacated Irrigation Dam | Concrete | 1.2 | Partial | Remove |
| Jungle/Windy Springs Creek | . 1 | Culvert | Steel | 0.15 | Partial | Modify |
| McKay Creek | 6.0 | Earthen Dam | Earth/Concrete | 4 0 | Complete | Leave |
| Wildhorse Creek | 0.1 | Vacated Irrigation Dam | Concrete | 0.7 | Partial | Remove |
| Wildhorse Creek | 18.8 | Bridge | concrete | 1.0 | Partial | Modify |
| Greasewood Creek | 0.4 | Irrigated Dam | Concrete | 0.6 | Partial | Modify |
| Mission Creek | 1.2 | Rip-rap | Concrete Blocks | 0.7 | Partial | Remove |
| Mission Creek | 1.4 | Bridge | Concrete | 0.5 | Partial | Modify |
| Mission Creek | 1.7 | Frame | Steel | 0.7 | Partial | Remove |
| Mission Creek | 3.3 | Culvert | Steel | 0.8 | Partial | Modify |
| Cottonwood Creek | 0.6 | Culvert | Steel | 0.8 | Partial | Modify |
| Cottonwood Creek | 0.9 | Water Pipe and Casing | concrete | 1.1 | Partial | Modify |
| Cottonwood Creek | 1.3 | Bridge | Concrete | 0.7 | Partial | Modify |
| Moonshine Creek | 1.0 | Bridge | Concrete | 1.2 | Partial | Modify |
| Coonskin Creek | .30 | Culvert | Steel | 0.5 | Partial | Modify |
| Camp Creek | .25 | Irrigation Dam | Concrete | 1.3 | Partial | Remove |
| Un-named Tributary at RM 1.5 of SF Umatilla River | 0.1 | Culvert | Steel | 0.5 | Complete | Modify |
| Whitman springs | 0.1 | Culvert | Steel | 0.5 | Complete | Modify |




Figure E-1. Length Frequency of Natural Juvenile Chinook Salmon and Natural Rainbow/Steelhead Trout captured during electrofishing in the Umatilla River, RM 81.8-89.6, 8/8-8/25, 1995. (95B-UMT1.CH3)




Figure E-2. Length Frequency of Mountain Whitefish captured during electrofishing in the Umatilla River, RM 81.8-89.6, 8/8-8/25, 1995. (95B-UMT2.CH3)




Figure E-3. Length Frequency of Natural **Rainbow/Steelhead** Trout captured during **electrofishing** in Moonshine Creek, RM O-4.4, 9/18-9/21, 1995. (95B-MNS1.CH3)




Figure E-4. Length Frequency of Natural Juvenile Coho Salmon and Natural **Rainbow/Steelhead** Trout captured during **electrofishing** in Mission Creek, RM O-4.3, 9/18-9/21, 1995. (95B-MSH1.CH3)




Figure E-5. Length Frequency of Natural Juvenile Coho Salmon and Natural Rainbow/Steelhead Trout captured during electrofishing in Cottonwood Creek, RM O-4.1, 7/5-8/1, 1995. (95B-CTT1.CH3)




Figure E-6. Length Frequency of Natural Juvenile Coho Salmon and Natural Rainbow/Steelhead Trout captured during electrofishing in Coonskin Creek, RM O-2.0, 6/29-7/18, 1995. (95B-CSK1.CH3)

APPENDIX F Adult Passage Examinations 1994-1995

Table F-1: Summer steelhead release dates, migrational timing, passage routes, and passage times (in days, hours and minutes) for Westland, Feed, and Stanfield Dams. Passage times between Three Mile Dam and Westland, Three Mile Dam and Stanfield, Westland and Feed, Feed and Stanfield, and Stanfield and ODFW (RM 56) is also included.

Westland (site 1)

| Ch/Code | R d . Date | R d . Time | First | | Last | | Westland Passage | | | Total Flows | | Avg: | | Westland to Feed | | Total Hours |
|---------|---------------|---------------|----------|-------|----------|-------|---------------------|-------|-------|-------------|------|------|---------|---------------------|-------|----------------|
| | | | Date | Time | Date | Time | Hrs/Min | Hours | (cfs) | T | amps | days | hrs/min | | | |
| 7/39 | 11/10/94 | 10:25 | 12/21/94 | 12:48 | 12/21/94 | 13:35 | 1 | 0 | 00:47 | 0.76 | 673 | 45.6 | 5 | 14:01 | 134 | |
| 7/40 | 11/17/94 | 10:05 | 12/21/94 | 15:42 | 12/21/94 | 18:35 | 1 | 0 | 00:53 | 0.88 | 673 | 45.0 | 5 | 06:27 | 126.5 | |
| 7/45 | 11/30/94 | 10:30 | 02/04/95 | 10:55 | 02/04/95 | 12:55 | 2 | 0 | 02:00 | 2 | 2650 | 45.4 | 0 | 20:36 | 20.6 | |
| 7/47 | 01/27/95 | 10:25 | 02/07/95 | 09:55 | 02/07/95 | 11:26 | 2 | 0 | 01:31 | 1.52 | 1760 | 44.8 | 0 | 02:56 | 2.933 | |
| 7/42 | 01/13/95 | 10:25 | 02/18/95 | 03:58 | 02/18/95 | 11:56 | 2 | 0 | 07:56 | 7.97 | 1160 | 46.3 | 0 | 20:20 | 20.33 | |
| 7/37 | 12/05/94 | 10:00 | 02/24/95 | 16:10 | 02/25/95 | 13:30 | 2 | 0 | 21:20 | 21.3 | 1640 | 46.6 | 1 | 01:08 | 25.13 | |
| 7/46 | 01/18/95 | 10:10 | 02/23/95 | 07:15 | 02/23/95 | 19:08 | 2 | 0 | 11:53 | 11.9 | 2210 | 46.7 | 0 | 06:28 | 6.467 | |
| 7/48 | 02/08/95 | 10:30 | 02/27/95 | 07:20 | 03/04/95 | 12:24 | 1 | 5 | 05:04 | 125 | 1263 | 44.4 | 4 | 23:04 | 1.181 | |
| 7/3 | 03/23/95 | 10:10 | 03/30/95 | 15:17 | 03/30/95 | 18:16 | 1 | 0 | 02:59 | 2.96 | 657 | 45.2 | 0 | 15:38 | 15.63 | |
| 7/85 | 03/14/95 | 10:20 | 03/27/95 | 15:04 | 03/27/95 | 18:31 | 2 | 0 | 03:27 | 3.45 | 1000 | 45.0 | 1 | 01:14 | 2523 | |
| 7/88 | 03/13/95 | 10:45 | 03/24/95 | 07:33 | 03/24/95 | 12:57 | 2 | 0 | 05:24 | 5.4 | 1550 | 43.2 | 0 | 01:49 | 1.617 | |
| 7/81 | 03/08/95 | 10:45 | 03/28/95 | 18:45 | 03/28/95 | 01:03 | 1 | 0 | 08:18 | 6.3 | 950 | 47.2 | 0 | 19:49 | 19.62 | |
| 7/5 | 03/27/95 | 10:30 | 04/06/95 | 08:54 | 04/06/95 | 08:00 | 2 | 0 | 01:08 | 1.1 | 666 | 40.7 | 0 | 03:30 | 3.5 | |
| 7/82 | 03/08/95 | 10:45 | 04/04/95 | 07:08 | 04/04/95 | 08:59 | 1 | 0 | 01:51 | 1.85 | 707 | 51.4 | 0 | 01:19 | 1.317 | |
| 7/22 | 04/07/95 | 10:25 | 04/13/95 | 14:35 | 04/13/95 | 15:23 | 2 | 0 | 00:48 | 0.6 | 1310 | 46.5 | 0 | 02:58 | 2.967 | |
| 7/13 | 03/30/95 | 11:00 | 04/12/95 | 17:28 | 04/13/95 | 09:56 | 2 | 0 | 18:28 | 16.5 | 1240 | 46.6 | 0 | 08:01 | 6.017 | |
| | | | | | | | Avg: | | 0.55 | | 13.1 | | 1.39 | | 33.33 | |

Feed Canal (site 2)

| Ch/Code | Rel. Date | Rel. Time | First | | Last | | Feed Passage | | | Total Flows | | Avfl. : | | Feed to Stanfield | | Total Hours |
|---------|-----------|-----------|----------|-------|----------|-------|-----------------|-------|-------|-------------|------|---------|------|----------------------|-------|----------------|
| | | | Date | Time | Date | Time | Hrs/Min | Hours | (cfs) | Temps | Days | hrs/min | | | | |
| 7/39 | 11/10/94 | 10:25 | 12/27/94 | 03:36 | 12/27/94 | 11:20 | 1 | 0 | 07:44 | 7.73 | 1162 | 46.8 | 16 | 11:05 | 395.1 | |
| 7/40 | 11/17/94 | 10:05 | 12/26/94 | 23:00 | 12/27/94 | 12:31 | 1 | 0 | 13:29 | 13.5 | 762 | 46.2 | 20 | 00:05 | 460.1 | |
| 7/45 | 11/30/94 | 10:30 | 02/05/95 | 09:31 | 02/05/95 | 18:14 | 2 | 0 | 08:43 | 6.72 | 2446 | 46.6 | 1 | 05:42 | 29.7 | |
| 7/47 | 01/27/95 | 10:25 | 02/07/95 | 14:22 | 02/28/95 | 08:08 | 1 | 16 | 18:46 | 451 | 1601 | 45 | 1 | 07:49 | 31.82 | |
| 7/42 | 01/13/95 | 10:25 | 02/18/95 | 08:16 | 02/18/95 | 14:53 | 2 | 0 | 08:37 | 6.62 | 1676 | 49 | 2 | 01:38 | 49.6 | |
| 7/37 | 12/05/94 | 10:00 | 02/26/95 | 14:38 | 03/10/95 | 13:04 | 1 | 11 | 22:28 | 266 | 774 | 46.2 | 0 | 22:21 | 22.35 | |
| 7/46 | 01/18/95 | 10:10 | 02/24/95 | 01:36 | 03/09/95 | 15:57 | 1 | 13 | 14:21 | 326 | 891 | 46.3 | 2 | 00:24 | 46.4 | |
| 7/48 | 02/08/95 | 10:30 | 03/09/95 | 11:28 | 03/09/95 | 12:04 | 1 | 0 | 00:38 | 0.6 | 552 | 49.6 | 1 | 01:54 | 25.9 | |
| 7/3 | 03/23/95 | 10:10 | 03/31/95 | 09:54 | 04/02/95 | 18:10 | 1 | 2 | 08:16 | 56.3 | 563 | 50.1 | 0 | 18:01 | 16.02 | |
| 7/85 | 03/14/95 | 10:20 | 03/28/95 | 19:45 | 04/01/95 | 13:03 | 1 | 3 | 17:18 | 69.3 | 621 | 46 | 0 | 06:48 | 6.6 | |
| 7/88 | 03/13/95 | 10:45 | 03/24/95 | 14:48 | 03/25/95 | 14:04 | 2 | 0 | 23:18 | 23.3 | 1406 | 43.4 | 0 | 20:53 | 20.66 | |
| 7/81 | 03/08/95 | 10:45 | 03/29/95 | 20:52 | 03/29/95 | 21:33 | 1 | 0 | 00:41 | 0.66 | 665 | 47.6 | 1 | 04:45 | 26.75 | |
| 7/5 | 03/27/95 | 10:30 | 04/06/95 | 11:30 | 04/07/95 | 18:20 | 1 | 1 | 08:50 | 30.6 | 860 | 50 | 3 | 17:01 | 69.02 | |
| 7/82 | 03/08/95 | 10:45 | 04/04/95 | 10:18 | 04/04/95 | 10:36 | 1 | 0 | 00:20 | 0.33 | 531 | 51.4 | 0 | 05:43 | 5.717 | |
| 7/22 | 04/07/95 | 10:25 | 04/13/95 | 18:21 | 04/14/95 | 08:11 | 2 | 0 | 11:50 | 11.6 | 1315 | 46.5 | 0 | 13:42 | 13.7 | |
| 7/1 | 03/30/95 | 11:00 | 04/13/95 | 17:57 | 04/14/95 | 15:32 | 1 | 0 | 21:35 | 21.6 | 1315 | 46.5 | 7 | 02:32 | 170.5 | |
| | | | | | | | Avg: | | 3.46 | | 83.4 | | 3.74 | | 89.77 | |

Stanfield (site 3)

| Ch/Code | Rel. Date | Rel. Time | First | | Last | | Stanfield Passage | | | Total Flows | | Avg: | | Stanfield to ODFW | | Total Hours |
|---------|-----------|-----------|----------|-------|----------|-------|----------------------|-------|-------|-------------|------|------|---------|----------------------|-------|----------------|
| | | | Date | Time | Date | Time | Hrs/Min | Hours | (cfs) | T | amps | Days | Hrs/Min | | | |
| 7/39 | 11/10/94 | 10:25 | 01/12/95 | 22:25 | 01/13/95 | 02:21 | 1 | 0 | 03:56 | 3.93 | 1075 | 42 | 14 | 17:18 | 353.3 | |
| 7/40 | 11/17/94 | 10:05 | 01/18/95 | 12:36 | 01/18/95 | 13:45 | 1 | 0 | 01:09 | 1.15 | 2260 | 42 | 33 | 20:39 | 612.6 | |
| 7/45 | 11/30/94 | 10:30 | 02/06/95 | 23:56 | 02/07/95 | 07:43 | 2 | 0 | 07:47 | 7.76 | 2145 | 43.5 | 17 | 04:48 | 412.6 | |
| 7/47 | 01/27/95 | 10:25 | 02/27/95 | 16:57 | 02/27/95 | 17:48 | 1 | 0 | 00:51 | 0.85 | 1490 | 45.5 | na | na | na | |
| 7/42 | 01/13/95 | 10:25 | 02/21/95 | 16:29 | 02/21/95 | 17:58 | 2 | 0 | 01:29 | 1.46 | 3420 | 47.3 | na | na | na | |
| 7/37 | 12/05/94 | 10:00 | 03/11/95 | 11:25 | 03/11/95 | 12:18 | 2 | 0 | 00:53 | 0.66 | 951 | 50.2 | na | na | na | |
| 7/46 | 01/18/95 | 10:10 | 03/11/95 | 16:21 | 03/11/95 | 16:57 | 2 | 0 | 00:36 | 0.6 | 651 | 50.2 | na | na | na | |
| 7/48 | 02/08/95 | 10:30 | 03/10/95 | 13:58 | 03/10/95 | 15:39 | 2 | 0 | 01:41 | 1.66 | 731 | 46.4 | na | na | na | |
| 7/3 | 03/23/95 | 10:10 | 04/03/95 | 12:11 | 04/03/95 | 12:34 | 1 | 0 | 00:23 | 0.36 | 662 | 54.7 | na | na | na | |
| 7/85 | 03/14/95 | 10:20 | 04/01/95 | 19:51 | 04/01/95 | 20:30 | 2 | 0 | 00:39 | 0.65 | 727 | 53.3 | 5 | 01:04 | 121.1 | |
| 7/88 | 03/13/95 | 10:45 | 03/26/95 | 10:57 | 03/26/95 | 12:11 | 2 | 0 | 01:14 | 1.23 | 1350 | 47.7 | 3 | 00:48 | 72.6 | |
| 7/81 | 03/08/95 | 10:45 | 03/31/95 | 02:18 | 03/31/95 | 03:27 | 2 | 0 | 01:09 | 1.15 | 724 | 52.4 | 3 | 08:39 | 76.65 | |
| 7/5 | 03/27/95 | 10:30 | 04/11/95 | 11:21 | 04/11/95 | 15:07 | 1 | 0 | 03:46 | 3.77 | 1460 | 51.7 | na | na | na | |
| 7/82 | 03/08/95 | 10:45 | 04/04/95 | 16:21 | 04/04/95 | 0.701 | 2 | 0 | 00:29 | 0.46 | 734 | 54.3 | 5 | 04:08 | 124.1 | |
| 7/22 | 04/07/95 | 10:25 | 04/14/95 | 19:53 | 04/15/95 | 0.716 | 2 | 0 | 21:18 | 21.3 | 1360 | 40.1 | na | na | na | |
| 7/13 | 03/30/95 | 11:00 | 04/21/95 | 18:04 | 04/21/95 | 0.77 | 2 | 0 | 00:25 | 0.42 | 904 | 64.7 | 5 | 02:13 | 122.2 | |
| | | | | | | | Avg: | | 0.12 | | 2.96 | | 10.9 | | 262.2 | |

ODFW (site 4)

| Ch/Code | Rel. Date | Rel. Time | First | | Last | | 3MD to Westland | | | 3MD to above Stfld | | Total Hours | | |
|--|-----------|-----------|----------|-------|----------|-------|--------------------|---------|-------|-----------------------|---------|----------------|--------|--|
| | | | Date | Time | Date | Time | Days | Hrs/Min | Hours | Days | Hrs/Min | | | |
| 7/39 | 11/10/94 | 10:25 | 01/27/95 | 18:39 | 01/27/95 | 18:56 | 41 | 02:23 | 966 | 63 | 15:56 | 1526 | | |
| 7/40 | 11/17/94 | 10:05 | 02/18/95 | 10:24 | 02/18/95 | 10:25 | 34 | 05:37 | 622 | 60 | 03:40 | 1444 | | |
| 7/45 | 11/30/94 | 10:30 | 02/24/95 | 12:31 | 02/24/95 | 13:45 | 66 | 00:25 | 1564 | 66 | 21:13 | 1653 | | |
| 7/47 | 01/27/95 | 10:25 | na | na | na | na | 10 | 23:30 | 263 | 31 | 07:23 | 751.4 | | |
| 7/42 | 01/13/95 | 10:25 | na | na | na | na | 35 | 17:33 | 656 | 39 | 07:33 | 043.5 | | |
| 7/37 | 12/05/94 | 10:00 | na | na | na | na | 81 | 08:10 | 1950 | 86 | 02:18 | 2306 | | |
| 7/46 | 01/18/95 | 10:10 | na | na | na | na | 35 | 21:05 | 661 | 52 | 06:47 | 1255 | | |
| 7/48 | 02/08/95 | 10:30 | na | na | na | na | 18 | 20:50 | 453 | 30 | 05:09 | 725.2 | | |
| 7/3 | 03/23/95 | 10:10 | na | na | na | na | 7 | 05:07 | 173 | 11 | 02:24 | 266.4 | | |
| 7/85 | 03/14/95 | 10:20 | 04/06/95 | 21:34 | 04/06/95 | 22:08 | 13 | 04:44 | 317 | 18 | 10:10 | 442.2 | | |
| 7/88 | 03/13/95 | 10:45 | 03/29/95 | 12:59 | 03/29/95 | 13:17 | 10 | 20:48 | 281 | 13 | 01:26 | 313.4 | | |
| 7/81 | 03/08/95 | 10:45 | 04/03/95 | 10:06 | 04/03/95 | 10:50 | 22 | 08:00 | 536 | 24 | 16:42 | 592.7 | | |
| 7/5 | 03/27/95 | 10:30 | na | na | na | na | 9 | 20:24 | 236 | 15 | 04:37 | 364.6 | | |
| 7/82 | 03/08/95 | 10:45 | 04/09/95 | 20:56 | 04/09/95 | 21:30 | 26 | 20:23 | 692 | 29 | 06:05 | 702.1 | | |
| 7/22 | 04/07/95 | 10:25 | na | na | na | na | 6 | 04:10 | 149 | 6 | 08:48 | 195.6 | | |
| 7/13 | 03/30/95 | 11:00 | 04/26/95 | 20:42 | 04/26/95 | 21:14 | 13 | 06:28 | 318 | 22 | 07:29 | 535.5 | | |
| File name: 9495data; * -trap and haul evaluation | | | | | | | 272 | | 654 | | 36.5 | | 1178.4 | |

Table F-2: Summer Steelhead release dates at Three Mile Falls Dam and days required to successfully migrate from Three Mile Falls Dam to S1(Westland), S2 (Feed Canal), S3 (Stanfield), and S4 (ODFW Rm 56), Umatilla River, 1994-95.

| CH/CODE | 3MD RELEASE DATE | 3MD TO SITE #1 DAYS | 3MD TO SITE #2 DAYS | 3MD TO SITE #3 DAYS | 3MD TO SITE #4 DAYS |
|---------|------------------|---------------------|---------------------|---------------------|---------------------|
| 7/39 | 11/10/94 | 41.1 | 46.7 | 63.5 | 78.4 |
| 7/40 | 11/17/94 | 34.2 | 39.5 | 60.1 | 94.0 |
| 7/45 | 11/30/94 | 66.0 | 67.0 | 68.6 | 86.1 |
| 7/47 | 01/27/95 | 10.9 | 11.2 | 31.3 | n/a |
| 7/42 | 01/13/95 | 35.7 | 36.9 | 39.3 | n/a |
| 7/37 | 12/05/94 | 81.2 | 83.2 | 96.1 | n/a |
| 7/46 | 01/18/95 | 35.8 | 36.6 | 52.3 | n/a |
| 7/48 | 02/08/95 | 18.8 | 29.0 | 30.1 | n/a |
| 7/3 | 03/23/95 | 7.2 | 8.0 | 11.1 | n/a |
| 7/85 | 03/14/95 | 13.1 | 14.4 | 18.4 | 23.5 |
| 7/88 | 03/13/95 | 10.8 | 11.2 | 13.0 | 16.1 |
| 7/81 | 03/06/95 | 22.3 | 23.4 | 24.6 | 28.0 |
| 7/5 | 03/27/95 | 9.8 | 10.0 | 15.0 | n/a |
| 7/82 | 03/06/95 | 28.8 | 29.0 | 29.2 | 34.4 |
| 7/22 | 04/07/95 | 6.1 | 6.3 | 7.4 | n/a |
| 7/13 | 03/30/95 | 13.2 | 14.3 | 22.3 | 27.4 |
| | AVERAGE: | 27.2 | 29.2 | 36.4 | 48.5 |

Filename: 9495days

Table F-3: Spring Chinook Salmon release dates, migrational timing, passage routes, and passage times (in days, hours and minutes) for Westland, Feed, and Stanfield Dams. Passage times between Three Mile Falls Dam and Westland, Three Mile Falls Dam and Stanfield, Westland and Feed, Feed and Stanfield, and Stanfield and ODFW (RM 56) is also included.

Westland (site 1)

| Ch/Code | Rel. Date | Rel. Time | First | | Last | | Route | Days | Westland Passage | | Total Hours | Avg. Flows (cfs) | Avg. Temps | Westland to Feed | | Total Hours |
|---------|-----------|-----------|----------|-------|----------|-------|-------|------|------------------|-------|-------------|------------------|------------|------------------|---------|-------------|
| | | | Date | Time | Date | Time | | | Hrs/Min | Hours | | | | days | hrs/min | |
| 13/32 | 04/10/95 | 10:00 | 04/19/95 | 18:18 | 04/18/95 | 18:40 | 1 | 0 | 01:22 | 1.37 | 911 | 46.64 | 0 | 18:20 | 18.33 | |
| 13/34 | 04/11/95 | 10:20 | 04/19/95 | 20:57 | 04/18/95 | 22:14 | 2 | 0 | 01:17 | 1.26 | 911 | 46.64 | 0 | 14:42 | 14.7 | |
| 13/36 | 04/13/95 | 10:30 | 04/23/95 | 09:57 | 04/23/95 | 11:33 | 1 | 0 | 01:36 | 1.6 | 797 | 54.27 | 0 | 21:49 | 21.82 | |
| 13/37 | 04/14/95 | 09:55 | 04/22/95 | 19:12 | 04/23/95 | 20:45 | 2 | 1 | 01:33 | 25.5 | 796 | 53.07 | na | na | na | |
| 13/38 | 04/18/95 | 10:13 | 04/23/95 | 03:18 | 04/23/95 | 12:23 | 2 | 0 | 09:05 | 9.06 | 787 | 54.27 | 0 | 08:59 | 6.983 | |
| 13/40 | 04/20/95 | 10:20 | 04/23/95 | 04:30 | 04/23/95 | 06:21 | 2 | 0 | 01:51 | 1.65 | 797 | 54.27 | 0 | 04:34 | 4.567 | |
| 13/41 | 04/18/95 | 10:15 | 04/23/95 | 06:56 | 04/23/95 | 08:30 | 1 | 0 | 01:34 | 1.57 | 797 | 54.27 | 0 | 03:51 | 27.85 | |
| 13/31 | 04/24/95 | 10:40 | 04/28/95 | 08:05 | 04/28/95 | 09:22 | 2 | 0 | 01:17 | 1.26 | 805 | 55.22 | 0 | 03:55 | 3.917 | |
| 13/35 | 04/13/95 | 10:30 | 04/28/95 | 13:45 | 04/28/95 | 14:35 | 2 | 0 | 00:50 | 0.63 | 605 | 55.22 | 0 | 13:25 | 13.42 | |
| 13/43 | 04/24/95 | 10:40 | 04/28/95 | 18:39 | 04/28/95 | 19:12 | 1 | 0 | 00:33 | 0.55 | 605 | 55.22 | 0 | 09:58 | 9.967 | |
| 13/42 | 04/28/95 | 10:10 | na | na | na | na | na | na | na | na | na | na | na | na | na | |
| | | | | | | | | Avg: | 0.16 | | 4.5 | | 0.56 | | 13.51 | |

Feed Canal (site 2)

| Ch/Code | Rel. Date | Rel. Time | First | | Last | | Route | Days | Feed Passage | | Total Hours | Avg. Flows (cfs) | Avg. Temps | Feed to Stanfield | | Total Hours |
|---------|-----------|-----------|----------|-------|----------|-------|-------|------|--------------|-------|-------------|------------------|------------|-------------------|---------|-------------|
| | | | Date | Time | Date | Time | | | Hrs/Min | Hours | | | | Days | hrs/min | |
| 13/32 | 04/10/95 | 10:00 | 04/20/95 | 14:00 | 04/24/95 | 04:30 | 2 | 3 | 14:30 | 66.5 | 738 | 51.04 | 0 | 11:49 | 11.52 | |
| 13/34 | 04/11/95 | 10:20 | 04/20/95 | 12:56 | 04/25/95 | 05:14 | 1 | 4 | 16:18 | 112 | 721 | 52.71 | 0 | 08:17 | 6.253 | |
| 13/36 | 04/13/95 | 10:30 | 04/24/95 | 09:22 | 04/24/95 | 22:29 | 1 | 0 | 13:07 | 13.1 | 889 | 52.32 | 0 | 13:58 | 13.97 | |
| 13/37 | 04/14/95 | 09:55 | na | na | na | na | na | na | na | na | na | na | na | na | na | |
| 13/38 | 04/18/95 | 10:13 | 04/23/95 | 19:22 | 04/24/95 | 15:16 | 1 | 0 | 19:54 | 19.9 | 705 | 53.3 | 0 | 09:19 | 9.317 | |
| 13/40 | 04/20/95 | 10:20 | 04/23/95 | 10:55 | 04/23/95 | 13:14 | 1 | 0 | 02:19 | 2.32 | 720 | 54.27 | 0 | 07:22 | 7.367 | |
| 13/41 | 04/19/95 | 10:15 | 04/24/95 | 12:21 | 04/26/95 | 13:41 | 1 | 2 | 01:20 | 48.3 | 700 | 64.7 | na | na | na | |
| 13/31 | 04/24/95 | 10:40 | 04/28/95 | 13:17 | 04/28/95 | 17:08 | 2 | 0 | 03:51 | 3.65 | 737 | 55.22 | 2 | 03:41 | 51.66 | |
| 13/35 | 04/13/95 | 10:30 | 04/27/95 | 04:00 | 04/27/95 | 04:48 | 1 | 0 | 00:48 | 0.6 | 798 | 55.74 | 4 | 13:03 | 109 | |
| 13/43 | 04/24/95 | 10:40 | 04/27/95 | 05:10 | 05/22/95 | 02:38 | 2 | 24 | 21:28 | 587 | 2772 | 52.57 | 0 | 08:15 | 8.25 | |
| 13/42 | 04/28/95 | 10:10 | 05/18/95 | 14:02 | 05/19/95 | 01:05 | 2 | 0 | 11:03 | 11.1 | 1060 | 55.53 | 0 | 13:00 | 13 | |
| | | | | | | | | Avg: | 3.74 | | 60.7 | | 1.06 | | 25.66 | |

Stanfield (site 3)

| Ch/Code | Rel. Date | Rel. Time | First | | Last | | Route | Days | Stanfield Passage | | Total Hours | Avg. Flows (cfs) | Avg. Temps | Stanfield to ODFW | | Total Hours |
|---------|-----------|-----------|----------|-------|----------|-------|-------|------|-------------------|-------|-------------|------------------|------------|-------------------|---------|-------------|
| | | | Date | Time | Date | Time | | | Hrs/Min | Hours | | | | Days | Hrs/Min | |
| 13/32 | 04/10/95 | 10:00 | 04/24/95 | 18:19 | 04/24/95 | 16:40 | 2 | 0 | 00:21 | 0.35 | 689 | 52.32 | 13 | 11:31 | 323.5 | |
| 13/34 | 04/11/95 | 10:20 | 04/25/95 | 13:31 | 04/25/95 | 14:00 | 1 | 0 | 00:29 | 0.46 | 675 | 56.57 | 8 | 04:21 | 186.4 | |
| 13/36 | 04/13/95 | 10:30 | 04/25/95 | 12:27 | 04/25/95 | 13:04 | 2 | 0 | 00:37 | 0.62 | 675 | 56.57 | 20 | 13:40 | 403.7 | |
| 13/37 | 04/14/95 | 09:55 | na | na | na | na | na | na | na | na | na | na | na | na | na | |
| 13/38 | 04/18/95 | 10:13 | 04/25/95 | 00:35 | 04/25/95 | 01:39 | 2 | 0 | 01:04 | 1.07 | 675 | 56.57 | 13 | 14:35 | 326.6 | |
| 13/40 | 04/20/95 | 10:20 | 04/23/95 | 20:36 | 04/24/95 | 08:57 | 2 | 0 | 12:21 | 12.3 | 705 | 53.3 | 2 | 19:10 | 67.17 | |
| 13/41 | 04/19/95 | 10:15 | na | na | na | na | na | na | na | na | na | na | na | na | na | |
| 13/31 | 04/24/95 | 10:40 | 04/28/95 | 20:49 | 04/28/95 | 23:39 | 2 | 0 | 02:50 | 2.83 | 1450 | 52.76 | 19 | 18:55 | 474.9 | |
| 13/35 | 04/13/95 | 10:30 | 05/01/95 | 17:51 | 05/02/95 | 11:35 | 2 | 0 | 17:44 | 17.7 | 3781 | 47.85 | 16 | 14:48 | 398.8 | |
| 13/43 | 04/24/95 | 10:40 | 05/22/95 | 10:53 | 05/22/95 | 11:14 | 2 | 0 | 00:21 | 0.35 | 657 | 60.5 | 2 | 02:25 | 50.42 | |
| 13/42 | 04/28/95 | 10:10 | 05/19/95 | 14:05 | 05/19/95 | 14:36 | 2 | 0 | 00:31 | 0.52 | 1006 | 57 | 4 | 12:15 | 106.2 | |
| | | | | | | | | Avg: | 0.17 | | 4.03 | | 10.2 | | 244 | |

ODFW (site 4)

| Ch/Code | Rel. Date | Rel. Time | First | | Last | | Days | 3MD to Westland | | Total Hours | Days | 3MD to above Sffld | | Total Hours |
|---------|-----------|-----------|----------|-------|----------|-------|------|-----------------|-------|-------------|-------|--------------------|-------|-------------|
| | | | Date | Time | Date | Time | | Hrs/Min | Hours | | | Hrs/Min | Hours | |
| 13/32 | 04/10/95 | 10:00 | 05/08/95 | 04:11 | 05/08/95 | 04:19 | 9 | 08:18 | 224 | 14 | 06:40 | 342.7 | | |
| 13/34 | 04/11/95 | 10:20 | 05/03/95 | 18:21 | 05/03/95 | 19:04 | 8 | 10:37 | 203 | 14 | 03:40 | 339.7 | | |
| 13/36 | 04/13/95 | 10:30 | 05/18/95 | 02:44 | 05/18/95 | 03:08 | 9 | 23:27 | 238 | 12 | 02:34 | 290.6 | | |
| 13/37 | 04/14/95 | 09:55 | na | na | na | na | 8 | 09:17 | 201 | na | na | na | | |
| 13/38 | 04/18/95 | 10:13 | 05/08/95 | 16:14 | 05/08/95 | 16:56 | 4 | 17:05 | 113 | 5 | 22:44 | 142.7 | | |
| 13/40 | 04/20/95 | 10:20 | 04/27/95 | 04:07 | 04/27/95 | 11:50 | 2 | 18:10 | 66.2 | 6 | 13:19 | 205.3 | | |
| 13/41 | 04/19/95 | 10:15 | | | | | 3 | 20:41 | 92.7 | 13 | 01:20 | 313.3 | | |
| 13/31 | 04/24/95 | 10:40 | 05/18/95 | 18:34 | 05/18/95 | 18:50 | 1 | 21:25 | 45.4 | 26 | 00:34 | 672.6 | | |
| 13/35 | 04/13/95 | 10:30 | 05/19/95 | 02:23 | 05/19/95 | 02:45 | 13 | 03:15 | 315 | 36 | 04:06 | 868.1 | | |
| 13/43 | 04/24/95 | 10:40 | 05/24/95 | 13:39 | 05/24/95 | 13:50 | 2 | 07:59 | 56 | 26 | 00:34 | 672.6 | | |
| 13/42 | 04/28/95 | 10:10 | 05/24/95 | 02:51 | 05/24/95 | 03:16 | na | na | na | 23 | 13:50 | 556.4 | | |
| | | | | | | | | Avg: | 6.46 | 158 | 16.3 | 440.4 | | |

File name: data9495, - trap and haul evaluation

Table F-4: Summer steelhead passage times (days, hours, minutes) and miles moved per day between Stanfield Dam and ODFW (RM 56), Passage Evaluation, Umatilla River, 1993-95.

1993-94

| Ch/Code | Rel. Date | Stanfield Lest | | ODFW First | | Days | Stan. to ODFW Passage | | Total | |
|---------|-----------|----------------|-------|------------|-------|------|-----------------------|-------|-----------|-----|
| | | Date | Time | Date | Time | | Hrs/Min | Hours | Miles/Day | |
| 7/1 | 10/19/94 | 04/02/94 | 15:06 | 04/16/94 | 15:25 | 14 | 00:19 | 336.3 | 1.7 | |
| 7/13 | 12/07/94 | 01/15/94 | 12:49 | 01/25/94 | 21:46 | 10 | 08:57 | 249 | 2.3 | |
| 7/4 | 12/1 3194 | 01/10/94 | 19:06 | 01/16/94 | 16:32 | 5 | 21:26 | 141.4 | 4.0 | |
| 7/5 | 01/07/94 | 01/13/94 | 11:53 | 01/25/94 | 01:53 | 11 | 14:00 | 278 | 2.0 | |
| 7/6 | 01/10/94 | 03/11/94 | 17:57 | 03/28/94 | 22:30 | 17 | 04:33 | 412.6 | 1.4 | |
| 7/10 | 04/25/94 | 04/27/94 | 02:30 | 04/30/94 | 00:35 | 2 | 22:05 | 70.08 | 8.1 | |
| 7/13 | 03/11/94 | 03/15/94 | 12:59 | 03/26/94 | 04:32 | 10 | 15:33 | 255.6 | 2.2 | |
| 7/14 | 03/11/94 | 03/27/94 | 23:50 | 03/31/94 | 00:25 | 3 | 00:35 | 72.58 | 7.8 | |
| 7/17 | 03/24/94 | 03/30/94 | 19:06 | 04/02/94 | 02:53 | 1 | 07:47 | 55.78 | 10.2 | |
| 7/18 | 03/28/94 | 04/21/94 | 00:33 | 04/22/94 | 23:12 | 2 | 22:39 | 46.65 | 12.1 | |
| 7/23 | 04/04/94 | 04/07/94 | 03:58 | 04/09/94 | 06:25 | | 05:55 | 53.92 | 10.5 | |
| 7/26 | 04/11/94 | 04/17/94 | 00:22 | 05/02/94 | 22:00 | 15 | 18:02 | 378 | 1.5 | |
| 7/27 | 04/14/94 | 04/17/94 | | 04/18/94 | 19:58 | 1 | 19:36 | 43.6 | 13.0 | |
| | | | | | | | | Avg: | 184.1 | 5.9 |

1994-95

| Ch/Code | Rel. Date | Stanfield Last | | ODFW First | | Days | Stan. to ODFW Passage | | Total | |
|---------|-----------|----------------|-------|------------|-------|------|-----------------------|-------|-----------|-----|
| | | Date | Time | Date | Time | | Hrs/Min | Hours | Miles/Day | |
| 7/39 | 11/10/94 | 01/13/95 | 02:21 | 01/27/95 | 19:39 | 33 | 17:18 | 353.3 | 1.6 | |
| 7/40 | 11/17/94 | 01/16/95 | 13:45 | 02/19/95 | 10:24 | | 20:39 | 812.6 | 0.7 | |
| 7/45 | 11/30/94 | 02/07/95 | 07:43 | 02/24/95 | 12:31 | 17 | 04:48 | 412.8 | 1.4 | |
| 7/85 | 03/14/95 | 04/01/95 | 20:30 | 04/06/95 | 21:34 | 5 | 01:24 | 121.1 | 4.7 | |
| 7/88 | 03/13/95 | 03/26/95 | 12:11 | 03/29/95 | 12:59 | 3 | 00:48 | 72.8 | 7.8 | |
| 7/81 | 03/06/95 | 03/31/95 | 03:27 | 04/03/95 | 10:06 | 3 | 06:39 | 78.65 | 7.2 | |
| 7/82 | 03/06/95 | 04/04/95 | 07:01 | 04/09/95 | 20:56 | 5 | 04:06 | 124.1 | 4.6 | |
| 7/13 | 03/30/95 | 04/21/95 | 07:7 | 04/26/95 | 20:42 | 5 | 02:13 | 122.2 | 4.6 | |
| | | | | | | | | Avg: | 262.2 | 4.1 |

Table F-5: Summer steelhead passage times (days, hours, minutes) and miles moved per day between the release site (Barnhart Nolin) and ODFW (RM 56), Upstream Transport Evaluation, Umatilla River, 1993-95.

1993-94

| Ch/Code | Rel. Site | Release | | ODFW First | | Days | Rel. Site to ODFW | | Total | |
|---------|-----------|----------|-------|------------|-------|------|-------------------|-------|-----------|-----|
| | | Date | Time | Date | Time | | Hrs/Min | Hours | Miles/Day | |
| 7/8 | Barnhart | 02/28/94 | 11:00 | 03/06/94 | 06:14 | 5 | 19:14 | 139.2 | 2.4 | |
| 7/10 | Nolin | 03/09/94 | 11:00 | 03/13/94 | 03:29 | 3 | 16:29 | 88.48 | 6.1 | |
| 7/12 | Barnhart | 03/10/94 | 11:10 | 03/13/94 | 20:47 | 3 | 09:37 | 81.62 | 4.1 | |
| 7/15 | Nolin | 03/14/94 | 11:00 | 03/24/94 | 02:41 | 9 | 15:41 | 231.7 | 2.3 | |
| 7/16 | Barnhart | 03/22/94 | 10:40 | 03/24/94 | 13:36 | 2 | 02:56 | 50.93 | 6.5 | |
| 7/21 | Nolin | 03/31/94 | 10:50 | 04/02/94 | 18:58 | 2 | 08:08 | 56.13 | 9.6 | |
| | | | | | | | | Avg: | | 5.2 |

1994-95

| Ch/Code | Rel. Site | Release | | ODFW First | | Days | Rel. Site to ODFW | | Total | |
|---------|-----------|----------|-------|------------|-------|------|-------------------|-------|-----------|-----|
| | | Date | Time | Date | Time | | Hrs/Min | Hours | Miles/Day | |
| 7/49 | Nolin | 02/27/95 | 11:00 | 03/27/95 | 19:53 | 28 | 08:53 | 680.9 | 0.5 | |
| 7/6 | Nolin | 03/27/95 | 11:30 | 03/31/95 | 20:11 | 4 | 08:41 | 104.7 | 3.2 | |
| 7/20 | Barnhart | 04/07/95 | 10:45 | 04/11/95 | 20:55 | 4 | 10:10 | 106.2 | 3.1 | |
| 7/38 | Barnhart | 11/10/94 | 10:30 | 01/29/95 | 23:21 | 80 | 1251 | 1933 | 0.2 | |
| | | | | | | | | Avg: | | 1.7 |

Table F--6: Spring Chinook Salmon passage times (days, hours, minutes) and miles moved per day between the release site (Barnhart) and ODFW (RM 56). Upstream Transport Evaluation, Umatilla River, 1993-94.

1993-94

| Ch/Code | Rel. Site | Release | | ODFW First | | Days | Rel. Site to ODFW | | Total | |
|---------|-----------|----------|-------|------------|-------|------|-------------------|-------|-----------|-----|
| | | Date | Time | Date | Time | | Hrs/Min | Hours | Miles/Day | |
| 13/21 | Barnhart | 05/02/94 | 11:30 | 05/05/94 | 23:01 | 3 | 11:31 | 83.52 | 4.0 | |
| 13/22 | Barnhart | 05/06/94 | 11:00 | 05/10/94 | 03:28 | 3 | 16:28 | 68.47 | 3.7 | |
| 13/44 | Barnhart | 05/10/94 | 13:30 | 05/12/94 | 23:03 | 2 | 09:33 | 57.55 | 5.8 | |
| 13/15 | Barnhart | 05/13/94 | 15:00 | 05/16/94 | 01:19 | 2 | 10:19 | 56.32 | 5.7 | |
| | | | | | | | | Avg: | 71.96 | 4.8 |

Table F-7: Spring Chinook Salmon passage times (days, hours, minutes) and miles moved per day between Stanfield Dam and ODFW (RM 56), Passage Evaluation, Umatilla River, 1993-95.

1993-94

| Ch/Code | Rel. Date | Stanfield Last | | ODFW First | | Days | Stan. to ODFW Passage | | Total | |
|---------|-----------|----------------|-------|------------|-------|------|-----------------------|-------|-----------|-----|
| | | Date | Time | Date | Time | | Hrs/Min | Hours | Miles/Day | |
| 13/14 | 04/14/94 | 04/20/94 | 10:20 | 04/24/94 | 07:34 | 3 | 21:14 | 93.23 | 6.1 | |
| 13/17 | 04/27/94 | 05/06/94 | 04:41 | 05/08/94 | 22:03 | 2 | 17:22 | 65.37 | 8.7 | |
| 13/18 | 04/29/94 | 05/23/94 | 17:39 | 05/25/94 | 17:06 | 1 | 23:27 | 47.45 | 11.9 | |
| | | | | | | | | Avg: | 68.66 | 8.9 |

1994-95

| Ch/Code | Rel. Date | Stanfield Last | | ODFW First | | Days | Stan. to ODFW Passage | | Total | |
|-------------------|-----------|----------------|-------|------------|-------|------|-----------------------|-------|-----------|-----|
| | | Date | Time | Date | Time | | Hrs/Min | Hours | Miles/Day | |
| 13/32 | 04/10/95 | 04/24/95 | 16:40 | 05/08/95 | 04:11 | 13 | 11:31 | 323.5 | 1.8 | |
| 13/34 | 04/11/95 | 04/25/95 | 14:00 | 05/03/95 | 18:21 | 8 | 04:21 | 196.4 | 2.9 | |
| 13/36 | 04/13/95 | 04/25/95 | 13:04 | 05/16/95 | 02:44 | 20 | 13:40 | 493.7 | 1.1 | |
| 13/38 | 04/18/95 | 04/25/95 | 01:39 | 05/08/95 | 16:14 | 13 | 14:35 | 326.6 | 1.7 | |
| 13/40 | 04/20/95 | 04/24/95 | 08:57 | 04/27/95 | 04:07 | 2 | 19:10 | 67.17 | 8.4 | |
| 13/31 | 04/24/95 | 04/28/95 | 23:39 | 05/18/95 | 18:34 | 19 | 18:55 | 474.9 | 1.2 | |
| 13/35 | 04/13/95 | 05/02/95 | 11:35 | 05/19/95 | 02:23 | 16 | 14:48 | 398.8 | 1.4 | |
| 13/43 | 04/24/95 | 05/22/95 | 11:14 | 05/24/95 | 13:39 | 2 | 02:25 | 50.42 | 11.2 | |
| 13/42 | 04/26/95 | 05/19/95 | 14:36 | 05/24/95 | 02:51 | 4 | 12:15 | 108.2 | 5.2 | |
| file name: 9395#2 | | | | | | | | Avg: | 271.1 | 3.9 |

Table F-8. Fall chinook salmon mainstem passage data at John Day, McNary, and Ice Harbor Dams, 1990-93.

| Year | Dam | Aug 1-15 | | Aug 16-31 | | Sep 1-15 | | Sep 16-30 | | Oct 1-15 | | Oct 16-31 | | Total No. |
|-------|------------|----------|-----|-----------|-----|----------|------|-----------|------|----------|------|-----------|------|-----------|
| | | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % | |
| 1990 | John Day | 2147 | 2.3 | 11223 | 12 | 49115 | 52.7 | 22393 | 24 | 6663 | 7.1 | 1652 | 1.8 | 93193 |
| | McNary | 2686 | 3.3 | 4504 | 5.5 | 40375 | 49.2 | 21343 | 26 | 10037 | 12.2 | 3053 | 3.7 | 81998 |
| | Ice Harbor | 102 | 1.9 | 202 | 3.7 | 1716 | 31.8 | 1598 | 29.6 | 1169 | 21.7 | 604 | 11.2 | 5391 |
| 1991 | John Day | 1132 | 1.4 | 3653 | 4.5 | 34358 | 42.7 | 30592 | 38 | 8434 | 10.5 | 2341 | 2.9 | 80510 |
| | McNary | 1340 | 1.8 | 2832 | 3.8 | 25055 | 33.9 | 31196 | 42.2 | 10638 | 14.4 | 2872 | 3.9 | 73933 |
| | Ice Harbor | 87 | 1.4 | 54 | 0.9 | 1989 | 32.5 | 2064 | 33.7 | 1367 | 22.3 | 563 | 9.2 | 6124 |
| 1992 | John Day | 1225 | 1.7 | 6320 | 8.6 | 33363 | 45.5 | 24777 | 33.8 | 6160 | 8.4 | 1413 | 1.09 | 73258 |
| | McNary | 1470 | 2.1 | 4294 | 6 | 26679 | 37.3 | 25282 | 35.3 | 11602 | 16.2 | 2280 | 3.2 | 71607 |
| | Ice Harbor | 67 | 1.2 | 156 | 2.8 | 1732 | 31.1 | 1984 | 35.6 | 1078 | 19.3 | 556 | 10 | 5573 |
| 1993 | John Day | 1761 | 2.6 | 8828 | 13 | 29623 | 43.9 | 22044 | 32.7 | 3805 | 5.6 | 1411 | 2.1 | 67472 |
| | McNary | 2137 | 3.3 | 6098 | 9.5 | 28042 | 43.6 | 20051 | 31.2 | 6182 | 9.6 | 1820 | 2.8 | 64327 |
| | Ice Harbor | 132 | 4.1 | 199 | 6.2 | 988 | 30.7 | 1099 | 34.1 | 539 | 16.7 | 262 | 8.1 | 3219 |
| Total | John Day | 6265 | 2 | 30024 | 9.5 | 146459 | 46.6 | 99806 | 31.7 | 25062 | 8 | 6817 | 2.2 | 314433 |
| | McNary | 7630 | 2.6 | 17728 | 6.1 | 12011 | 41.2 | 97872 | 33.5 | 38459 | 13.2 | 10025 | 3.4 | 291865 |
| | Ice Harbor | 388 | 1.9 | 611 | 3 | 6425 | 31.6 | 6745 | 33.2 | 4153 | 20.5 | 1985 | 9.8 | 20307 |

F-6

file name: chfmnstm

Table F-9: Percent of Fall Chinook Salmon homing to the Umatilh River versus straying into fish hatcheries and spawning grounds above McNary Dam. Average attraction flows exiting the Umatilla River during September are also included. Numbers represent estimated coded-wire tag recoveries.

| Recovery Year | No. Above McNary | No. to Uma. R. | Total No. | Percent Home | Percent Stray | Avg. Flow Sept 1-15 | Avg. Flow Sept 16-30 |
|---------------|------------------|----------------|-----------|--------------|---------------|---------------------|----------------------|
| 1990 | 152 | 223 | 375 | 59.5 | 41 | 4 cfs | 21 cfs |
| 1991 | 182 | 145 | 327 | 44.3 | 56 | 50 cfs | 130 cfs |
| 1992 | 92 | 29 | 121 | 24 | 76 | 1.5 cfs | 1 cfs |
| 1993 | 67 | 39 | 106 | 36.8 | 83 | 78 cfs | 100 cfs |
| 1994 | 88 | 110 | 198 | 55.6 | 44 | 59 cfs | 62 cfs |

Table F-10. Umatilla River fall chinook salmon homing and straying rates for acclimated (Minthom) versus direct (near Minthom) releases. Numbers represent estimated coded-wire tag recoveries.

| Brood Yr. Tag Code | Rel. Loc. | No. Tagged | Rel. Age | No. Above McNary | No. to Uma. R. | Percent Home | Percent Stray |
|--------------------|-------------|------------|----------|------------------|----------------|--------------|---------------|
| 87 539-41 | Minthom | 13260 | o++ | 6 | 2 | 25.0 | 75.0 |
| 8 7 536-38 | Nr. Minthom | 73148 | o++ | 24 | 49 | 67.1 | 32.9 |
| 8 8 753,54,57 | Minthom | 76824 | o++ | 11 | 13 | 54.2 | 45.8 |
| 8 8 758,60,63 | Nr. Minthom | 76425 | o++ | 11 | 9 | 45.0 | 55.0 |
| 89 325-27 | Minthom | 66426 | o++ | 2 | 7 | 77.8 | 22.2 |
| 89 322-24 | Nr. Minthom | 70450 | o++ | 4 | 1 | 20.0 | 80.0 |
| 9 0 563,601,602 | Minthom | 76411 | o+ | 15 | 15 | 50.0 | 50.0 |
| 90 560-62 | Nr. Minthom | 73454 | o+ | 20 | 14 | 41.2 | 58.8 |

file name: 9495chfl

Table F-I 1: Umatilla River homing and straying data for yearling (1 +) fall chinook salmon (includes acclimated and direct releases). Numbers represent estimated coded-wire tag recoveries.

| Brood Yr. | Tag Code | Rel. Loc. | No. Tagged | Rel. Age | No. Above McNary | No. To U m a . R. | % home | % stray |
|-----------|------------|-----------|------------|----------|------------------|-------------------|--------|---------|
| 84 | 073327 | Bon/Minth | 88396 | 1+ | 101 | 55 | 35.3 | 64.7 |
| 85 | 073823-27 | Minthom | 49635 | 1+ | 53 | 100 | 65.4 | 34.6 |
| 85 | 073828-32 | Bonifer | 50492 | 1+ | 36 | 63 | 63.6 | 36.4 |
| 86 | 074038-39 | Minthorn | 81046 | 1+ | 67 | 234 | 77.7 | 22.3 |
| 86 | 074036-37 | Bonifer | 77914 | 1+ | 39 | 170 | 81.3 | 18.7 |
| 91 | 071460,461 | RM 73.5 | 47102 | 1+ | 1 | 5 | 83.3 | 16.7 |

Table F-12: Umatilla River homing and straying data for sub-yearling (O+,O+ +) fall chinook salmon (includes acclimated and direct releases). Numbers represent estimated coded-wire tag recoveries.

| Brood Yr. | Tag Code | Rel. Loc. | No. Tagged | Rel. Age | No. Above McNary | No. To U m a . R. | % home | % stray |
|-----------|---------------|-------------|------------|----------|------------------|-------------------|--------|---------|
| 89 | 075403-05 | RM 70-79 | 159020 | o+ | 46 | 27 | 37.0 | 63.0 |
| 89 | 075325-27 | Minthorn | 66426 | 0++ | 2 | 24 | 92.3 | 7.7 |
| 89 | 075322-24 | Nr. Mintorn | 70450 | 0++ | 4 | 1 | 20.0 | 80.0 |
| 90 | 075563,601-02 | Minthom | 76411 | 0+ | 16 | 9 | 36.0 | 64.0 |
| 90 | 075560-62 | Nr. Minthom | 73454 | 0+ | 20 | 14 | 41.2 | 58.8 |
| 91 | 071429-38 | RM 42.5 | 304968 | 0+ | 0 | 2 | 100.0 | 0.0 |
| 90 | 075225-26 | RM 70-79 | 103980 | 0+ | 15 | 18 | 54.5 | 45.5 |
| 90 | 075328 | RM 70-79 | 48266 | 0+ | 14 | 13 | 48.1 | 51.9 |
| 90 | 075449,50,51 | RM 70-79 | 152739 | 0+ | 33 | 38 | 53.5 | 46.5 |
| 90 | 070016 | RM 70-79 | 48301 | 0+ | 13 | 7 | 35.0 | 65.0 |

file name:9495chf2

Table F-13: Umatilla River homing and straying data for coho salmon. Numbers represent estimated coded-wire tag recoveries only.

| Brood Yr. | Tag Code | No. Tagged | Rel. Location | No. to Uma. R. | No. to Cascade | No. to Other | Percent Home | Percent Stray |
|-----------|-----------|------------|---------------|----------------|----------------|--------------|--------------|---------------|
| 67 | 074809 | 27062 | Nr. Minthom | 19 | 4 | 0 | 82.6 | 17.4 |
| 87 | 74610-11 | 53155 | Minthom | 75 | 18 | 2 | 78.9 | 21.1 |
| 88 | 074814-1s | 55259 | Minthom | 175 | 93 | 32 | 58.3 | 41.7 |
| 88 | 074813 | 26881 | RM 63-70 | 72 | 31 | 5 | 66.7 | 33.3 |
| 89 | 075535 | 24584 | Minthom | 6 | 0 | 0 | 100.0 | 0.0 |
| 89 | 075534 | 25338 | RM 56-60 | 8 | 3 | 0 | 72.7 | 27.3 |
| 89 | 075533 | 25407 | RM 63-70 | 12 | 0 | 0 | 100.0 | 0.0 |
| 90 | 075620 | 27908 | RM 56 | 45 | 12 | 2 | 76.3 | 23.7 |
| 90 | 075621-22 | 55163 | RM 60 | 119 | 31 | 4 | 77.3 | 22.7 |
| 91 | 071521 | 28273 | RM 60 | 36 | 0 | 0 | 100.0 | 0.0 |
| 91 | 07X22-23 | 55895 | RM 42 | 76 | 0 | 0 | 100.0 | 0.0 |

Table F-14: Umatilla River coho salmon homing and straying data for acclimated versus direct releases. Numbers represent estimated coded-wire tag recoveries.

| Brood Yr. | Tag Code | No. Tagged | Rel. Location | No. to Uma. R. | No. to Other | Total No. | Percent Home | Percent Stray |
|-----------|----------|------------|---------------|----------------|--------------|-----------|--------------|---------------|
| 87 | 074609 | 27062 | Nr. Minthom | 19 | 4 | 23 | 41.3 | 58.7 |
| 87 | 074610 | 26416 | Minthom | 37 | 8 | 45 | 41.1 | 58.9 |
| 87 | 074611 | 26739 | Minthom | 36 | 12 | 50 | 38.0 | 62.0 |
| 88 | 074614 | 28033 | Minthom | 81 | 4s | 129 | 31.4 | 68.6 |
| 88 | 074813 | 26881 | Nr. Minthom | 72 | 36 | 108 | 33.3 | 66.7 |
| 88 | 074815 | 27226 | Minthom | 94 | 77 | 171 | 27.5 | 72.5 |
| 89 | 075535 | 24584 | Minthom | 6 | 0 | 6 | 50.0 | 50.0 |
| 89 | 075534 | 25905 | RM 56-60 | 8 | 3 | 11 | 36.4 | 63.6 |
| 89 | 075533 | 24851 | RM 63-70 | 12 | 0 | 12 | 50.0 | 50.0 |

file name: 9495chol

Table F–15: Percent of Spring Chinook Salmon homing to the Umatilla River versus straying into fish hatcheries and spawning grounds above and below McNary Dam. Numbers represent estimated coded–wire tag recoveries.

| Recovery Year | No. Above McNary | No. to Uma. R. | No. to Other | Total No. | Percent Home | Percent Stray |
|---------------|------------------|----------------|--------------|-----------|--------------|---------------|
| 1990 | 9 | 770 | 4 | 783 | 98.3 | 9.5 |
| 1991 | 0 | 710 | 1 | 711 | 99.9 | 0.1 |
| 1992 | 22 | 326 | 3 | 351 | 92.9 | 22.9 |
| 1993 | 17 | 753 | 1 | 771 | 97.7 | 17.1 |
| 1994 | 13 | 157 | 0 | 170 | 92.4 | 13.0 |


file name: 9495chs1

Figure F-1

Summer Steelhead Mean Passage Times

for Westland, Feed, and Stanfield Diversion Dams

Umatilla River, 1993-95



File name: avg9495




Figure F-2. Summer Steelhead migrational routes for Westland, Feed and Stanfield Dams, 1993-95.
File name: ladders

Figure F-3. Radio telemetry data depicting average migrational times (hours and minutes) for Summer Steelhead between dams versus passage times over dams, Umatilla River 1993-1995.





Figure F-4

Spring Chinook Mean Passage Times for Westland, Feed, and Stanfield Diversion Dams Umatilla River, 1993-95



F-14






Figure F-5. Spring Chinook migrational routes for Westland, Feed and Stanfield Dams, 1993-95.

File name: ladder#1

Figure F-6

Summer Steelhead Migrational Behavior Umatilla River 1993-94

F-16




Passage Evaluation
Release site RM4-3MD
File Name sts9394

Figure F-7

Summer Steelhead Migrational Behavior Umatilla River 1994-95

F-17




Channel/Code
- 7 1 3 9 - □ 7/40 - 7 1 4 2 - ◆ 7/81

Passage Evaluation
Release site RM4-3MD
File Name sts9495

Figure F-8

Fall Chinook and Coho Returns Versus Flows Umatilla River 1992

F-18



Flows measured at Umatilla
File name: 92chfflw




Figure F-9

Fall Chinook and Coho Returns Versus Flows Umatilla River 1993

F-19





Figure F- 10

Fall Chinook and Coho Returns Versus Flows Umatilla River 1994

F-20







Figure F-1 1

Summer Steelhead Returns Versus Flows Umatilla River 1992-93


F-21



Flows measured at Umatilla
File name: 83stflw

Figure F- 12


Summer Steelhead Returns Versus Flows Umatilla River 1993-94



F-22

Summer Steelhead Returns Versus Flows Umatilla River 1994-95

Figure F-13



Date

% freq - flows

Flows measured at Umatilla
File name: 95btrflw

Figure F- 14

Spring Chinook Salmon Versus Flows Umatilla River 1993

F-24





Figure F-15

Spring Chinook Salmon Versus Flows Umatilla River 1994





CHS — flows

File name:

Figure F-16

Spring Chinook Salmon Versus Flows Umatilla River 1995

Flows measured at Umatilla
File name: 95chaflw



APPENDIX G
Spawning Survey Data for 1993-1994

Table G-1. Summary of Summer Steelhead Escapement Surveys, Umatilla River Basin, 1995.

| Redd # | River Mile | Area Description | Habitat Type | Date Observed | Total Steelhead On Redds On That Date | Total Steelhead Holding On That Date |
|--|------------|---|--------------|---------------|---------------------------------------|--------------------------------------|
| CAMP CREEK - MOUTH TO FORKS TO WATERFALL | | | | | | |
| Surveyed March 10, 28*, April 17 | | | | | | |
| 1 | 0.4 | Red Cabin | Riffle | 3/28 | 1 | |
| 2 | 1.0 | | Tailout | 4/17 | | |
| 3 | 1.5 | | Tailout | 4/17 | | |
| 4 | 1.6 | | Tailout | 4/17 | | |
| 5 | 2.9 | | Tailout | 4/17 | | 1 |
| NORTH FORK MEACHAM CREEK - MOUTH TO POT CREEK (5.0 Miles) | | | | | | |
| Surveyed March 30*, April 3* | | | | | | |
| 6 | 1.7 | | | 3/30 | 3 | |
| 7 | 2.0 | | | 3/30 | | |
| 8 | 2.1 | | Tailout | 3/30 | | |
| 9 | 2.5 | Small anabranch | Riffle | 3/30 | | |
| 10 | 2.7 | Small anabranch | Riffle | 3/30 | | |
| 11 | 3.1 | Cabin above Bear Creek | Riffle | 3/30 | | |
| 12 | 3.2 | 100 yards above cabin | Riffle | 3/30 | | |
| 13 | 3.3 | 141 yards above cabin | Riffle | 3/30 | | |
| 14 | 3.4 | 175 yards above cabin | Tailout | 3/30 | | |
| 15 | 4.0 | .5 miles above Forest Service upper fence | Riffle | 3/30 | | |
| 16 | 4.0 | 8 yards above redd #15 | Riffle | 3/30 | | |
| 17 | 4.1 | .6 miles above Forest Service upper fence | Tailout | 3/30 | | |
| 18 | 4.1 | 100 yards upstream of redd #17 | Tailout | 3/30 | | |
| 19 | 4.8 | | Riffle | 4/3 | | |

Table G- 1. Continued

| Redd # | River Mile | Area Description | Habitat Type | Date Observed | Total Steelhead On Redds On That Date | Total Steelhead Holding On That Date |
|--|------------|--|--------------|---------------|---------------------------------------|--------------------------------------|
| SQUAW CREEK - MOUTH TO LITTLE SQUAW CREEK (6.7 Miles) | | | | | | |
| Surveyed March 8, 27, 28, April 4, 5, 6, May 18 | | | | | | |
| 20 | 0.2 | 50 yard below Highway Bridge | Riffle | 4/6 | 19 | 2 |
| 21 | 0.3 | Old pipe trap site | Riffle | 4/6 | | |
| 22 | 0.4 | 250 yards above Highway Bridge | Tailout | 4/6 | | |
| 23 | 0.4 | 300 yards above Highway Bridge | Riffle | 4/6 | | |
| 24 | 0.5 | Below Walt Farrow's (WF) house | Riffle | 4/6 | | |
| 25 | 0.5 | Below WF house | Riffle | 4/6 | | |
| 26 | 0.6 | Same area as redd #25 | Riffle | 4/6 | | |
| 27 | 0.6 | Below WF house, redd not visible after high water of 3/14-23 | Riffle | 3/8 | 3 | |
| 28 | 0.9 | 175 yards below WF house | Tailout | 4/6 | | |
| 29 | 1.0 | 70 yards below WF house | Riffle | 4/6 | | |
| 30 | 1.0 | 50 yards below WF house | Riffle | 4/6 | | |
| 31 | 1.2 | 10 yards below Bedrock Falls above WF house | Riffle | 4/6 | | |
| 32 | 1.3 | In anabranch | Riffle | 3/8 | | |
| 33 | 1.3 | Mile 1.9 below Bachelor canyon | Riffle | 4/6 | | |
| 34 | 1.3 | Mile 1.9 below Bachelor canyon | Riffle | 4/6 | | |
| 35 | 1.6 | 20 yards below redd #34 | Riffle | 4/6 | | |
| 36 | 1.6 | 1.6 miles below Bachelor canyon | Riffle | 4/6 | | |
| 37 | 1.7 | 1.5 miles below Bachelor canyon | Riffle | 4/6 | | |
| 38 | 1.7 | Visible after high water of 3/14-23 | Riffle | 3/8 | | |
| 39 | 1.9 | 41 yards above falls - not visible after high water of 3/14-23 | Riffle | 3/8 | | |
| 40 | 1.9 | 80 yards above falls | Riffle | 4/6 | | |
| 41 | 2.1 | 303 yards below Cliff Picard's old cabin | Riffle | 3/8 | | |
| 42 | 2.1 | 300 yards below Cliff Picard's old cabin | Tailout | 4/6 | | |
| 43 | 2.5 | 200 yards below old log cabin with silver roof | Riffle | 3/28 | 4 | 2 |

Table G-1. Continued

| Redd # | River Mile | Area Description | Habitat Type | Date Observed | Total Steelhead On Redds On That Date | Total Steelhead Holding On That Date |
|---------------|-------------------|---|---------------------|----------------------|--|---|
| 44 | 2.6 | 16 yards below old log cabin with silver roof | Riffle | 3/28 | | |
| 45 | 2.6 | Across from old log cabin with silver roof | Riffle | 4/6 | | |
| 46 | 2.8 | 200 yards below new log home | Tailout | 4/6 | | |
| 47 | 3.1 | 150 yards below Bachelor canyon | Riffle | 4/6 | | |
| 48 | 3.2 | 100 yards below Bachelor canyon | Riffle | 4/6 | | |
| 49 | 3.5 | 507 yards above Bachelor canyon | Riffle | 4/6 | | |
| 50 | 4.0 | 50 yards below first crossing | Tailout | 3/27 | | |
| 51 | 4.0 | 33 yards above first crossing | Tailout | 3/8 | | |
| 52 | 4.1 | 150 yards above first crossing | Riffle | 4/6 | | |
| 53 | 4.1 | 175 yards above first crossing | Tailout | 3/27 | | |
| 54 | 4.1 | 200 yards above first crossing | Riffle | 3/27 | | |
| 55 | 4.2 | 250 yards above first crossing | Tailout | 3/27 | | |
| 56 | 5.0 | 100 yards above 2nd crossing - not visible after high water of 3/14-23 | Riffle | 3/8 | | |
| 57 | 5.1 | 125 yards above second crossing | Riffle | 3/27 | | |
| 58 | 5.2 | Third crossing - redd not visible - truck drove over | Riffle | 3/8 | | |
| 59 | 5.5 | 500 yards above third crossing | Tailout | 3/27 | | |
| 60 | 6.0 | 75 yards above excellent old spawning area - not visible after high water | Riffle | 3/8 | | |
| 61 | 6.0 | 150 yards above excellent old spawning area | Riffle | 5/18 | | |
| 62 | 6.5 | Big pool on corner - 300 yards below Little Squaw Creek | Tailout | 3/8 | | |
| 63 | 6.5 | Spawning in same place as redd #62 | Tailout | 4/6 | | |
| 64 | 6.7 | 100 yards below Little Squaw Creek confluence | Riffle | 3/8 | | |

Table G-1. Continued

| Redd # | River Mile | Area Description | Habitat Type | Date Observed | Total Steelhead On Redds On That Date | Total Steelhead Holding On That Date |
|--|------------|---|--------------|---------------|---------------------------------------|--------------------------------------|
| BUCKAROO CREEK - MOUTH TO 3.0 MILES UPSTREAM Surveyed March 9, 24, April 14 | | | | | | |
| 65 | 0.0 | 23 yards above mouth - not visible -high w. of 3/14-23 | Riffle | 3/9 | | |
| 66 | 0.3 | Across from yellow house | Riffle | 3/9 | | |
| 67 | 0.6 | 200 yards above first road crossing - not visible • high water of 3/14-23 | Riffle | 3/9 | | |
| 68 | 0.6 | 75 yards above redd #67 | Tailout | 3/24 | 1 | |
| 69 | 1.1 | Falls pool - not visible - high water of 3/14-23 | Tailout | 3/9 | | |
| 70 | 1.1 | Falls pool | Tailout | 3/24 | | |
| NORTH FORK - UMATILLA RIVER - MOUTH TO 1.5 MILES ABOVE COYOTE CREEK Surveyed March 29*, April 19 | | | | | | |
| 71 | 0.1 | NF Gage | Tailout | 3/29 | | |
| SOUTH FORK - UMATILLA RIVER - MOUTH TO 1.0 MILES ABOVE SHIMMIEHORN CREEK Surveyed March 29* | | | | | | |
| 72 | 0.9 | 0.9 miles above mouth | Riffle | 3/29 | | |
| 73 | 1.2 | 1.2 miles above mouth | Riffle | 3/29 | | 2 |
| 74 | 1.2 | 1.2 miles above mouth | Riffle | 3/29 | | |
| 75 | 1.4 | 1.4 miles above mouth | Riffle | 3/29 | | |
| MEACHAM CREEK - MOUTH TO 18.2 MILES UPSTREAM Surveyed April 18* | | | | | | |
| 76 | 13.9 | | Riffle | 4/18 | 4 | 1 |
| 77 | 13.6 | NF railroad bridge | Riffle | 4/18 | | |
| 78 | 13.5 | | Riffle | 4/18 | | |
| 79 | 13.5 | | Riffle | 4/18 | | |
| 80 | 13.1 | | Riffle | 4/18 | | |
| 81 | 12.3 | 200 yards above white RR switch building | Riffle | 4/18 | | |
| 82 | 12.3 | 50 feet downstream | Riffle | 4/18 | | |
| 83 | 12.2 | 100 yards downstream | Riffle | 4/18 | | |
| 84 | 11.2 | .5 miles above Duncan | Riffle | 4/18 | | |
| 85 | 11.1 | | Riffle | 4/18 | | |
| 86 | 10.8 | | Riffle | 4/18 | | |
| 87 | 10.7 | Duncan | Riffle | 4/18 | | |
| BOSTON CANYON CREEK - MOUTH TO FORKS, Surveyed March 13, 21, 31, April 14 | | | | | | |
| MINTHORN SPRINGS CREEK, Surveyed April 14 | | | | | | |
| MAINSTEM - RM 80.0 -74.5, Surveyed May 30 | | | | | | |

* Partial Survey

Table G-2. Comparison of Umatilla River Adult Summer Steelhead Released above Three Mile Falls Dam, Redds and Redds per Mile surveyed, 1985 - 1995 (* estimated).

| YEAR | STEELHEAD ESCAPEMENT | | REDDS OBSERVED | MILES SURVEYED | REDDS PER MILE SURVEYED |
|------|----------------------|----------|-----------------|----------------|-------------------------|
| | NATURAL | HATCHERY | | | |
| 1985 | 3197* | 0 | 33 | 23.5 | 1.4 |
| 1986 | 2885* | 0 | 134 | 20.9 | 6.4 |
| 1987 | 3444 | 0 | 156 | 52.5 | 3.0 |
| 1988 | 2144 | 160 | 275 | 61.0 | 4.5 |
| 1989 | 1934 | 353 | 128 | 50.2 | 2.5 |
| 1990 | 1290 | 102 | High Water | High Water | High Water |
| 1991 | 623 | 234 | High Water | High Water | High Water |
| 1992 | 2007 | 315 | 300 | 67.2 | 4.4 |
| 1993 | 1166 | 455 | 51 - High Water | 46.6 | High Water |
| 1994 | 852 | 252 | 235 | 75.6 | 3.1 |
| 1995 | 784 | 530 | 126 | 35.3 | 3.6 |

Table G-3. Summary of Summer Steelhead Escapement Survey Data in the Umatilla River Basin, 1985-1995.

| Year | Squaw Creek | | | Buckaroo Creek | | | Meacham Creek | | | NF Meacham Creek | | | Camp Creek | | | Boston Canyon Creek | | | NF Umatilla | | |
|--|---------------------------------------|-----|-------|----------------|-----|-------|---------------|-----|-------|------------------|-----|-------|------------|-----|-------|---------------------|-----|-------|-------------|-----|-------|
| | Redds | STS | Miles | Redds | STS | Miles | Redds | STS | Miles | Redds | STS | Miles | Redds | STS | Miles | Redds | STS | Miles | Redds | STS | Miles |
| 1985 | 14 | 3 | 5.0 | 2 | 0 | 2.0 | 0 | 0 | 1.5 | 1 | 8 | 3.0 | 4 | 2 | 2.5 | 10 | 9 | 1.0 | | | |
| 1986 | 25 | 0 | 3.5 | 3 | 0 | 2.0 | 49 | 2 | 6.4 | 27 | 0 | 3.0 | 8 | 7 | 2.5 | 8 | 0 | 1.0 | | | |
| 1987 | 25 | 13 | 6.6 | 0 | 0 | 2.0 | 49 | 0 | 9.0 | 7 | 2 | 3.0 | 12 | 3 | 2.5 | 0 | 0 | 1.0 | 6 | 2 | 2.5 |
| 1988 | 95 | 0 | 6.6 | 20 | 3 | 3.5 | 51 | 1 | 9.0 | 10 | 0 | 3.0 | 6 | 0 | 2.5 | 2 | 0 | 1.0 | 1 | 0 | 2.5 |
| 1989* | 46 | 0 | 6.6 | 10 | 2 | 3.5 | 24 | 0 | 9.0 | 4 | 2 | 3.0 | 1 | 0 | 4.0 | 9 | 0 | 1.0 | 3 | 0 | 1.5 |
| 1990 | High water and poor survey conditions | | | | | | | | | | | | | | | | | | | | |
| 1991 | High water and poor survey conditions | | | | | | | | | | | | | | | | | | | | |
| 1992 | 77 | 10 | 6.7 | 5 | 0 | 3.0 | 120 | 39 | 18.0 | 30 | 18 | 5.0 | 8 | 9 | 2.5 | 0 | 0 | 1.0 | 17 | 3 | 2.5 |
| 1993* | 10 | 12 | 6.7 | 6 | 4 | 3.0 | 6 | 5 | 15.8 | 3 | 1 | 3.3 | 7 | 4 | 2.5 | 6 | 3 | 1.0 | | | |
| 1994 | 36 | 4 | 6.7 | 0 | 0 | 3.0 | 40 | 5 | 18.2 | 11 | 6 | 5.0 | 6 | 2 | 2.5 | 3 | 4 | 1.0 | 4 | 0 | 4.0 |
| 1995** | 45 | 21 | 6.7 | 6 | 1 | 3.0 | 12 | 5 | 3.1 | 14 | 3 | 5.0 | 5 | 1 | 2.5 | 0 | 0 | 1.0 | 1 | 1 | 2.0 |
| <p>NOTES: 1) Variability in areas surveyed, surveyors and survey conditions make direct comparisons of redd data difficult. 2) Steelhead observed were number observed during peak survey. 3) 1992 - Fifteen redds observed in mainstem not listed. 4) 1994 - Five redds observed in mainstem not listed. 5) *High water was believed to wash out some redds. 6) **High water after April 18 washed out redds previously marked - good surveys before the washout. 7) Steelhead redds have also been observed in the following tributaries that are not annually surveyed: Duncan Canyon Creek, East Fork Meacham Creek, Owsley Creek, Buck Creek, Thomas Creek, Moonshine and Westgate Canyon Creek.</p> | | | | | | | | | | | | | | | | | | | | | |
| <p>REAS PRESENTLY SURVEYED: Squaw Creek - Mouth to Little Squaw Creek Confluence - 6.7 miles Buckaroo Creek - Mouth to top of Timber Breakout Meadow - 3.0 miles Meacham Creek - Mouth to 18.2 miles upstream - Top of USFS Habitat Improvement Area North Fork Meacham Creek - Mouth to Pot Creek Confluence - 5.0 miles Camp Creek - Mouth to Large Fork - 2.5 miles Boston Canyon - Mouth to Forks - 1.0 miles North Fork Umatilla - Mouth to 1.0 miles above Coyote Creek - 4.0 miles South Fork Umatilla - Mouth to Forks - 3.2 miles Ryan Creek - Mouth to 3.0 miles upstream - 3.0 miles (lower .3 miles not currently surveyed - private land) Minthom Springs - Mouth to Confluence of Umatilla - .3 miles Pearson Creek - Mouth to 6.0 miles upstream - Culvert Crossing - 6.0 miles West Birch Creek - Bridge Creek to RM 16.0 East Birch Creek - RM 8.5- RM 15.0</p> | | | | | | | | | | | | | | | | | | | | | |

Table G-3. Continued

| Year | South Fork Umatilla | | | Ryan Creek | | | Minthorn Springs | | | Pearson Creek | | | West Birch Creek | | | East Birch Creek | | |
|--------|---------------------------------------|-----|-------|------------|-----|-------|------------------|-----|-------|---------------|-----|-------|------------------|-----|-------|------------------|------|-------|
| | Redds | STS | Miles | Redds | STS | Miles | Redds | STS | Miles | Redds | STS | Miles | Redds | STS | Miles | Redds | STS | Miles |
| 1985 | | | | 2 | 0 | 2.0 | | | | | | | | | | | | |
| 1986 | | | | 13 | 0 | 2.0 | | | | | | | | | | | | |
| 1987 | 3 | 0 | 3.0 | 10 | 0 | 2.0 | | | | 22 | 0 | 6.0 | | | 11 | 0 | 5.5 | |
| 1988 | 5 | 1 | 2.0 | 9 | 0 | 2.0 | | | | 15 | 13 | 6.0 | | | 39 | 10 | 11.0 | |
| 1989* | 7 | 0 | 2.0 | 16 | 0 | 3.0 | | | | | | | | | | | | |
| 1990 | High water and poor survey conditions | | | | | | | | | | | | | | | | | |
| 1991 | High water and poor survey conditions | | | | | | | | | | | | | | | | | |
| 1992 | 15 | 9 | 4.2 | 3 | 0 | 2.0 | 5 | 0 | .2 | 1 | 1 | 6.0 | 0 | 0 | 4 | 0 | 1.0 | |
| 1993* | 8 | 4 | 4.2 | | | | | | | 3 | 5 | 8.0 | 3 | 0 | 11 | 2 | 4.5 | |
| 1994 | 8 | 0 | 4.2 | 3 | 0 | 3.0 | 1 | 2 | .2 | 31 | 9 | 5.0 | 20 | 5 | 61 | 9 | 7.0 | |
| 1995** | 4 | 2 | 3.2 | | | | | | | 8 | 1 | 2.0 | | | 31 | 5 | 6.5 | |

*High water was believed to wash out some redds. **High water after April 18 washed out redds previously marked - good surveys before the washout.

Table G-4. Summary of Spring Chinook Salmon Escapement Survey Data, Umatilla River Basin, 1995.

| Redd # | River Mile | Area Description | Habitat Type | Date Observed | Prespawning Mortalities Sampled | | Spawned-Outs Sampled | |
|--|-------------|--|-------------------|---------------|---------------------------------|---|----------------------|-----------|
| | | | | | F | M | F | M |
| NORTH FORK OF THE UMATILLA RIVER, RM 0.0 TO RM 3.0 | | | | | | | | |
| SURVEYED JULY 11, 31, AUGUST 7, 14, 21, 28, SEPTEMBER 5, 12 | | | | | | | | |
| 21 | 0.1 | Just below above highway highway Bridge Bridge | Riffle Riffle | 9/5 9/5 | 0 | 1 | 8 | 5 (1) |
| 3 | 0.5 | 250 yards below index site | Tailout | 8/14 | | | | |
| 4 | 0.7 | Lower index site | Riffle | 8/28 | | | | |
| 5 | 0.8 | 100 yards above index site | Riffle | 9/5 | | | | |
| 6 | 0.9 | 250 yards below Bear's old start | Riffle | 8/21 | | | | |
| 7 | 0.9 | 200 yards below Bear's old start | Riffle | 8/14 | | | | |
| 8 | 1.0 | Camping area | Riffle | 8/28 | | | | |
| 9 | 1.4 | Mile 1.4 | Riffle | 8/14 | | | | |
| 10 | 1.5 | Mile 1.5 | Riffle | 8/28 | | | | |
| 11 | 1.6 | Mile 1.6 | Riffle | 8/28 | | | | |
| 12 | 2.0 | Mile 2.0 (200 yards above good old area) | Riffle | 8/21 | | | | |
| 13 | 2.8 | Mile 2.8 | Riffle | 8/14 | | | | |
| MAINSTEM UMATILLA RIVER, RM 89.6 TO RM 86.6 (FORKS TO BAR M ENTRANCE) | | | | | | | | |
| SURVEYED JULY 6, 31, AUGUST 8, 14, 24, 28, SEPTEMBER 1, 8, 12 | | | | | | | | |
| 14 | 89.5 | 30 feet below Forks | Riffle | 7/3 | 1 | 5 | 6 (1) | 17 24 (1) |
| 15 | 89.5 | 35 feet below Forks | Riffle | 8/28 | | | | |
| 16 | 89.5 | First habitat structure below Forks | Tailout | 8/21 | | | | |
| 17 | 89.5 | First habitat structure below Forks | Tailout | 8/24 | | | | |
| 18 | 89.5 | First habitat structure below Forks | Tailout | 9/6 | | | | |
| 19 | 89.4 | 100 yards below Forks | Riffle | 8/21 | | | | |
| 20 | 89.3 | Second habitat structure | Tailout | 8/21 | | | | |
| 21 | 89.1 | Just above third habitat structure | I Riffle I | 8/18 | | | | |
| 22 | 89.1 | Just above third habitat structure | Riffle | 8/21 | | | | |
| 23 | 88.3 | Mile 1.2 below Forks | Riffle | 8/21 | | | | |
| 24 | 88.0 | Top of big braid - at beaver diggings | Tailout | 8/28 | | | | |
| II 25 I | 88.0 | Top end of big braid | Tailout I | 8/21 I | | | | |
| 26 | 88.0 | Big braid | Tailout | 9/1 | | | | |
| 27 | 88.0 | Big braid | Riffle | 8/24 | | | | |
| 28 | 87.9 | Big braid | Tailout | 9/1 | | | | |
| 29 | 87.9 | Big braid | Riffle | 9/1 | | | | |
| 30 | 87.9 | Bottom of big braid | Riffle | 8/28 | | | | |
| 31 | 87.7 | River Mile 87.7 | Riffle | 9/8 | | | | |
| 32 | 87.5 | Upper tin shed | Riffle | 8/28 | | | | |
| 33 | 87.5 | Upper tin shed | Riffle | 8/28 | | | | |
| 34 | 86.3 | 125 yards below footbridge at Bar M | Tailout | 9/1 | | | | |
| MAINSTEM UMATILLA, RM 86.6 TO 83.6 | | | | | | | | |
| SURVEYED JULY 6, AUGUST 1, 8, 22, 31, SEPTEMBER 6, 14 | | | | | | | | |
| 35 | 85.9 | Area start riffle | Riffle | 9/6 | I 2 | 2 | 9 | 12 (2) |
| 36 | 85.8 | In beaver workings | Rime | 9/6 | | | | |
| 37 | 85.7 | River Mile 85.7 | Rime | 8/31 | | | | |
| 38 | 84.8 | Stage coach stop | Rime | 8/31 | | | | |
| 39 | 84.8 | Stage coach stop | Riffle | 8/22 | | | | |
| 40 | 84.6 | Lower stage coach stop | Rime | 8/31 | | | | |

Table G-4. Continued

| Redd # | River Mile | Area Description | Habitat Type | Date Observed | Prespawning Mortalities Sampled | | Spawned Out | |
|---|------------|--|--------------|---------------|---------------------------------|-------|-------------|--------|
| | | | | | F | M | F | M |
| 41 | 84.4 | Log truck house | Rifle | 9/6 | | | | |
| 42-43 | 84.4-83.7 | Log A-Frame truck Gulch house | Rime Rifle | 8/31/83 | | | | |
| 44 | 83.7 | A-Frame Gulch | Rifle | 9/6 | | | | |
| MAINSTEM UMATILLA, RM 83.6 TO 80.0 | | | | | | | | |
| SURVEYED JUNE 13, JULY 5, AUGUST 1, 15, 22, 31 SEPTEMBER 6, 14 | | | | | | | | |
| 45 | 82.3 | Homemade fence | Tailout | 9/14 | 1 | 2 | 9 | 12 (2) |
| 46 | 81.9 | Corner above Dabulskis | Rifle | 8/31 | | | | |
| 47 | 81.8 | 150 yards downstream | Rifle | 8/31 | | | | |
| 48 | 81.4 | London bridge | Rifle | 8/31 | | | | |
| 49 | 81.3 | Footbridge | Rime | 9/14 | | | | |
| 50 | 81.0 | Gage | Tailout | 8/31 | | | | |
| 51 | 81.0 | 100 feet below Gage | Rifle | 8/31 | | | | |
| 52 | 80.8 | 100 feet above lower structure at Emmit Williams | Rime | 8/31 | | | | |
| 53 | 80.8 | 100 feet above lower structure at Emmit Williams | Rifle | 9/14 | | | | |
| 54 | 80.7 | Below lower structure at Emmit Williams | Tailout | 9/14 | | | | |
| 55 | 80.5 | River Mile 80.5 | Rifle | 9/14 | | | | |
| 56 | 80.3 | New house above corn cob county | Rifle | 9/14 | | | | |
| 57 | 80.3 | New house above corn cob county | Rifle | 9/14 | | | | |
| MAINSTEM UMATILLA RM 80.0 TO 76.7 (FRED GRAY'S BRIDGE TO SQUAW CREEK) SURVEYED MAY 24, 30, JUNE 13, 21, 27, JULY 5-7, AUGUST 2, 9, 15, 23, 30, SEPTEMBER 7, 13, 18 | | | | | | | | |
| 58 | 79.8 | 300 yards below Fred Gray's bridge | Rifle | 9/13 | 13 | 6 (4) | | |
| 59 | 79.7 | 200 yards above rotary screw trap (RST) at Fred Gray's | Rifle | 9/7 | | | | |
| 60 | 79.7 | 200 yards above RST at Fred Gray's | Rifle | 9/13 | | | | |
| 61 | 79.5 | 125 feet above RST | Rime | 9/7 | | | | |
| 62 | 79.5 | 125 feet above RST | Rime | 9/7 | | | | |
| 63 | 79.5 | 115 feet above RST | Rifle | 9/18 | | | | |
| 64 | 79.4 | 75 yards below RST | Rime | 9/13 | | | | |
| 65 | 79.3 | 200 yards below RST | Tailout | 9/13 | | | | |
| 66 | 79.3 | 225 yards below RST | Tailout | 9/7 | | | | |
| 67 | 79.3 | 230 yards below RST | Rifle | 9/13 | | | | |
| 68 | 79.2 | 250 yards below RST | Tailout | 9/7 | | | | |
| 69 | 79.2 | 275 yards below RST | Rime | 9/18 | | | | |
| 70 | 79.0 | 100 feet above Meacham Creek con. | Rifle | 9/7 | | | | |
| 71 | 78.8 | 250 feet below Meacham Creek con. | Rifle | 9/13 | 11 | 9 (4) | 7 | 3 |
| 72 | 77.5 | 125 feet above Gibbon RR crossing | Rime | 9/13 | | | | |
| 73 | 77.2 | New house | Rifle | 9/7 | | | | |
| 74 | 77.2 | New house | Rifle | 9/7 | | | | |
| 75 | 77.1 | 100 yards below new house | Rifle | 9/7 | | | | |
| 76 | 77.1 | 100 yards below new house | Rifle | 8/30 | | | | |
| 77 | 77.0 | 300 yards below new house | Rime | 9/7 | | | | |

Table G-4. Continued

| Redd # | River Mile | Area Description | Habitat Type | Date Observed | Prespawn Mortality Sampled Type | | Spawned Outs Sampled | |
|---|------------|------------------------------------|--------------|---------------|---------------------------------|-------|----------------------|-------|
| | | | | | F | M | F | M |
| MAINSTEM UMATILLA, RM 76.7 TO RM 73.5 (SQUAW CREEK TO THORNHOLLOW BRIDGE) SURVEYED MAY 30, JUNE 28, AUGUST 2, 9, 18, 23, SEPTEMBER 7, 13, 18 | | | | | | | | |
| 78 | 74.7 | Twin bluffs above Wither's | Rifle | 9/18 | 15 | 3 (1) | 4 | 9 (1) |
| 79 | 74.5 | Above Wither's pool | Rime | 9/13 | | | | |
| 80 | 74.3 | 300 yards below Wither's pool | Rifle | 9/13 | | | | |
| 81 | 73.6 | 200 yards above Thornhollow bridge | Rifle | 9/13 | | | | |
| MAINSTEM UMATILLA, RM 73.5 TO RM 70.0 (THORNHOLLOW BRIDGE TO LOUIE DICK'S FENCE), SURVEYED JUNE 14, 29, JULY 28 | | | | | | | | |
| MAINSTEM UMATILLA, RM 70.0 TO RM 64.5 (LOUIE DICK'S FENCE TO MINTHORN SPRINGS), SURVEYED JUNE 30, JULY 28 | | | | | | | | |
| MAINSTEM UMATILLA, RM 64.5 TO RM 59.5 (MINTHORN SPRINGS TO MISSION BRIDGE), SURVEYED JUNE 7, AUGUST 17 | | | | | | | | |
| MAINSTEM UMATILLA, RM 59.5 TO RM 55.5 (MISSION BRIDGE TO HIGHWAY 11 BRIDGE), SURVEYED JULY 24 | | | | | | | | |
| MAINSTEM UMATILLA, RM 49.0 TO RM 33.8 (REITH BRIDGE TO CUNNINGHAM SHEEP BRIDGE), SURVEYED AUGUST 3, 4 | | | | | | | | |
| MEACHAM CREEK - MOUTH TO RM 3.0, SURVEYED JUNE 1, 15, JULY 7, 26, AUGUST 11, 16, SEPTEMBER 11, 19 | | | | | | | | |
| 82 | 2.0 | Mile 2.0 | Tailout | 9/11 | 0 | 0 | 1 | 1 |
| 83 | 2.9 | Mile 2.9 | Rifle | 9/19 | | | | |
| MEACHAM CREEK - RM 3.0 TO RM 6.0 SURVEYED JUNE 20, JULY 7, 26, AUGUST 10, 16 SEPTEMBER 11, 19 | | | | | | | | |
| 85 | 3.1 | Mile 3.1 | Rifle | 9/19 | 0 | 0 | 1 | 1 |
| 86 | 3.5 | Mile 3.5 | Tailout | 9/11 | | | | |
| 87 | 5.0 | Mile 5.0 | Rifle | 9/11 | | | | |
| 88 | 5.8 | Mile 5.8 | Rifle | 9/11 | | | | |
| MEACHAM CREEK - RM 6.0 TO RM 9.8 SURVEYED JUNE 20, JULY 12, 26*, SEPTEMBER 11*, 19 | | | | | | | | |
| 89 | 6.1 | | Rifle | 9/19 | 0 | 0 | 2 | 0 |
| 90 | 9.8 | 100 feet below Duncan Bridge | Tailout | 9/19 | | | | |
| MEACHAM CREEK - RM 9.8 to RM 13.8 (NORTH FORK), SURVEYED JUNE 26, JULY 12 | | | | | | | | |
| MEACHAM CREEK - RM 13.8 to 19.8, SURVEYED JULY 25 | | | | | | | | |
| NORTH FORK MEACHAM CREEK - RM 0.0 TO 3.0 (BEAR CREEK), SURVEYED JULY 25 | | | | | | | | |

| Total Prespawning Mortalities Sampled | | | Total Spawned Outs Sampled | |
|---------------------------------------|--------|-------------|----------------------------|---------|
| Females | Males | Unknown Sex | Females | Males |
| 41 | 30 (9) | 1 | 73 | 72 (11) |

*Partial survey

() jack salmon which were included in total

Table G-5. Disposition of Umatilla River Spring Chinook Salmon above **Three Mile Falls** Dam, 1989-1995.

| YEAR | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 |
|--|------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Total Observed at TMD | 164 | 2190 | 1330 | 464 | 1221 | 271 | 496 |
| Chinook Sacrificed/Mort. at TMD | 36 | 26 | 234 | 200 | 165 | 31 | 56 |
| Chinook Taken For Brood Stock | 0 | 200 | 0 | 0 | 0 | 0 | 0 |
| Number Released Above TMD | 128 | 1965 | 1096 | 264 | 1056 | 234 | 424 |
| Number Released at TMD | | | | | 9 | 6 | 16 |
| Number of Adipose Clipped Fish Released Above TMD | 3 | 685 | 479 | 135 | 603 | 133 | 156 |
| Estimated Harvest Above TMD | ? | ? | ? | CLOSED | 191 | CLOSED | 0 |
| Number of Chinook Sampled on Spawning Grounds | 6 | 272 | 264 | 79 | 474 | 113 | 217 |
| Percent Recovered (all chinook) | 4.7 | 13.8 | 24.1 | 29.9 | 44.9 | 47.1 | 49.3 |
| Number of Ad. Clipped Chinook Recovered | 0 | 83 | 136 | 39 | 356 | 50 | 78 |
| Percent Recovered (ad. clipped) | 0.0 | 12.1 | 28.4 | 28.9 | 59.0 | 37.6 | 50.0 |
| Prespawning Mortalities Examined | 0 | 0 | 88 | 22 | 125 | 20 | 72 |
| Spawned Out Carcasses Examined | 0 | 0 | 130 | 48 | 338 | 93 | 145 |
| Redds Observed | 14 | 287 | 144 | 59 | 224 | 74 | 90 |
| Spawned Out Females Sampled | -- | -- | 81 | 37 | 205 | 56 | 73 |

Table G-6. Umatilla River Spring Chinook Salmon Redd Distributions, 1989-1995.

| YEAR | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 |
|-------------------------------|--|------------|------------|-----------|------------|-----------|-----------|
| Total # Redds Observed | 14 | 287 | 144 | 59 | 224 | 74 | 90 |
| RIVER SECTION | NUMBER OF REDDS OBSERVED / PERCENT BY REACH | | | | | | |
| North Fork Umatilla River | 0 / 0 | 68 / 23.5 | 13 / 9.0 | 10 / 16.9 | 27 / 12.1 | 16 / 21.6 | 13 / 14.4 |
| River Mile 86 to 89.5 | 14 / 100 | 174 / 60.3 | 21 / 14.6 | 13 / 22.0 | 25 / 11.2 | 13 / 17.6 | 21 / 23.3 |
| River Mile 83 to 86 | | | 29 / 20.1 | 15 / 25.4 | 14 / 6.5 | 6 / 8.1 | 10 / 11.1 |
| River Mile 80 to 83 | 0 / 0 | | 26 / 18.1 | 13 / 22.0 | 31 / 13.8 | 9 / 12.2 | 13 / 14.4 |
| River Mile 78.9 to 80 | 0 / 0 | | 20 / 13.9 | 6 / 10.2 | 39 / 17.4 | 14 / 18.9 | 13 / 14.4 |
| River Mile 76.7 to 78.9 | 0 / 0 | | 36 / 12.5 | | | | |
| River Mile 73.6 to 76.7 | 0 / 0 | | 0 / 0 | 0 / 0 | 25 / 11.1 | 2 / 2.7 | 4 / 4.4 |
| River Mile 70.0 to 73.6 | 0 / 0 | 0 / 0 | 0 / 0 | 0 / 0 | 0 / 0 | 0 / 0 | 0 / 0 |
| River Mile 67.5 to 70.0 | 0 / 0 | 0 / 0 | 0 / 0 | 0 / 0 | 0 / 0 | 0 / 0 | 0 / 0 |
| River Mile 63.8 to 67.5 | 0 / 0 | 0 / 0 | 0 / 0 | 0 / 0 | 0 / 0 | 0 / 0 | 0 / 0 |
| River Mile 59.5 to 63.8 | 0 / 0 | 0 / 0 | 0 / 0 | 0 / 0 | 0 / 0 | 0 / 0 | 0 / 0 |
| Meacham Creek (RM 1-15) | 0 / 0 | 11 / 3.7 | 35 / 24.3 | 1 / 1.7 | 63 / 28.1 | 14 / 18.9 | 9 / 10.0 |

Table G-7. Minimum Estimate of Fall Chinook Salmon and Coho **Salmon** Adult Returns to the Umatilla River, 1989-1994. (Excludes **Jacks**)

| YEAR | ADULTS ENUMERATED AT THREE MILE DAM | ADULTS FOUND BELOW THREE MILE DAM | TOTAL | PERCENT SAMPLED BELOW DAM |
|----------------|-------------------------------------|-----------------------------------|-------|---------------------------|
| COHO | | | | |
| 1989 | 4,154 | 44 | 4,198 | 1.0% |
| 1990 | 409 | 2 | 411 | 0.5% |
| 1991 | 1,732 | 107 | 1,839 | 5.8% |
| 1992 | 356 | 22 | 378 | 5.8% |
| 1993 | 1,531 | 122 | 1,653 | 7.4% |
| 1994 | 984 | 19 | 1,003 | 1.9% |
| CHINOOK | | | | |
| 1989 | 271 | 89 | 360 | 27.2% |
| 1990 | 329 | 110 | 439 | 25.1% |
| 1991 | 522 | 16 | 538 | 3.0% |
| 1992 | 225 | 85 | 310 | 27.4% |
| 1993 | 412 | 70 | 482 | 14.5% |
| 1994 | 688 | 23 | 711 | 3.2% |

Table G-8. Summary of Fall Chinook and Coho Salmon Escapement Data, Umatilla River Basin, 1994.

| REDD # | RIVER MILE | SPECIES | AREA DESCRIPTION | DATE OBSERVED | FISH ON REDDS | | FISH HOLDING | | FISH SAMPLED | | |
|---|------------|---------|--|---------------|---------------|------|--------------|------|--------------|------|---|
| | | | | | FALL CHINOOK | COHO | FALL CHINOOK | COHO | FALL CHINOOK | COHO | |
| FRED GRAY'S BRIDGE TO SQUAW CREEK CONFLUENCE, Surveyed November 21 | | | | | | | | | | | |
| 1 | 78.8 | Chin | 250 yards below Meacham Creek confluence | 11/21 | 5 | 0 | 0 | 0 | 1 | 0 | |
| 2 | 71.4 | Chin | 200 yards below Gibbon RR siding | 11/21 | | | | | | | |
| 3 | 71.4 | Chin | 200 yards below Gibbon RR siding | 11/21 | | | | | | | |
| SQUAW CREEK TO THORNHOLLOW BRIDGE, Surveyed November 21 | | | | | | | | | | | |
| No redds observed in area | | | | | | 0 | 0 | 0 | 0 | 1 | 0 |
| THORNHOLLOW BRIDGE TO LOUIE DICK'S FENCE, Surveyed November 18 | | | | | | | | | | | |
| 4 | 73.1 | Chin | .4 miles below Thornhollow bridge | 11/18 | 19 | 2 | 0 | 0 | 0 | 0 | |
| 5 | 72.8 | Coho | .7 miles below Thomhollow bridge | 11/18 | | | | | | | |
| 6 | 72.7 | Chin | .8 miles below Thornhollow bridge | 11/18 | | | | | | | |
| 1 | 72.7 | Chin | .8 miles below Thomhollow bridge | 11/18 | | | | | | | |
| 8 | 72.7 | Chin | .8 miles below Thomhollow bridge | 11/18 | | | | | | | |
| 9 | 72.7 | Chin | .8 miles below Thomhollow bridge | 11/18 | | | | | | | |
| 10 | 71.7 | Chii | Highway - RR crossing | 11/18 | | | | | | | |
| 11 | 71.7 | Chin | Highway - RR crossing | 11/18 | | | | | | | |
| 12 | 71.7 | Chin | 200 feet below Highway - RR crossing | 11/18 | | | | | | | |
| 13 | 71.3 | Chin | .2 miles below Thomhollow RR bridge | 11/18 | | | | | | | |
| 14 | 71.2 | Chin | .3 miles below Thornhollow RR bridge | 11/18 | | | | | | | |
| 15 | 71.2 | Chin | .3 miles below Thombollow RR bridge | 11/18 | | | | | | | |
| 16 | 71.0 | Chin | Behind Darryl's house | 11/18 | | | | | | | |
| 17 | 70.7 | Chin | 40 yards below lower Thornhollow release site (LTRS) | 11/18 | | | | | | | |
| 18 | 70.6 | Chin | 150 yards below LTRS | 11/18 | | | | | | | |
| 19 | 70.6 | Chin | 150 yards below LTRS | 11/18 | | | | | | | |

Table G-8. Continued

| Redd # | River Mile | Species | Area Description | Date Observed | Fall Chinook on Reds | Coho on Reds | Fall Chinook Holding | Coho Holding | Fall Chinook Sampled | Coho Sampled |
|---|------------|---------|--|---------------|----------------------|--------------|----------------------|--------------|----------------------|--------------|
| 20 | 70.6 | Chin | 150 yards below LTRS | 11118 | | | | | | |
| 21 | 70.3 | Chin | .3 miles above Louis Dick's fence | 11118 | | | | | | |
| 22 | 70.3 | Chin | .3 miles above Louie Dick's fence | 11118 | | | | | | |
| 23 | 70.2 | Chin | .2 miles above Louie Dick's fence above finish cover | 11118 | | | | | | |
| 24 | 70.1 | Chin | 100 yards above Louie Dick's fence | 11/18 | | | | | | |
| LOUIE DICK'S FENCE TO CAYUSE RR BRIDGE, Surveyed November 14, 29 | | | | | | | | | | |
| 25 | 69.5 | Chin | .5 miles below Louie Dick's fence | 11114 | 3 | 2 | 0 | 8 | 4 | 0 |
| 26 | 69.5 | Chin | .5 miles below Louie Dick's fence | 11/14 | | | | | | |
| 27 | 69.3 | Chin | .75 miles below Louie Dick's fence | 11114 | | | | | | |
| 28 | 69.0 | unknw | 1.0 miles below Louie Dick's fence | 11114 | | | | | | |
| 29 | 69.0 | unknw | 1.0 miles below Louie Dick's fence | 11129 | | | | | | |
| 30 | 67.5 | Coho | 50 yards below Cayuse bridge | 11114 | | | | | | |
| 31 | 67.4 | unknw | .1 miles below Cayuse bridge | 11114 | | | | | | |
| 32 | 66.5 | Coho | 1 mile below Cayuse bridge | 11114 | | | | | | |
| CAYUSE RR BRIDGE TO MINTHORN SPRINGS, Surveyed November 14, 29 | | | | | | | | | | |
| 33 | 66.6 | unknw | .4 miles below Cayuse RR bridge | 11129 | 9 | 6 | 1 | 0 | 2 | 2 |
| 34 | 66.6 | Coho | .4 miles below Cayuse RR bridge | 11/29 | | | | | | |
| 35 | 66.6 | Coho | .4 miles below Cayuse RR bridge | 11114 | | | | | | |
| 36 | 66.6 | unknw | .4 miles below Cayuse RR bridge | 11114 | | | | | | |
| 37 | 66.5 | Coho | .5 miles below Cayuse RR bridge | 11/14 | | | | | | |
| 38 | 66.3 | Chin | .7 miles below Cayuse RR bridge | 11114 | | | | | | |
| 39 | 66.3 | unknw | .7 miles below Cayuse RR bridge | 11114 | | | | | | |
| 40 | 65.0 | Chin | 2.0 miles below Cayuse RR bridge | 11114 | | | | | | |
| 41 | 65.0 | Chin | 2.0 miles below Cayuse RR bridge | 11129 | | | | | | |

Table G-8. Continued

| Redd # | River Mile | Species | Area Description | Date Observed | Fall Chinook on Redds | Coho on Redds | Fall Chinook Holding | Coho Holding | Fall Chinook Sampled | Coho Sampled |
|--|------------|-------------|---|---------------|-----------------------|---------------|----------------------|--------------|----------------------|--------------|
| 42 | 64.7 | Coho | Anabmnc above Minthom Springs | 11/29 | | | | | | |
| 43 | 64.7 | Chin | Anabmnc above Minthom Springs | 11129 | | | | | | |
| 44 | 64.7 | Chin | Anabmnc above Minthom Springs | 11114 | | | | | | |
| 45 | 64.6 | Chin | Mainstem -just downstream | 11114 | | | | | | |
| 46 | 64.6 | Chin | Mainstem -jbstw nstream | 11114 | | | | | | |
| 47 | 64.6 | Chin | Mainstem - just downstream | 11/14 | | | | | | |
| MINTHORN SPRINGS TO MISSION HOLES, Surveyed November 16, 28 | | | | | | | | | | |
| 48 | 64.5 | Coho | Minthorn Springs Creek - 50 yards below facility | 11/16 | 36 | 7 | 5 | 0 | 17 | 7 |
| 49 | 64.5 | Coho | Minthom Springs Creek - 125 yards above facility | 11/16 | | | | | | |
| 50 | 64.5 | Coho | Minthom Springs Creek - 50 yards above mouth | 11116 | | | | | | |
| 51 | 64.5 | Chin | Minthom Mainstem | 11/16 | | | | | | |
| 52 | 64.4 | Chin | 100 yards below Minthom Springs Creek | 11116 | | | | | | |
| 53 | 64.4 | Chin | 100 yards below Minthom Springs Creek | 11116 | | | | | | |
| 54 | 64.4 | Chin | 175 yards below Minthom Springs Creek | 11116 | | | | | | |
| 55 | 64.3 | Chin | 250 yards below Minthom Springs Creek | 11128 | | | | | | |
| 56 | 64.3 | Chin | 250 yards below Minthom Springs Creek | 11116 | | | | | | |
| 57 | 64.3 | Chin | 300 yards below Minthom Springs Creek | 11116 | | | | | | |
| 58 | 64.3 | Chin | 300 yards below Minthom Springs Creek | 11116 | | | | | | |
| 59 | 64.3 | Chin | 320 yards below Minthom Springs Creek | 11/28 | | | | | | |
| 60 | 64.3 | Chin | 360 yards below Minthom Springs Creek | 11/16 | | | | | | |
| 61 | 64.3 | Chin | 360 yards below Minthom Springs Creek | 11116 | | | | | | |
| 62 | 64.3 | Chin | 360 yards below Minthom Springs Creek | 11116 | | | | | | |
| 63 | 64.3 | Chin | 360 yards below Minthom Springs Creek | 11116 | | | | | | |
| 64 | 64.3 | Chin | 360 yards below Minthom Springs Creek | 11116 | | | | | | |

Table G-8. Continued

| Redd # | River Mile | Species | Area Description | Date Observed | Fall Chinook on Redds | Coho on Redds | Fall Chinook Holding | Coho Holding | Fall Chinook Sampled | Coho Sampled |
|--|------------|---------|--|---------------|-----------------------|---------------|----------------------|--------------|----------------------|--------------|
| 65 | 64.1 | Chin | 600 yards below Minthorn Springs Creek | 11/16 | | | | | | |
| 66 | 64.0 | Chin | 750 yards below Minthorn Springs Creek | 11/16 | | | | | | |
| 67 | 63.9 | Chin | 1000 yards below Minthorn Springs Creek | 11/28 | | | | | | |
| 68 | 63.9 | Coho | 1000 yards below Minthorn Springs Creek | 11/16 | | | | | | |
| 69 | 60.5 | Chin | 440 yards above Mission swim hole access | 11/28 | | | | | | |
| 70 | 60.3 | Chin | 225 yards above Mission swim hole access | 11/28 | | | | | | |
| 71 | 60.3 | Chin | 200 yards above Mission swim hole access | 11/16 | | | | | | |
| 72 | 60.3 | Chin | 200 yards above Mission swim hole access | 11/16 | | | | | | |
| 73 | 60.3 | Chin | 200 yards above Mission swim hole access | 11/16 | | | | | | |
| 74 | 60.3 | Chin | 200 yards above Mission swim hole access | 11/16 | | | | | | |
| 75 | 60.3 | Chin | 200 yards above Mission swim hole access | 11/16 | | | | | | |
| 76 | 60.3 | Chin | 200 yards above Mission swim hole access | 11/16 | | | | | | |
| 77 | 60.3 | Chin | 200 yards above Mission swim hole access | 11/16 | | | | | | |
| 78 | 60.3 | Chin | 200 yards above Mission swim hole access | 11/16 | | | | | | |
| 79 | 60.3 | Chin | 167 yards above Mission swim hole access | 11/16 | | | | | | |
| 80 | 60.3 | Chin | 167 yards above Mission swim hole access | 11/16 | | | | | | |
| 81 | 60.3 | Chin | 100 yards above Mission swim hole access | 11/16 | | | | | | |
| MISSION HOLES TO BEDROCK CORNER, Surveyed November 17 | | | | | | | | | | |
| 82 | 60.2 | Chin | 50 feet above Mission swim hole access | 11/16 | | | | | | |
| 83 | 60.2 | Chin | Mission swim hole access (SHA) | 11/17 | 11 | 1 | 1 | 0 | 0 | 2 |
| 84 | 60.1 | Coho | 150 yards below SHA | 11/17 | | | | | | |
| 85 | 60.1 | Chin | 155 yards below SHA | 11/17 | | | | | | |
| 86 | 60.1 | Chin | 155 yards below SHA | 11/17 | | | | | | |
| 87 | 59.8 | Chin | .4 miles below SHA | 11/17 | | | | | | |

Table G-8. Continued

| Redd # | River Mile | Species | Area Description | Date Observed | Fall Chinook on Redds | Coho on Redds | Fall Chinook Holding | Coho Holding | Fall Chinook Sampled | Coho Sampled |
|--|------------|---------|-------------------------------------|---------------|-----------------------|---------------|----------------------|--------------|----------------------|--------------|
| 88 | 59.8 | Chin | .4 miles below SHA | 11/17 | | | | | | |
| 89 | 59.8 | Coho | .4 miles below SHA | 11/17 | | | | | | |
| 90 | 59.8 | Chin | .4 miles below SHA | 11/17 | | | | | | |
| 91 | 59.8 | Chin | .4 miles below SHA | 11/17 | | | | | | |
| 92 | 59.8 | Chin | .4 miles below SHA | 11/17 | | | | | | |
| 93 | 59.7 | Chin | .5 miles below SHA | 11/17 | | | | | | |
| 94 | 59.7 | Chin | .5 miles below SHA | 11/17 | | | | | | |
| 95 | 59.7 | Chin | 125 yards above finish | 11/17 | | | | | | |
| 96 | 59.7 | Chin | 115 yards above finish | 11/17 | | | | | | |
| MCKAY DAM TO UMATILLA CONFLUENCE, Surveyed November 1 | | | | | | | | | | |
| 97 | 3.8 | Coho | Above Carl Scheeler's house | 11/1 | 2 | 3 | 0 | 2 | 2 | 6 |
| 98 | 3.8 | Chin | Above Carl Scheeler's house | 11/1 | | | | | | |
| 99 | 2.0 | Coho | McKay Park lower road to confluence | 11/1 | | | | | | |
| 100 | 2.0 | Coho | McKay Park lower road to confluence | 11/1 | | | | | | |
| 101 | 2.0 | Coho | McKay Park lower road to confluence | 11/1 | | | | | | |
| 102 | 2.0 | Coho | McKay Park lower road to confluence | 11/1 | | | | | | |
| 103 | 2.0 | Coho | McKay Park lower road to confluence | 11/1 | | | | | | |
| 104 | 2.0 | Coho | McKay Park lower road to confluence | 11/1 | | | | | | |
| 105 | 2.0 | Coho | McKay Park lower road to confluence | 11/1 | | | | | | |
| BARNHART RELEASE SITE TO 3.7 MILES UPSTREAM, Surveyed November 22, December 7 | | | | | | | | | | |
| 106 | 45.5 | Chin | RM 45.5 | 11/22 | 5 | 0 | 0 | 0 | 12 | 4 |
| 107 | 43.5 | Chin | 200 yards below Bedrock bridge | 11/22 | | | | | | |
| 108 | 42.0 | Chin | .25 miles above Barnhart | 12/7 | | | | | | |

Table G-8. Continued

| Redd # | River Mile | Species | Area Description | Date Observed | Fall Chinook on Redds | Coho on Redds | Fall Chinook Holding | Coho Holding | Fall Chinook Sampled | Coho Sampled | |
|---|------------|---------|--|---------------|-----------------------|---------------|----------------------|--------------|----------------------|--------------|--|
| | | | | | | | | | | | |
| BARNHART RELEASE SITE TO 3.0 MILES DOWNSTREAM, Surveyed November 22, December 8 No redds observed | | | | | | | | | | | |
| 3.0 MILES BELOW BARNHART TO STANFIELD JUVENILE RETURN, Surveyed December 12 | | | | | | | | | | | |
| 109 | 36.6 | Chin | 650 yards below Yokum Bridge | 12/12 | 0 | 0 | 0 | 0 | 4 | 3 | |
| STANFIELD JUVENILE BYPASS RETURN TO ECHO BRIDGE, Surveyed November 23, December 14 | | | | | | | | | | | |
| 110 | 31.6 | Coho | .8 miles below Stanfield Return | 12/14 | 0 | 1 | 0 | 0 | 1 | 7 | |
| 111 | 28.1 | Coho | 200 yards below Cold Springs Diversion | 11/23 | | | | | | | |
| 112 | 27.4 | Coho | 300 yards above Westland | 11/23 | | | | | | | |

Table G-9. Fall Chinook and Coho Salmon Escapement Surveys, 1989-1994

| YEAR | MILES SURVEYED | REDDS | OBSERVED LIVE FISH | | | | RECOVERED CARCASSES | | | |
|----------------------------|----------------|-------|--------------------|------|---------|-------|---------------------|------|---------|-----|
| | | | CHP | COHO | UNKNOWN | TOTAL | CHP | COHO | UNKNOWN | SUM |
| ABOVE THREE MILE FALLS DAM | | | | | | | | | | |
| 1989 | 32.5 | 92 | 5 | 30 | 0 | 35 | 20 | 37 | 10 | 67 |
| 1990 | 42.8 | 50 | 19 | 3 | 11 | 33 | 12 | 6 | 1 | 19 |
| 1991 | 29.0 | 18 | 12 | 15 | 1 | 28 | 5 | 11 | 1 | 17 |
| 1992 | 9.0 | 12 | 0 | 11 | 3 | 14 | 2 | 8 | 1 | 11 |
| 1993 | 42.0 | 44 | 0 | 12 | 0 | 12 | 1 | 14 | 0 | 15 |
| 1994 | 42.3 | 112 | 97 | 33 | 0 | 130 | 49 | 41 | 0 | 90 |
| BELOW THREE MILE FALLS DAM | | | | | | | | | | |
| 1989 | 2.5 | - | 8 | 4 | 15 | 27 | 92 | 52 | 17 | 161 |
| 1990 | 2.5 | - | 15 | 9 | 11 | 35 | 120 | 5 | 8 | 133 |
| 1991 | 2.5 | - | 16 | 68 | 0 | 84 | 16 | 107 | 1 | 124 |
| 1992 | 2.5 | - | 50 | 19 | 0 | 69 | 88 | 22 | 0 | 110 |
| 1993 | 2.5 | - | 6 | 23 | 0 | 29 | 50 | 122 | 0 | 172 |
| 1994 | 2.5 | - | 13 | 13 | 0 | 26 | 25 | 19 | 0 | 44 |

Table G-10. Average Fecundity of Salmonids Returning to the Umatilla River, 1990-1995.

| SPECIES | STOCK | BROOD YEAR | FECUNDITY | MEAN FECUNDITY |
|----------------|-----------------|------------|---------------|----------------|
| Steelhead | Umatilla | 1990 | 5870 | 5669 |
| Steelhead | Umatilla | 1991 | 6412 | |
| Steelhead | Umatilla | 1992 | 5545 | |
| Steelhead | Umatilla | 1993 | 5435 | |
| Steelhead | Umatilla | 1994 | 4884 | |
| Steelhead | Umatilla | 1995 | 5870 | |
| Spring Chinook | Carson | 1991 | 4387 | 4376 |
| Spring Chinook | Carson | 1992 | 3991 | |
| Spring Chinook | Carson | 1993 | 4653 | |
| Spring Chinook | Carson | 1994 | 4328 | |
| Spring Chinook | Carson | 1995 | 4519 | |
| Fall Chinook | Upriver Brights | 1991 | 3783 | 3735 |
| Fall Chinook | Upriver Brights | 1992 | 3373 | |
| Fall Chinook | Upriver Brights | 1993 | 4050 | |
| Coho | Tanner Creek | 1993 | 2356 | 2356 |
| Coho | Tanner Creek | 1995 | not available | |

Table G- 11. Spring Chinook Salmon Escapement Data Umatilla River, 1095.

| MEHP | FL | SCALES | SEX | AREA CODE | AREA SAMPLED | HATCHERY/BROOD TAG CODE | MARKS | SPAWNING STATUS | DATE | REMARKS |
|------|------|--------|-----|-----------|---|-------------------------|-------|-----------------|----------|---|
| 820 | | yes | M | 01 | NF - old good area log jam | | | partial | 8/21/95 | Dead 2 days - upper caudal punch |
| 770 | | yes | M | 03 | NF Index Area | | | partial | 8/28/95 | Dead 1 week+ |
| 780 | 855 | yes | M | 04 | Corporation | | | partial | 08/14/95 | Dead 1 day |
| 455 | | no | M | 12 | 40 feet above Fred Gray's Bridge | | LV | Partial | 9/18/95 | |
| 645 | 810 | yes | M | 12 | RST to Meacham Con. | | RV | Partial | 9/13/95 | |
| 670 | 860 | yes | M | 15 | 1.3 miles below Squaw Creek | | RV | Partial | 9/18/95 | Dead 2 days |
| 625 | 800 | yes | M | 15 | 1.0 miles below Squaw Creek | | RV | Partial | 9/18/95 | Dead 1 day |
| 615 | 820 | yes | M | 16 | Wither's Pool | | RV | Partial | 9/13/95 | |
| 480 | | no | M | 09 | Corner above Dubalski's | | LV | Partial | 9/14/95 | |
| 480 | 565 | no | M | 12 | RST to Meacham Con. | | LV | Partial | 9/13/95 | |
| 405 | | no | J | 09 | Dubalski's Dam | | LV | Partial | 9/14/95 | |
| 840 | 1080 | yes | M | 12 | Fred Gray's Trap | | | Partial | 9/11/95 | Dead 1 day |
| 440 | 550 | no | J | 05 | Big Braid | | LV | Partial | 9/11/95 | Dead 1 day |
| 590 | | yes | F | 17 | 60 feet below Thornhollow Bridge | | RV | PM | 7/25/95 | Dead 1 + week |
| 635 | 760 | no | F | 11 | Lower Emmitt Williams | | RV | PM | 9/14/95 | Poached-Ripid open |
| 660 | 620 | yts | F | 06 | .1miles above Bar M | | RV | PM | 7/31/95 | Bad gills- dead 2 days |
| 635 | 760 | yes | F | 13 | 75 yards below Meacham Con | | RV | PM | 8/23/95 | Dead 2 days- good gills |
| 675 | 650 | no | M | 11 | 200 yards above Emmitt Williams | | RV | PM | 9/14/95 | Dead 1 day |
| 655 | 615 | no | F | 12 | 100 yards below Fred Gray's Outlet | | RV | PM | 0/1 8/95 | Bad Gills |
| 650 | | yes | F | 14 | Gibbon RR Siding | | RV? | PM | 8/30/95 | Dead 4 days-bad gills - poor RV |
| 220 | | no | M | 13 | just below Meacham Con | | RV | PM | 7/05/95 | Dead one day |
| 680 | | yes | M | 10 | London Bridge | | | PM | 8/01/95 | Dead 1 + weeks |
| 640 | 790 | yes | M | 14 | Gibbon RR Siding | | RV | PM | 7/27/95 | Sad gills- dead 3+ days |
| 715 | 915 | no | M | 14 | 150 yards below New House | | RV | PM | 9/18/95 | |
| 605 | 740 | yes | F | 13 | 250 yards below Meacham Con. | | RV | PM | 8/15/95 | Dead 4 days |
| 585 | | no | F | 15 | .75 miles below Squaw Creek | | RV | PM | 8/18/95 | Dead 1 week+ lost scale envelope at Wither's |
| 680 | | yes | F | 16 | 150 yards above Thornhollow Bridge | | RV | PM | 8/14/95 | Heavy fungus on head- many deep head cuts from jumping |
| 615 | 740 | yes | F | 15 | 1.0 miles below Squaw Creek | | ?? | PM | 9/13/95 | |
| 440 | 550 | no | M | 12 | RST to Meacham Con. | | LV | PM | 9/13/95 | Dead 1 day |
| 500 | 835 | no | M | 13 | Mtacham Con to Squaw Creek | | LV | PM | 9/13/95 | |
| 395 | 490 | no | M | 12 | 250 yards below RST | | LV | PM | 9/13/95 | |
| 465 | 560 | no | J | 13 | Old Meacham Con. | | LV | PM | 9/7/95 | Died today- bad gills |
| 475 | 610 | no | M | 12 | 250 yards below RST | | LV | PM | 9/13/95 | |
| 510 | 620 | yes | M | 15 | Below split channel merge - below Squaw | | LV | PM | 9/7/95 | Very old mort - Radio 7-23 |
| 415 | 520 | no | J | 13 | Gibbon RR Siding | | LV | PM | 8/12/95 | Dead several days- no scales |
| 620 | 760 | yes | F | 14 | 150 yards below new house | | RV | PM | 8/23/95 | Died today- fungused gills- green color on skeins and liver |
| 655 | | yes | F | 14 | Gibbon RR siding | | RV | PM | 8/27/95 | Dead one week+ couldn't tell cause |
| 940 | | yes | M | 12 | First corner below RG Bridge | | NM? | PM | 7/27/95 | Possible poor RV clip-dead 1 week+ |
| 460 | 570 | no | J | 13 | Gibbon RR Siding | | LV | PM | 9/12/95 | Dead several days- no scales |
| 705 | 690 | yts | M | 14 | Gibbon RR Siding | | RV | PM | 9/7/95 | Dead 2 days- old shaker injury |
| 660 | 615 | yes | F | 15 | Wither's | | RV | PM | 9/7/95 | |
| 660 | 615 | yes | F | 13 | Gibbon RR Siding | | RV | PM | 8/23/95 | Dead 2-3 days- gills good-fungus patches on sltd(2) |
| ? | | no | M | 13 | just below Mtacham Con. | | ? | PM | 7/05/95 | poached mort??only gut track present |
| 480 | | no | J | 05 | Big Braid | | LV | PM | 9/8/95 | Dead 5 days |
| 790 | 1010 | yes | M | 03 | .5 miles above NF Mouth | | | PM | 8/07/95 | Dead 1 day - dorsal+ ventral fungus-radio tagged 13-35 |
| 605 | 965 | yes | F | 04 | 100 yards below Forks-Umatilla | | | PM | 7/06/95 | A few jump marks on head- 5 days old |
| 745 | 910 | yes | F | 12 | RST to Mtacham Con. | | | PM | 9/13/95 | |
| 690 | 625 | yes | F | 04 | 50 yards below NF | | | PM | 8/28/95 | |
| 600 | 960 | yes | M | 05 | Big Braid | | | PM | 9/1/95 | Dead 1 week+ |
| 675 | 640 | yes | M | 12 | RST | | | PM | 9/18/95 | W/OUT FISH?? |
| 670 | 610 | yes | F | 15 | Below split channel merge-below Squaw | | | PM | 9/7/95 | |
| 640 | | yes | F | 07 | 200 yards below Bar M | | | PM | 7/28/95 | Habitat survey |
| 710 | 695 | yts | F | 07 | .2 milts below Bar M | | | PM | 8/08/95 | Dead 3 days |
| 600 | | yes | M | 08 | Upper Bar M Horse Crossing | | | PM | 8/08/95 | Dead 1 week+ + + |
| 745 | 925 | yes | M | 07 | .7 miles below Bar M | | | PM | 8/15/95 | |
| 765 | 976 | yes | M | 05 | .5 miles below Umatilla National Forest | | | PM | 8/02/95 | Dead 1 week+Habitat Survey |
| 705 | | yes | M | 05 | Braided area below Forks | | | PM | 8/24/95 | Very old mort |
| 666 | 765 | yes | F | 15 | .8 miles below Squaw Crtk | | | PM | 8/23/95 | Dead 3 days |

G-20

Table G-1 1. Continued

| MEHP | FL | SCALES | SEX | AREA CODE | AREA SAMPLE | HATCHERY/BROOD TAG CODE | MARKS | STATUS | DATE | REMARKS |
|-------|------|--------|-----|-----------|--|-------------------------|-------|--------|----------|--|
| Adult | | no | F | 18 | Wither's Swim Hole | | | PM | 8/23/95 | Trail of eggs m wing up bank- animal |
| 660 | | yes | F | 14 | 75 feet above Squaw Creek Con. | | | PM | 8/13/95 | nose about gone, dead 2 days- gill fungus-marks on head behind eye |
| 665 | 795 | yes | F | 25 | Meacham - AM 3.0 | | RV | R10 | 9/21/95 | |
| 765 | | yes | F | 04 | 300 yards below Forks | | | R10 | 8/07/95 | Dead 2 days |
| 680 | 780 | yes | F | 05 | Mile 1.2 below Forks | | | R100 | 9/1/95 | Dead 2 days- Large growth on right side |
| 610 | | no | F | 11 | RM 80.3 | | RV | R100 | 9/14/95 | |
| 670 | | yes | F | 15 | .5 miles below Squaw Creek | | RV | R12 | 9/13/95 | |
| 650 | | yes | F | 12 | Fred Gray's Rotary Trap | | | R20 | 9/28/95 | |
| 760 | 890 | yes | F | 04 | .2 miles below Forks | | | R20 | 8/21/95 | |
| 855 | | no | F | 12 | 275 yards below Rotary Trap | | RV | R20 | 9/18/95 | |
| 610 | | yes | F | 16 | Thornhollow Bridge | | RV | R30 | 9/22/95 | Radio Tagged |
| 620 | | no | F | 12 | RST Fred Gray's | | RV | R50 | 9/13/95 | Dead 1 day |
| 610 | 1030 | yes | M | 05 | 115 yards below Big Braid | | | SO | 9/1/95 | Dead 2 days |
| 785 | 1000 | yes | M | 04 | 200 yards below Forks | | | SO | 8/24/95 | good gills- died today |
| 855 | 1090 | yes | M | 04 | .2 miles below Forks | | | SO | 8/24/95 | |
| 855 | 1070 | yes | M | 04 | Corporation | | | SO | 08/14/95 | Dead 3 days |
| 745 | | yes | F | 03 | NF-250 yards below Bear's start | | | SO | 8/28/95 | Dead 1 day |
| 670 | 850 | yes | M | 05 | Just upstream of Larson's Driveway | | | SO | 9/8/95 | Dead 1 week+ |
| 455 | 580 | yes | J | 08 | 1.5 miles below Bar M | | LV | SO | 9/14/95 | |
| 480 | 565 | yes | J | 08 | 1.9 miles below Bar M | | LV | so | 9/14/95 | |
| 810 | 1015 | ye * | M | 04 | 200 yards below Forks | | | so | 8/24/95 | good gills- died three days ago |
| 440 | 550 | yes | M | 16 | 2.1 miles below Squaw Creek | | LV | SO | 9/13/95 | |
| 795 | | yes | M | 03 | NF- .4 miles above mouth | | | SO | 8/21/95 | dead 4 days |
| 685 | 840 | yes | F | 05 | 2.0 miles below NF | | | SO | 9/12/95 | just below Big Braid |
| 805 | 870 | yes | r | 04 | 300 yards below NF | | | so | 8/28/95 | |
| 490 | 570 | no | M | 11 | Lower Emmitt Williams | | LV | SO | 9/14/95 | |
| 435 | | no | J | 03 | NF-250 yards below Bear's start | | LV | so | 8/28/95 | Near SO female- dead several days |
| 710 | | yes | F | 04 | 25 yards below 2nd habitat structure below Forks | | | SO | 9/1/95 | Dead 1 day |
| 710 | | no | F | 12 | RST to Meacham Con. | | | SO | 9/13/95 | |
| 475 | 575 | | F | 05 | Big Braid | | LV | SO | 9/8/95 | Dead 2 days |
| 780 | | yes | F | 01 | .2 miles below Coyote Creek | | | SO | 8/28/95 | Dead 1 day |
| 605 | 745 | yes | F | 07 | Below Bar M | | | | 9/14/95 | |
| 770 | | yes | F | 02 | Mile 1.6 below Coyote Creek | | | SO | 9/5/95 | Sacrificed- last day of life |
| | | no | M | 17 | 100 yards below Thornhollow Bridge | | | SO | 9/28/95 | no scale envelope |
| 800 | 1020 | yes | M | 05 | Big Braid | | | SO | 9/1/95 | Sacrificed |
| 840 | | yes | F | 03 | 500 yards below Bear's start | | RV | so | 8/28/95 | Dead 1 day |
| 670 | | yes | F | 04 | Below first habitat structure below Forks | | RV | so | 9/5/95 | Dead 2 days |
| 690 | 890 | no | M | 12 | 50 yards below Outlet Fred Gray's | | RV | SO | 9/18/95 | |
| 695 | 890 | no | M | 12 | Outlet Fred Gray's | | RV | so | 9/18/95 | |
| 670 | 845 | yes | M | 12 | Rotary Trap | | RV | SO | 9/20/95 | |
| 640 | 800 | no | M | 12 | 50 yards below RST | | RV | so | 9/18/95 | Shaker |
| 690 | 870 | yes | M | 24 | Meacham Creek-RR Bridge below Bon | | RV | SO | 9/19/95 | |
| 630 | 770 | no | M | 05 | Mile 1.7BF | | RV | so | 9/1/95 | Dead 3 days |
| 650 | 800 | no | F | 14 | New House | | RV | SO | 9/18/95 | |
| 675 | 780 | yes | F | 15 | Wither's | | RV | SO | 9/18/95 | |
| 500 | 555 | yes | M | 04 | .2 miles below Forks | | RV | so | 8/24/95 | |
| 615 | 745 | yes | F | 15 | 1.5 miles below Squaw Creek | | RV | so | 9/13/95 | |
| 610 | | no | F | 29 | Meacham Creek- mile 6.1 | | RV | SO | 9/19/95 | Dead 5 days |
| 665 | 810 | yes | F | 05 | 2.0 miles below NF | | RV | SO | 9/12/95 | just below Big Braid |
| 810 | 1060 | yes | M | 05 | Big Braid | | | | 9/8/95 | Dead 1 seek+ |
| 745 | 940 | yes | M | 05 | Big Braid | | | | 9/8/95 | Dead 2 days |
| 685 | 820 | yes | F | 04 | 400 yards below NF | | | SO | 8/28/95 | |
| 775 | 970 | yes | M | 01 | NF- good old area | | | SO | 8/28/95 | Dead 2 days |
| 810 | 1030 | yes | M | 04 | Corner below 3rd habitat structure below Forks | | | SO | 9/1/95 | Dead 2 days |
| 685 | 720 | yes | M | 07 | .1 miles below Bar M | | | so | 8/31/95 | |
| 705 | 900 | yes | M | 04 | 200 yards below Forks | | RV | so | 9/1/95 | Dead 2 days- Tail punch 1 in |
| 790 | | yes* | M | 08 | Clark's Bridge | | | so | 9/8/95 | Dead 5 days |
| 615 | 785 | yes | M | 08 | 1.5 miles below Bar M | | RV | so | 9/14/95 | |

| MEHP | FL | SCALES | SEX | AREA | | HATCHERY/BROOD TAG CODE | MARKS | SPAWNING | | DATE | REMARKS |
|------|------|--------|-----|------|--|-------------------------|--------|----------|---------|----------|-------------------------|
| | | | | CODE | AREA SAMPLED | | | STATUS | DATE | | |
| 650 | 775 | yes | F | 11 | Corn Cob County | | RV | SO | | 9/20/95 | |
| 600 | | yes | F | 03 | 200 yards below Bear's start | | RV | SO | | 9/5/95 | Dead 3 days |
| 625 | 790 | no | M | 05 | Tin Shed - mile 2.0 BF | | RV | SO | | 9/8/95 | Dead 1 week - no scales |
| 750 | | no | F | 13 | 50 yards below old Meacham Con. | | RV | SO | | 9/18/95 | |
| 615 | 725 | yes | F | 13 | Gibbon Store | | RV | SO | | 9/27/95 | |
| 605 | 715 | yes | F | 32 | Meacham Creek - Duncan Bridge | | RV | SO | | 9/27/95 | |
| 645 | | no | F | 10 | 40 yards below Footbridge | | RV | SO | | 9/14/95 | |
| 680 | | no | F | 10 | London Bridge | | RV | SO | | 9/14/95 | |
| 665 | | yes | F | 11 | Emmitt Williams | | RV | SO | | 9/20/95 | |
| 645 | | no | F | 14 | 200 yards above Squaw Creek Con. | | RV | SO | | 9/13/95 | |
| 625 | 770 | no | F | 12 | 100 yards above RST | | RV | SO | | 9/18/95 | |
| 605 | 770 | no | M | 05 | Mile 1.9 BF | | RV | SO | | 9/8/95 | Died today - no scales |
| 660 | 830 | yes | M | 15 | Split channel merge below Squaw Creek | | RV | SO | | 9/7/95 | |
| 660 | 840 | yes | M | 05 | 100 yards below Big Braid | | RV | SO | | 9/8/95 | Dead 5 days |
| 605 | 750 | yes | M | 16 | Thornhollow Bridge | | RV | SO | | 10/02/95 | |
| 670 | 810 | yes | F | 12 | 200 yards above Rotary Trap - FG | | RV | SO | | 9/7/95 | Dead 1 day - bad gills |
| 700 | | yes | F | 14 | 225 yards below new house | | RV | SO | | 9/7/95 | Dead 3 days |
| 615 | | yes | F | 12 | RST | | RV | SO | | 9/18/95 | |
| 665 | | yes | F | 05 | 30 yards below Big Braid | | RV | SO | | 9/8/95 | Dead 3 days |
| 640 | | no | F | 10 | 200 yards below Footbridge | | RV | SO? | | 9/14/95 | |
| 670 | | no | M | 12 | 300 yards below Fred Gray's Outlet | | | SO? | | 9/22/95 | old mort |
| 680 | | no | F | 11 | 200 yards below Lower Emmitt Williams | | RV | SO? | | 9/14/95 | |
| 405 | | no | J | 10 | Larson's to Fred Gray's Bridge | | ?? | ?? | | 9/14/95 | Eaten By Crayfish |
| 885 | 855 | yes | M | 15 | 1.3 miles below Squaw Creek | BON-91 | 071455 | 95J2241 | PM | | 9/13/95 |
| 690 | 840 | yes | F | 12 | 300 yards below Fred Gray's Outlet | BON-91 | 071455 | 95J2292 | PM | | 9/18/95 |
| 640 | 795 | yes | F | 18 | Thornhollow Bridge | BON-91 | 071455 | 95J2214 | PM | | 9/22/95 |
| 610 | 740 | yes | F | 07 | .8 miles below Bar M | BON-91 | 071455 | 95J2239 | R20 | | 8/31/95 |
| 670 | | yes | F | 12 | 100 feet above Rotary Screw Trap - FG | BON-91 | 071455 | 95J2278 | R60 | | 9/8/95 |
| 705 | 835 | yes | F | 08 | 1.5 miles below Bar M | BON-91 | 071455 | 95J2248 | SO | | 9/14/95 |
| 600 | | no | M | 05 | Mile 1.7 BF | BON-91 | 071455 | 95J2265 | SO | | 9/8/95 |
| 570 | 720 | no | M | 07 | .1 miles below Bar M | BON-91 | 071455 | 95J2245 | SO | | 9/14/95 |
| 680 | 795 | no | F | 11 | Lower Emmitt Williams | BON-91 | 071455 | 95J2234 | SO | | 9/14/95 |
| 680 | 825 | yes | F | 05 | Big Braid | BON-91 | 071455 | 95J2251 | SO | | 9/1/95 |
| 645 | 830 | yes | M | 05 | Mile 1.8 BF | BON-91 | 071455 | 95J2266 | SO | | 9/8/95 |
| 680 | 760 | no | F | 08 | Below Bar M | BON-91 | 071455 | 95J2222 | ?? | | 9/14/95 |
| 615 | 790 | no | M | 13 | 100 yards above Gibbon RR Siding | BON-91 | 071455 | 95J2294 | ?? | | 9/18/95 |
| 625 | | no | M | 11 | RM 80.3 | BON-91 | 071456 | 95J2237 | Partial | | 9/14/95 |
| 770 | 845 | yes | F | 12 | 200 Yards below Fred Gray's release site | BON-91 | 071456 | 95J2201 | PM | | 5/30/95 |
| 635 | 800 | yes | F | 13 | Old Meacham Con. | BON-91 | 071456 | 95J2202 | PM | | 6/13/95 |
| 625 | 790 | yes | M | 12 | First corner below RG Bridge | BON-91 | 071456 | 95J2207 | PM | | 7/27/95 |
| 690 | 830 | yes | F | 08 | Behind Bar M | BON-91 | 071456 | 95J2221 | PM | | 8/22/95 |
| 680 | | yes | F | 12 | Fred Gray's Trap | BON-91 | 071456 | 95J2220 | SO | | 9/11/95 |
| 670 | 850 | yes | M | 14 | Gibbon RR Siding | BON-91 | 071456 | 95J2295 | SO | | 9/18/95 |
| 715 | 920 | yes | M | 09 | 1.0 miles below Bar M | BON-91 | 071456 | 95J2247 | SO | | 9/14/95 |
| 625 | | yes | F | 05 | Mile 1.2 BF | BON-91 | 071456 | 95J2284 | SO | | 9/8/95 |
| 740 | 900 | yes | F | 12 | 250 yards above Meacham Creek Confluence | BON-99-MEACHAM | 075440 | 95J2218 | PM | | 8/09/95 |
| 615 | 730 | yes | F | 12 | 100 yards above Meacham Creek Con. | UM-91 | 075740 | 95J2206 | PM | | 7/05/95 |
| 675 | 815 | yes | F | 15 | 1.0 miles below Squaw Creek | UM-91 | 075741 | 95J2224 | PM | | 8/02/95 |
| 650 | 795 | yes | M | 08 | Bar M Barn | UM-91 | 075741 | 95J2217 | PM | | 8/08/95 |
| 610 | 750 | yes | F | 15 | 1.4 miles below Squaw Creek | UM-91 | 075741 | 95J2231 | R 3000 | | 9/18/95 |
| 630 | 780 | yes | M | 15 | Beaver Farm | UM-91 | 075741 | 95J2216 | SO | | 9/20/95 |
| 580 | | no | F | 28 | Meacham Creek - mile 5.8 | UM-91 | 075741 | 95J2296 | SO | | 9/19/95 |
| 630 | | yes | F | 14 | Meacham Con. to Squaw Creek | UM-91 | 075741 | 95J2230 | SO | | 9/13/95 |
| 645 | | no | F | 01 | 200 yards below old good spawning area | UM-91 | 075742 | 95J2255 | SO | | 9/5/95 |
| 640 | 770 | yes | F | 08 | 1.7 miles below Bar M | UM-91 | 075742 | 95J2205 | SO | | 9/14/95 |
| 600 | 725 | yes | F | 07 | Just below Bar M Driveway | UM-91 | 075742 | 95J2242 | SO | | 9/14/95 |
| 835 | 1040 | yes | M | 04 | 500 yards below NF | BON-90-MEACHAM | 075828 | 95J2238 | SO | | 8/28/95 |
| 820 | 1040 | yes | M | 12 | RST to Meacham Con | BON-90-MEACHAM | 075830 | 95J2273 | Partial | | 9/13/95 |

APPENDIX H
Emigrant Trapping Tables and Figures

Table H-1. Summary of Trap Catch Data from the Bar&art, **Tumla** and **Imeques** Traps sites, **1994/95**; Expanded Migration Estimates Include Days the Traps were not Operated within the Trapping Dates.

| | TRAPS | | |
|--|-------------------------|-------------------------|-------------------------|
| | BARNHART (RM 42.2) | TUMLA (RM 76) | IMEQUES (79.5) |
| Trapping Dates | 03/05/95 to 06/01/95 | 09/22/94 to 01/13/95 | 05/05/95 to 06/16/95 |
| Trapping days over total days | 87 / 125 | 63 / 113 | 43 / 43 |
| Natural Chinook | | | |
| Number Captured | 247 | 1,368 | 102 |
| Number Marked and Released | 112 | 1,207 | 95 |
| Total Number Recaptured | 5 | 348 | 10 |
| Average % Recaptured | 4.5% | 28.9% | 10.5% |
| Expanded Migration Estimate | 14,542 | 11,035 | 1093 |
| Mean Fork Length (mm) | 94.2 | 93.8 | 70.9 |
| Number Measured | 134 | 1363 | 100 |
| Sample Standard Deviation | 18.3 | 8.2 | 9.8 |
| Average % Containment | 87% | 72% | 85% |
| Number of containment trials | 4 | 12 | 5 |
| Natural Rainbow/Steelhead | | | |
| Number Captured | 105 | 596 | 304 |
| Number Marked and Released | 52 | 516 | 273 |
| Total Number Recaptured | 3 | 47 | 18 |
| Average % Recaptured | 5.7% | 9.9% | 6.6% |
| Expanded Migration Estimate | 4,789 | 14,029 | 7,435 |
| Mean Fork Length (mm) | 165 | 115.5 | 106 |
| Number Measured | 64 | 596 | 301 |
| Sample Standard Deviation | 33.2 | 35.2 | 27.4 |
| Average % Containment | 100% | 44% | 78% |
| Number of containment trials | 2 | 13 | 4 |
| CONTINUED ON THE FOLLOWING PAGE | | | |

| TABLE H-1 CONTINUED | TRAPS | | |
|----------------------------------|-----------------------|------------------|----------------------|
| | BARNHART (RM 42.2) | TUMLA (RM 76) | IMEQUES (RM 79.5) |
| Natural Coho Captured | 5 | 0 | 0 |
| Mean Fork Length (mm) | 111 | 94 | |
| Range (mm) | 66-139 | 92-95 | |
| Hatchery Chinook Captured | 6,265 | 41 | 289 |
| Marked and Released | 684 | 0 | 263 |
| Recaptured | 18 | 0 | 44 |
| Average % Recaptured | 2.6% | | 16.7% |
| Expanded Migration Estimate | 626,876 | | 1,728 |
| Mean Fork Length (mm) | 140 | 142 | 128 |
| Number Measured | 445 | 107 | 5 |
| Standard Deviation or Range | 26.8 | 29 | 83-240 (mm) |
| Hatchery STS Captured | 467 | 0 | 0 |
| Marked and Released | 258 | 0 | 0 |
| Recaptured | 6 | | |
| Average % Recaptured | 2.3% | | |
| Expanded Migration Estimate | 52,844 | | |
| Mean Fork Length (mm) | 213 | | |
| Number Measured | 267 | | |
| Sample Standard Deviation | 20.1 | | |
| Hatchery Coho Captured | 16,844 | 0 | 0 |
| Marked and Released | 3047 | 0 | 0 |
| Recaptured | 226 | | |
| Average % Recaptured | 7.4% | | |
| Expanded Migration Estimate | 599,000 | | |
| Mean Fork Length (mm) | 138 | | |
| Number Measured | 638 | | |
| Sample Standard Deviation | 10.7 | | |
| Bull Trout | 0 | 15 | 4 |
| Mean Fork Length (mm) | | 281.7 | 158.8 |
| Range (mm) | | 220-395 | 147-175 |
| Whitefish | 0 | 36 | 0 |
| Redside Shiner | 296 | 1,065 | 151 |
| Sucker | 63 | 71 | 154 |
| Dace | 262 | 1,289 | 2,653 |
| Sculpin | 12 | 694 | 63 |
| Squawfish | 30 | 84 | 26 |
| Chiselmouth | 52 | 8 | 39 |
| Yellow Perch | 1 | 0 | 0 |
| Brown Bullhead | 2 | 0 | 0 |

Table H-2. Estimated number of adult natural steelhead that would have been produced in the absence of the supplementation project (TMD = Three Mile Falls Dam, * = assuming same survival rates as cohorts, ** = portion of run contributed by each brood year estimated from scale samples).

| Return Year | Brood Year | Total Natural Steelhead Enumerated at TMD | | | | Natural Brood Taken | | | | Natural Steelhead Less Brood Taken | | Total Natural Steelhead Return | | | |
|--|------------|---|------|------|------|---------------------|------|------|------|------------------------------------|------|--------------------------------|------|------|------|
| | | 1981 | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | | 1991 | 1992 | 1993 |
| 80/81 | 1298 | 768 | 1264 | 2314 | 3197 | 2885 | 3444 | 2316 | 2104 | 1422 | 724 | 2247 | 1297 | 946 | 875 |
| 81/82 | 1264 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 82/83 | 1264 | 404 | 1385 | 482 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 83/84 | 2314 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 84/85 | 3197 | 1535 | 1215 | 1096 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 85/86 | 2885 | 404 | 1385 | 482 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 86/87 | 3444 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 87/88 | 2316 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 88/89 | 2104 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 89/90 | 1422 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 90/91 | 724 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 91/92 | 2247 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 92/93 | 1297 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 93/94 | 946 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 94/95 | 875 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 80/81 | 1298 | 768 | 1264 | 2314 | 3197 | 2885 | 3444 | 2316 | 2104 | 1422 | 724 | 2247 | 1297 | 946 | 875 |
| 81/82 | 1264 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 82/83 | 1264 | 404 | 1385 | 482 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 83/84 | 2314 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 84/85 | 3197 | 1535 | 1215 | 1096 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 85/86 | 2885 | 404 | 1385 | 482 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 86/87 | 3444 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 87/88 | 2316 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 88/89 | 2104 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 89/90 | 1422 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 90/91 | 724 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 91/92 | 2247 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 92/93 | 1297 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 93/94 | 946 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| 94/95 | 875 | 879 | 1535 | 1215 | 1096 | 1096 | 324 | 1309 | 1112 | 880 | 1010 | 295 | 1010 | 199 | 880 |
| Total Adult Progeny (Natural Production) | | 2818 | 3082 | 3073 | 2716 | 2089 | 1584 | 1201 | 1536 | 1609 | 1070 | | | | |
| Adult to Adult Percent Survival | | 220 | 507 | 279 | 120 | 68 | 56 | 36 | 70 | 83 | 81 | | | | |
| Additional Adults Produced Had Brood Not Been Taken for Supplementation (Through 1994/1995 Return) * | | 176 | 811 | 449 | 62 | 71 | 39 | 53 | 93 | 133 | 87 | | | | |
| Number of Hatchery Steelhead Collected at Three Mile Falls Dam (Through 1994/95 Return) | | | | | | | | | | | | | | | |
| Total, 3306 | | | | | | | | | | | | | | | |
| Total, 2844 (Estimated Natural Adult Steelhead Lost Because of Broodstock Mining) | | 32 | 87 | 43 | 59 | 133 | 59 | 133 | 87 | 32 | | | | | |
| Mean, 152% (Adult to Adult Survival) | | 81 | 1070 | 1070 | 1070 | 1070 | 1070 | 1070 | 1070 | 1070 | 1070 | 1070 | 1070 | 1070 | 1070 |




Figure H-1. Length Frequencies of Juvenile Natural Chinook Salmon Captured by the Rotary Screw Traps in the Umatilla River; Tumla Trap (RM 76, n=1363) from September 22, 1994 to January 13, 1995; Imeques Trap (RM 79.5, n= 100) from May 5, 1995 to June 16, 1995, and Barnhart Trap (RM 42.2, n= 134) from March 5, 1995 to June 1, 1995 (TPCN945L.CH3).




Figure H-2. Length Frequency of Juvenile Hatchery Chinook Salmon Captured by the Rotary Screw Traps in the Umatilla River; Tumla Trap (RM 76, n= 107) from September 22, 1994 to January 13, 1995, and Barnhart Trap (RM 42.2, n=445) from March 5, 1995 to June 1, 1995 (TPCH945L.CH3).




Figure H-3. Length Frequencies of Juvenile Hatchery Coho Salmon Captured by the Rotary Screw Traps in the Umatilla River, **Barnhart Trap (RM 42.2, n=638)** from March 5, 1995 to June 1, 1995 (TPHH945L.CH3).




Figure H-4. Length Frequencies of Juvenile Natural and Hatchery Summer Steelhead Captured by the Rotary Screw Traps in the Umatilla River; **Tumla Trap (RM 76, n=596)** from September 22, 1994 to January 13, 1995; **Imeques Trap (RM 79.5, n=301)** from May 5, 1995 to June 16, 1995, and **Barnhart Trap (RM 42.2, Natural n=64, Hatchery n=267)** from March 5, 1995 to June 1, 1995 (TPSN945L.CH3).




Figure H-5. Bamhart Trap (RM 42.2) from March 5, 1995 to June 1, 1995, Total Salmonid Catch, River Discharge (1000 CFS), Days When Most or All of the Catch Escaped, Days Trap Operated, Days When Trap was Checked but Catch was Held Over to the Next Day (TB945TFC.CH3).




Figure H-6. Bamhart Trap (RM 42.2) from March 5, 1995 to June 1, 1995, Total Salmonid Catch and Water Temperatures (C), (TB945TC2.CH3).




Figure H-7. Imeques Trap (RM 79.5) from May 5, 1995 to June 16, 1995, Total Salmonid-Catch, River Discharge (1000 CFS), Days Trap Operated, Days When Trap was Checked but Catch was Held Over to the Next Day (TI945TFC.CH3).




Figure H-8. Imeques Trap (RM 79.5) from May 5, 1995 to June 16, 1995, Estimated Number of Salmonids Migrating Past Trap (CHS = spring chinook; STS = summer steelhead; CH = hatchery spring and/or fall chinook), (TI945EC2.CH3).




Figure H-9. Tumla Trap (RM 76) from September 22, 1994 to January 13, 1995, Total Salmonid Catch, River Discharge (100 CFS), Water Temperature (C), Days When Most or All of the Catch Escaped, Days Trap Operated, Days When Trap was Checked but Catch was Held Over to the Next Day (TT945TFC.CH3).




Figure H-10. Tumla Trap (RM 76) from September 22, 1994 to January 13, 1995, Estimated Number of Salmonids Migrating Past Trap (CH = natural chinook; STS = natural summer steelhead; TT945TF2.CH3).

APPENDIX I

Age and Growth Tables

Table I-1. Age Summary by Sex of the Umatilla River Wild Summer Steelhead Escapement in the Umatilla River, 1995.

| AGE | | 1.1 | 1.2 | 2.1 | 2.2 | 3.1 | 3.2 | Total |
|--------|-----|-----|-----|------|------|------|------|-------|
| FEMALE | n = | 0 | 0 | 11 | 9 | 6 | 7 | 33 |
| | % = | 0 | 0 | 33.3 | 27.2 | 18.2 | 21.2 | 100 |
| MALE | n = | 0 | 0 | 8 | 8 | 3 | 4 | 23 |
| | % = | 0 | 0 | 34.8 | 34.8 | 13.0 | 17.4 | 100 |
| TOTAL | n = | 0 | 0 | 19 | 17 | 9 | 11 | 56 |
| | % = | 0 | 0 | 33.9 | 30.4 | 16.1 | 19.6 | 100 |

Table I-2. Brood Year of the 1995 Umatilla River Wild Summer Steelhead Escapement.

| BROOD YEAR | | 1991 | 1990 | 1989 | Total |
|------------|-----|------|------|------|-------|
| FEMALE | n = | 11 | 15 | 7 | 33 |
| | % = | 33.3 | 45.5 | 21.2 | 100 |
| MALE | n = | 8 | 11 | 4 | 23 |
| | % = | 34.8 | 47.8 | 17.4 | 100 |
| TOTAL | n = | 19 | 26 | 11 | 56 |
| | % = | 33.9 | 46.4 | 19.6 | 100 |

Table I-3. Freshwater Age Data of the 1995 Wild Summer Steelhead Escapement in the Umatilla River.

| AGE | | 1 | 2 | 3 | Total |
|--------|-----|---|------|------|-------|
| FEMALE | n = | 0 | 20 | 13 | 33 |
| | % = | 0 | 60.6 | 39.4 | 100 |
| MALE | n = | 0 | 16 | 7 | 23 |
| | % = | 0 | 69.6 | 30.4 | 100 |
| TOTAL | n = | 0 | 36 | 20 | 56 |
| | % = | 0 | 64.3 | 35.7 | 100 |

Table I-4. Ages Based on Scale Analysis and Expansions Based on Comparisons of Age Versus Fork Length of Juvenile **Rainbow/Steelhead** Sampled in Various Tributaries of the Umatilla River, 1995.

UMATILLA RIVER, AUGUST 8 -25, 1995

| Age | n = | Range(mm) | Mean(mm) | S.D. | L/F-Age Expansion | Percent |
|-----|-----|-----------|----------|------|-------------------|---------|
| 0+ | 76 | 36-95 | 63.6 | 14.0 | 1291 | 68.0 |
| 1+ | 82 | 92-182 | 123.7 | 22.4 | 509 | 26.8 |
| 2+ | 30 | 132-258 | 186.9 | 26.8 | 93 | 4.9 |
| 3+ | 3 | 190-240 | 215.7 | 20.4 | 5 | .3 |

MISSION CREEK, SEPTEMBER 5-13, 1995

| Age | n = | Range(mm) | Mean(mm) | S.D. | L/F-Age Expansion | Percent |
|-----|-----|-----------|----------|------|-------------------|---------|
| 0+ | 25 | 56-111 | 85.1 | 13.8 | 116 | 57.4 |
| 1+ | 25 | 89-242 | 178.8 | 38.0 | 63 | 31.2 |
| 2+ | 13 | 160-290 | 224.2 | 34.8 | 23 | 11.4 |

COTTONWOOD CREEK, JULY-6 AUGUST 1,1995

| Age | n = | Range(mm) | Mean(mm) | S.D. | L/F-Age Expansion | Percent |
|-----|-----|-----------|----------|------|-------------------|---------|
| 0+ | 12 | 51-100 | 70.5 | 13.5 | 87 | 50.9 |
| 1+ | 18 | 100-188 | 143.3 | 21.1 | 63 | 36.8 |
| 2+ | 9 | 140-222 | 181.2 | 22.8 | 20 | 11.7 |
| 3+ | 1 | 216 | | | 1 | .6 |

MOONSHINE CREEK, SEPTEMBER 18-21, 1995

| Age | n = | Range(mm) | Mean(mm) | S.D. | L/F-Age Expansion | Percent |
|-----|-----|-----------|----------|------|-------------------|---------|
| 0+ | 36 | 48-120 | 86.7 | 14.8 | 258 | 69.9 |
| 1+ | 33 | 118-194 | 158.3 | 21.1 | 97 | 26.3 |
| 2+ | 6 | 212-240 | 226.2 | 8.5 | 14 | 3.8 |

MOONSHINE CREEK, SEPTEMBER 18-21, 1995

| Age | n = | Range(mm) | Mean(mm) | S.D. | L/F-Age Expansion | Percent |
|-----|-----|-----------|----------|------|-------------------|---------|
| 0+ | 11 | 42-65 | 55.1 | 7.1 | 83 | 26.8 |
| 1+ | 56 | 83-182 | 120.9 | 23.1 | 195 | 62.9 |
| 2+ | 11 | 118-243 | 175.5 | 35.7 | 31 | 10.0 |
| 3+ | 0 | | | | | |
| 4+ | 1 | 327 | | | 1 | .3 |

Table I-5. Bull Trout Biological Data, 1994-1995.

| FORK LENGTH | AGE | SEX | AREA CAPTURED | DATE | REMARKS |
|--------------------|------------|------------|---------------------------------|-------------|---------------------------|
| 165 | 2+ | I | RM 79.5-Rotary Screw Trap-(RST) | 05/16/95 | Live |
| 170 | 2+ | | RM 88.4-Biological Survey | 08/23/95 | Live |
| 220 | 2+ | | RM 89.2-Biological Survey | 08/25/95 | Live |
| 222 | 2+ | | RM 79.5 (RST) | 09/27/95 | Live |
| 233 | 2+ | | RM 89.2-Biological Survey | 08/25/95 | Live |
| 245 | 2+ | | RM 79.5 (RST) | 11/02/95 | Live |
| 254 | 2+ | | RM 79.5 (RST) | 09/23/95 | Live |
| 258 | 2+ | | RM 79.5 (RST) | 11/13/95 | Live |
| 268 | 2+ | | RM 79.5 (RST) | 11/10/95 | Live |
| 270 | 2+ | Male | RM 2.0-North Fork Umatilla | 08/15/94 | Hooking Mortality-Spawner |
| 225 | 3+ | | RM 88 A-Biological Survey | 08/25/95 | Live |
| 265 | 3+ | | RM 87.7-Biological Survey | 08/22/95 | Live |
| 285 | 3+ | | RM 79.5 (RST) | 11/10/95 | Live |
| 288 | 3+ | | RM 79.5 (RST) | 10/05/95 | Live |
| 290 | 3+ | | RM 79.5 (RST) | 10/23/95 | Live |
| 320 | 3+ | | RM 79.5 (RST) | 10/23/95 | Live |
| 390 | 4+ | Female | RM 79.5- 25 feet above RST | 06/01/94 | Lure in throat |

APPENDIX J

Table J-1. Summary of Landmarks and their Associated River Miles, Umatilla River Basin.

| Location / Landmark | RM | Location / Landmark | RM |
|-------------------------------|-----------|----------------------------------|-----------|
| Three Mile Falls Dam | 3.7 | Gibbon Railroad Yard | 78.4 |
| Horse Ranch | 5 | Mouth Of Meacham Creek | 79.0 |
| Tree Farm | 5.5 | Imeques C-mem-i&kern | 79.5 |
| House on Bluff | 7.4 | Fred Gray's Bridge | 80.0 |
| South Park Bridge | 8.8 | Emmit Williams Place | 81.1 |
| Boyd's Return | 9 | London Bridge | 81.4 |
| Boyd's Dam | 10.2 | Reservation Boundary--Ryan Creek | 81.8 |
| Lookinglass Road | 11.3 | Larson's Driveway | 83.1 |
| Maxwell Dam | 15.2 | Stage Coach Stop House | 84.8 |
| Simplot | 17 | Bar M Driveway | 85.9 |
| Stanfield Bridge | 23 | Bear Creek | 86.8 |
| I-84 Bridge | 24.2 | Old Silver Building | 87.1 |
| Dillon Dam | 24.6 | Corporation Hole | 88.5 |
| Echo Bridge | 26.3 | Umatilla Mainstem Forks | 89.5 |
| Westland Dam | 27.2 | North Fork Umatilla River | 0-10 |
| Coldsprings Dam | 28.2 | Coyote Creek | 2.5 |
| Stanfield Dam | 32.4 | Woodward Creek | 5.7 |
| Yoakum | 37 | South Fork Umatilla River | 0-10 |
| Barnhart Bridge | 42.2 | Buck Creek | 0.5 |
| Forth's Diversion | 46.9 | Thomas Creek | 3.3 |
| Mouth of Birch Creek | 48.3 | Shimmiehom Creek | 4.6 |
| PGG Building | 51 | Meacham Creek | 0-36 |
| ODFW, Receiver Site #4 | 56 | Boston Canyon Creek | 2.2 |
| Pendleton Ready Mix | 57 | Bonifer Acclimation Site | 2.3 |
| Mission Bridge | 59.5 | Line Creek | 5.0 |
| Minthom Springs | 64.5 | Camp Creek | 10.9 |
| Cayuse Railroad Bridge | 67.0 | Duncan | 12.0 |
| Cayuse Highway Bridge | 67.5 | North Fork Meacham Creek | 15.0 |
| Louie Dick's Fence | 70.0 | East Meacham Creek | 18.5 |
| Thomhollow Railroad Bridge | 71.0 | Butcher Creek | 21.5 |
| Badger Comer | 71.8 | Meacham | 30.0 |
| Thomhollow Highway Bridge | 73.5 | North Fork Meacham Creek | 0-9.5 |
| Weathers's Place | 74.5 | Bear Creek | 3.0 |
| Mouth of Squaw Creek | 76.7 | Pot Creek | 5.2 |

Table J-2. Abbreviations Used in this Paper.

| | |
|--------------|---|
| BOR | US Bureau of Reclamation |
| BPA | Bonneville Power Administration |
| CTUIR | Confederated Tribes of the Umatilla Indian Reservation |
| CWT | Coded Wire Tags |
| DEQ | Department of Environmental Quality |
| MEHP | Mid-eye to Hypural Plate |
| ODFW | Oregon Department of Fish and Wildlife |
| RM | River Mile |
| TMD | Three Mile Dam |
| UBNPME | Umatilla Basin Natural Production Monitoring and Evaluation Project |
| UMEOC | Umatilla Monitoring Evaluation and Oversight Committee |
| USFS | US Forest Service |
| USGS | US Geological Survey |