

LOWER SNAKE RIVER COMPENSATION PLAN:
Oregon Summer Steelhead Evaluation Studies
2015 Annual Progress Report

Oregon Department of Fish and Wildlife
Fish Research and Development, NE Region



Lance R. Clarke
Holly M. Stanton
Michael W. Flesher
Debra L. Eddy
Richard W. Carmichael



This program receives federal financial assistance from the U.S. Fish and Wildlife Service and prohibits discrimination on the basis of race, color, national origin, age, sex, or disability. If you believe that you have been discriminated against as described above in any program, activity, or facility, or if you desire further information, please contact ADA coordinator, Oregon Department of Fish and Wildlife, Oregon Department of Fish and Wildlife 4034 Fairview Industrial Drive SE, Salem, OR 97302, 503-947-6000, or contact the Office for Human Resources, U.S. Fish and Wildlife Service, Region 1, 911 NE 11th Avenue, Portland, OR 97232-4181 or call (503)231-6136. This report is available at: <http://www.fws.gov/lsnakecomplan/Reports/ODFWreports.html>

Front cover photo: Big Canyon Facility on the Willowa River, August 2011. Photo by Chad Aschenbrenner.

ANNUAL PROGRESS REPORT

FISH RESEARCH PROJECT
OREGON

PROJECT TITLE: Lower Snake River Compensation Plan: Oregon Summer Steelhead
Evaluation Studies

CONTRACT NUMBER: F14AC00042 and F16AC00030

PROJECT PERIOD: January 1, 2015 through December 31, 2015

Prepared By: Lance R. Clarke
Holly M. Stanton
Michael W. Flesher
Debra L. Eddy
Richard W. Carmichael

September 2017

Oregon Department of Fish and Wildlife
4034 Fairview Industrial Drive SE
Salem, OR 97302

This project was financed by the U.S. Fish and Wildlife Service under
the Lower Snake River Compensation Plan.

PREFACE

The purpose of this progress report is to provide summary information for Lower Snake River Compensation Plan (LSRCP) summer steelhead (*Oncorhynchus mykiss*) programs operated by ODFW in the Grande Ronde and Imnaha river basins during 2015. These ongoing monitoring programs provide technical, logistical, and biological information to managers charged with maintaining viable salmon and steelhead populations and associated fisheries in northeast Oregon. This report is organized into fish culture monitoring for juveniles, adults, CWT recoveries, and estimates for total escapement. During the period covered in this report, steelhead from the 2010-2012 broods returned to spawn, and steelhead from the 2014 brood were released as smolts. Adult steelhead that returned to spawn were used to create the 2015 brood.

ACKNOWLEDGMENTS

We would like to thank hatchery managers Ron Harrod and Marc Garst, as well as Terry Blessing, Chad Aschenbrenner and many other hatchery personnel who exhibited great dedication and provided essential assistance. Numerous personnel from the Oregon Department of Fish and Wildlife, U.S. Fish and Wildlife Service, the Nez Perce Tribe, and the Confederated Tribes of the Umatilla Indian Reservation provided enthusiastic support. We also thank Chris Starr, Joe Krakker, Steve Yundt, Julie Collins, and Rod Engle, who provided administrative and technical support. This project was funded by the U.S. Fish and Wildlife Service under the Lower Snake River Compensation Plan, contract numbers F14AC00042 and F16AC00030, a cooperative agreement with the Oregon Department of Fish and Wildlife.

CORRECTION

Appendix A has corrections to Tables 7 and 8 of the 2014 Annual Progress Report.

CONTENTS

PREFACE	i
ACKNOWLEDGMENTS	ii
CORRECTION	ii
CONTENTS.....	iii
LIST OF FIGURES	iv
LIST OF TABLES	v
EXECUTIVE SUMMARY	1
Objectives	1
Accomplishments and Findings.....	1
INTRODUCTION	2
RESULTS AND DISCUSSION	3
Juveniles.....	3
Adults.....	4
Experimental Group Returns	5
Compensation Area Goals	6
REFERENCES	28

LIST OF FIGURES

<u>Figure</u>		<u>Page</u>
1.	Length-at-age relationships based on scale analysis for hatchery adult returns of one freshwater age (top) and two freshwater age (middle) Wallowa stock summer steelhead from 1991 to 2014, and in 2015 (bottom)	8
2.	Length-at-age relationships based on scale analysis for hatchery adult returns of one freshwater age (top), and two and three freshwater age (middle) Imnaha stock summer steelhead from 1991 to 2014, and in 2015 (bottom)	9
3.	Smolt-to-adult survival (SAS) for Wallowa and Imnaha stock summer steelhead, 1985-2010 brood years	10
4.	Smolt-to-adult return (SAR) to the compensation area above Lower Granite Dam for Wallowa and Imnaha stock summer steelhead, 1985-2010 brood years	10
5.	Progeny-to-parent ratios for Little Sheep Creek summer steelhead, 1987-2010 brood years	11
6.	Returns of naturally produced summer steelhead to Little Sheep Creek, run years 1984-85 to 2014-15	11
7.	Hatchery returns to the compensation area above Lower Granite Dam for Wallowa and Imnaha Stock summer steelhead.....	12

LIST OF TABLES

<u>Table</u>	<u>Page</u>
1. Summary of egg collection and juvenile survival for 2014 brood year summer steelhead released in the Grande Ronde and Imnaha river basins at LSRCP facilities in 2015	13
2. Estimates of fin clip quality and coded-wire tag retention for 2014 brood year summer steelhead reared at Irrigon Fish Hatchery and released in 2015.....	14
3. Details of experimental and production groups of 2014 brood year summer steelhead released in the Grande Ronde (Wallowa stock) and Imnaha (Imnaha stock) river basins in 2015.....	15
4. Density ($\pm 95\%$ confidence interval) and mean fork length (standard deviation in parentheses) of residual hatchery steelhead, wild rainbow trout/juvenile steelhead and Juvenile Chinook Salmon from index sites on Deer (Grande Ronde basin) and Little Sheep (Imnaha basin) creeks in 2015.....	16
5. Travel days to Lower Granite Dam (LGD) and percent of the PIT tagged releases that were uniquely detected at the four lower Snake River dams for Little Sheep Creek smolts that departed the acclimation pond during the early, middle, and late periods of the volitional release.....	16
6. Number of PIT tags released and unique adult PIT tag detections at Bonneville and Lower Granite dams during the 2014-15 run year by stock and brood year	17
7. Timing of adult steelhead returns to LSRCP facilities in 2015 by location and origin	18
8. Number, disposition, and mean fork length (mm) of adult steelhead that returned to LSRCP facilities in 2015 by stock, origin, estimated age (freshwater:saltwater), and gender	19
9. Number of adult summer steelhead trapped at the Little Sheep Creek Facility weir that were either outplanted to Big Sheep Creek or passed above the weir, and were subsequently recaptured, 1999-2015	21
10. Summary of summer steelhead spawning surveys in Deer Creek above the Big Canyon Facility weir, 2002-2015	22
11. Spawning summaries for summer steelhead at LSRCP facilities in 2015.	23

LIST OF TABLES (continued)

<u>Table</u>	<u>Page</u>
12. Summary of anadromous adult recoveries of coded-wire tagged (CWT) Willowa stock summer steelhead for the 2014-15 run year	24
13. Summary of anadromous adult recoveries of coded-wire tagged (CWT) Imnaha stock summer steelhead for the 2014-15 run year	25
14. Harvest and escapement distribution of adult summer steelhead by recovery location for the 2014-15 run year using the PSMFC and ODFW mark recovery databases	26
15. Harvest and escapement distribution of adult summer steelhead by age and recovery location for the 2014-15 run year using the PSMFC and ODFW mark recovery databases	27

EXECUTIVE SUMMARY

Objectives

1. Document summer steelhead rearing and release activities at all LSRCP facilities.
2. Determine optimum rearing and release strategies that will produce maximum survival to adulthood for hatchery-produced summer steelhead smolts.
3. Document summer steelhead adult returns by stock to each LSRCP broodstock collection facility.
4. Determine if the total production of summer steelhead adults meets mitigation goals, and index annual smolt survival and adult returns to Lower Granite Dam for production groups.
5. Participate in planning activities associated with anadromous fish production and management in the Grande Ronde and Imnaha river basins, and participate in ESA permitting, consultation, and rearing activities.
6. Monitor natural spawning of summer steelhead in selected areas within the Grande Ronde basin.
7. Determine the number of summer steelhead harvested annually and angler effort in recreational fisheries on the Grande Ronde, Wallowa, and Imnaha rivers.

Accomplishments and Findings

We accomplished each of our objectives for 2015. In this report, we present data and results for objectives 1, 2, 3, 4, and 6. To accomplish objective 5, project staff participated in planning and coordination with co-managers to develop and write the annual operation plan (available at: www.fws.gov/lsnakecomplan/Reports/AOPreports.html). Data and results for objective 7 are published in separate annual creel survey reports (e.g., *Flesher et al. 2015*).

The production goal of 800,000 Wallowa stock smolts was achieved in 2015, with 860,995 smolts released. The Imnaha stock production goal of 215,000 smolts was also achieved with 247,642 smolts released.

In 2015, 2,210 and 1,105 Wallowa stock hatchery steelhead returned to Wallowa Fish Hatchery and the Big Canyon Facility, respectively. We trapped 17 natural steelhead at Wallowa Fish Hatchery and 58 natural steelhead at the Big Canyon Facility, which were released to spawn naturally. At the Little Sheep Creek Facility, we trapped 1,080 Imnaha stock hatchery and 93 natural steelhead adults. Of these, a total of 71 hatchery and 57 natural steelhead were released above the weir, and 355 hatchery steelhead were outplanted to Big Sheep Creek. During spawning in the spring of 2015, we collected 643,900 Wallowa stock production eggs, 630,200 Wallowa fall broodstock eggs, and 333,600 Imnaha stock eggs.

In the 2014-15 run year, the compensation area goal of 9,184 Wallowa stock adults above Lower Granite Dam was met and the Imnaha stock goal of 2,000 adults was also met. We have met the Wallowa stock compensation area goal twelve times in our program history, and the Imnaha stock compensation area goal fourteen times. We estimate that 9,300 Wallowa stock hatchery steelhead (101.2% of goal), and 4,017 Imnaha stock hatchery steelhead (200.9% of goal) returned to the LSRCP compensation area in 2015.

INTRODUCTION

The objectives of this report are to document fish culture practices, describe adult returns, and assess progress toward meeting LSRCP goals for Grande Ronde and Imnaha steelhead (*Oncorhynchus mykiss*). We report on juvenile steelhead rearing and release activities for the 2014 brood year (BY) released in 2015. Included are collection, spawning, and adult characteristics for the 2015 returns, returns from experimental releases, supplementation in Little Sheep Creek, and success toward achieving compensation goals.

The Grande Ronde and Imnaha river steelhead hatchery programs were initiated in 1976 and 1982 in response to the rapid decline in Snake River steelhead abundance. Annual adult mitigation, brood year specific smolt-to-adult return and total smolt-to-adult survival rates, and annual smolt production goals were established to compensate for the estimated annual loss of 48% of adult production. Adaptive management has resulted in current interim smolt production goals of 800,000 (ODFW Wallowa stock released into the Grande Ronde) and 215,000 (Imnaha stock) smolts; less than the original goals of 1,350,000 and 330,000 smolts. Based on original smolt production goals it was assumed that 27,552 Wallowa stock and 6,000 Imnaha stock adults would be produced annually. Furthermore, 66.7% of these fish were expected to be harvested below the compensation area, defined as the watershed above Lower Granite Dam, resulting in compensation area adult return goals of 9,184 Wallowa stock and 2,000 Imnaha stock.

In general, the data in this report were derived from hatchery inventories and standard databases (e.g., Pacific States Marine Fisheries Commission Regional Mark Information System (RMIS), ODFW mark recovery) or through standard measuring techniques. As such, specific protocols are usually not described. In cases where expansions of data or unique methodologies were used, protocols are described in more detail. Additional descriptions of protocols can be found in our work statements (Carmichael et al. 2012, Carmichael et al. 2013). Coded-wire tag (CWT) data collected from 2015 adult returns were used to evaluate smolt-to-adult survival rates in experimental rearing and release groups. In 2015, the only experimental treatments from which fish returned were second generation progeny from early returning (fall-collected) broodstock. In 2015, smolts were released at Wallowa Hatchery that were third generation progeny of early returning (fall-collected) broodstock for an experimental comparison with progeny of standard production broodstock. Methods for the fall broodstock experiment are described in Warren et al. (2011a). Analysis of specific survival studies will be completed and published in separate reports once all brood years have returned and CWT data are complete for each experiment. In addition, much of the data that we discuss in this report will be used in separate and specific evaluations of ongoing supplementation programs for steelhead in the Imnaha River basin. We began culture evaluations in 1983 and have dramatically improved

many practices. Progress for work completed in previous years is presented in annual progress reports (Carmichael and Wagner 1983; Carmichael and Messmer 1985; Carmichael et al. 1986a; 1987; 1988a; 1999; 2004; 2005a; 2005b; Clarke et al. 2014; Clarke et al. 2015; Flesher et al. 2005a; 2009a; Gee et al. 2007; 2008; Messmer et al. 1989; 1990; 1991; 1992; 1993; Jonasson et al. 1994; 1995; 1996; Ruzycki et al. 2003; Warren et al. 2009; 2010; 2011a; 2011b; 2012; 2013; Whitesel et al. 1993), annual creel survey reports (Carmichael et al. 1988b; 1989; 1990; Flesher et al. 1991; 1992; 1993; 1994; 1995; 1996; 1997; 1999; 2000; 2001; 2004a; 2004b; 2005b; 2007; 2008a; 2008b; 2009b; 2010; 2011; 2012; 2013; 2014), a United States vs. Oregon production report (Carmichael et al. 1986b), a five-year study plan (Carmichael 1989), and journal articles (Clarke et al. 2010; 2011; 2014).

RESULTS AND DISCUSSION

Juveniles

Wallowa stock egg-to-eyed embryo survival for the 2014 BY was 91.1%, within the range of recent brood years (1993-2014 BY range = 71.8-93.8%), and embryo-to-smolt survival was 81.6%, within the range of recent brood years (1993-2014 BY range = 65.0-98.3%; Table 1). Imnaha stock egg-to-embryo survival for the 2014 BY was 91.3%, within the range of recent brood years (1993-2014 BY range = 76.7-92.1%), and embryo-to-smolt survival was 93.4%, within the range of recent brood years (1993-2014 BY range = 79.5-98.5%; Table 1). We released 860,995 Wallowa stock smolts in 2015, exceeding our production goal of 800,000 smolts. For the Imnaha stock, we released 247,642 Imnaha stock smolts, which also exceeded our production goal of 215,000 smolts (Tables 1 and 3). Hatchery managers attempt to meet production goals every year; however, variation in mortality at various stages of rearing, from fertilized eggs to acclimated smolts, results in fewer or more fish being released in any given year. Managers periodically adjust the number of eggs collected based on recent hatchery performance.

Beginning with BY 2013 releases, a programmatic decision was made to eliminate ventral fin clipping of steelhead for purposes of identifying the presence of coded-wire tags. Electronic scanning is now used to detect wire in hatchery fish harvested in fisheries and recovered at hatchery traps. However, one raceway of coded-wire tagged Wallowa production stock continue to be left ventral fin clipped (AdLV and CWT) to assess the effect that ventral clips have on smolt-to-adult survival, and Wallowa fall broodstock continue to be right ventral clipped so that returning adults may be visually identified at hatchery weirs and collected for broodstock. Hatchery fish continue to be adipose fin clipped. To evaluate different rearing and release strategies, we tagged and released six groups of Wallowa stock steelhead and one group of Imnaha stock steelhead smolts with adipose clips and coded-wire-tags (Ad and CWT), and four groups of Wallowa fall broodstock with adipose-right ventral clips and coded-wire tags (AdRV and CWT; Table 2). We marked 99.6 and 100% of Wallowa and Imnaha stock smolts with an adipose fin clip, which was within the range of recent brood years for Wallowa stock (1993-2013 BY range = 95.6-99.9%) and within the range of recent brood years for Imnaha stock (1993-2013 BY range = 96.1-100.0). Fin clip quality and tag retention for release groups averaged 98.7% for Wallowa stock, within the range of recent years (1993-2013 BY range = 89.1-99.3%)

and 99.6% for Imnaha stock, which slightly exceeded the range of recent years (1993-2013 BY range = 84.7-99.5%). Details of experimental and production releases for the 2014 BY, including the number of fish implanted with passive integrated transponder (PIT) tags, are shown in Table 3.

Densities of residual hatchery steelhead averaged 3.8 fish/100m² at index sites in the Grande Ronde basin in 2015 (Table 4), whereas wild *O. mykiss* averaged 6.5 fish/100m². In the Imnaha basin, densities of residual hatchery steelhead and wild *O. mykiss* were 16.1 and 1.8 fish/100m². Since sampling for residual hatchery steelhead began in 1996, we have observed a clear pattern of higher densities of residual hatchery steelhead than wild *O. mykiss* in the Imnaha basin index sites.

Adults

Returning PIT-tagged adults from the 2009 to 2011 broods were detected at mainstem dams during the 2014-15 run year. Of the 479 Wallowa stock adults detected at Bonneville Dam on the Columbia River, 350 were detected at Lower Granite Dam on the Snake River. For the Imnaha stock, 369 of the 530 adults detected at Bonneville Dam were detected at Lower Granite Dam (Table 6). Weirs were installed to capture adult steelhead on 27 January at Wallowa Fish Hatchery, 17 February at Big Canyon Facility, and 20 February at Little Sheep Creek Facility (Table 7). Returns to the Little Sheep Creek Facility were predominantly hatchery fish, with 93 (7.9%) natural steelhead. Similar to Little Sheep Creek, most of the adults that returned to the Big Canyon Facility were of hatchery origin, with only 58 (5.0%) natural steelhead. In addition, 15 (0.7%) natural steelhead returned to Wallowa Fish Hatchery. Fifty six percent of hatchery adults that returned to Wallowa Fish Hatchery and Big Canyon Facility spent one year in the ocean (Table 8) and 71.5% of hatchery fish that returned to Little Sheep Creek Facility spent one year in the ocean before returning. Of the natural origin fish, 62% (58 of 93), 64% (37 of 58), and 47% (8 of 17) of the Little Sheep Creek Facility, Big Canyon Facility, and Wallowa Fish Hatchery, respectively, spent one year in saltwater before returning.

The majority of hatchery adults that returned to Wallowa Fish Hatchery in 2015 were spawned or killed (Table 8). In 2015, Big Canyon Facility hatchery returns were not needed for the Grande Ronde steelhead hatchery program due to the large number of adults returning to Wallowa Fish Hatchery. We outplanted 247 adult hatchery steelhead from Wallowa Fish Hatchery and the Big Canyon Facility to local ponds for harvest opportunities. However, fish captured at Big Canyon Facility are no longer returned to the Wallowa River for further angling opportunities. At the Big Canyon Facility, 58 natural fish were passed above the weir to spawn naturally. We retained 8.8% of the hatchery fish and 38.7% of the natural fish for spawning at Little Sheep Creek Facility, and outplanted hatchery adults to Big Sheep Creek to spawn naturally. Ten of the 355 outplanted fish (2.8%) were recaptured at least once at the Little Sheep Creek Facility in 2015. Fifty-seven natural and 71 hatchery adults were released above the weir in Little Sheep Creek to spawn naturally. In addition, 19 natural males were spawned and then passed above the weir, resulting in 52% of fish above the weir being of hatchery origin. Of the 147 fish passed into Little Sheep Creek, 1 fell back and was recaptured at the weir (Table 9). Length-at-age data for Wallowa and Imnaha stock adults are presented in Figures 1 and 2, respectively.

We conducted multiple spawning surveys for steelhead that were passed above the Big Canyon Facility weir into Deer Creek using protocols described in Gee et al. (2008). In 2015, forty-nine redds were counted, which was 144% of the total number of redds constructed, assuming that each female constructs one redd.

In 2015, we reached our egg take goal for the Wallowa stock with 1,274,100 green eggs collected. Of these, 643,900 were for production and 630,200 were for the fall broodstock evaluation. We collected 333,600 green Imnaha stock eggs, thus we exceeded our goal of 313,850 eggs. Mortality from green egg-to-eyed embryo from six weekly spawns ranged from 3.1-6.8% for Wallowa production stock, 0.0-14.0% for fall broodstock, and from 3.2-14.9% for Imnaha stock (Table 11).

Experimental Group Returns

The number of coded-wire tag (CWT) and adipose-clipped adults that were harvested or returned to collection sites is used to estimate various performance parameters. These numbers allow us to monitor our success toward meeting the LSRCP goals, to estimate stray rates, and to determine the contribution to recreational, tribal, and commercial fisheries. They also provide the basis for evaluating the success of experimental rearing and release strategies. Recoveries for each CWT code were summarized from the CWT recovery database maintained by PSMFC, ODFW's mark recovery database, and from data reported by the Washington Department of Fish and Wildlife and Idaho Department of Fish and Game. Our protocol was to collect and enumerate all fish marked with a CWT when they were spawned, dispatched, or died. A summary of these data is provided in this report. Final analyses, results, and discussion of production and release strategies will be presented in special reports once all adults have returned from the experimental groups.

Adults from BY 2010 to 2012 returned during the 2014-15 run year, including the ninth year of adult returns from the Wallowa fall broodstock experiment. Of approximately 200,000 total coded-wire-tagged fish released for both production and fall brood groups, a total of 992 Wallowa fall brood and 2,015 Wallowa production CWTs were recovered (Table 12). We had Wallowa stock recoveries from 21 CWT codes (Table 12) and Imnaha stock recoveries from two CWT codes (Table 13).

Adult return data from the first generation of the fallbrood experiment became complete in fiscal year 2014, with results showing that average smolt-to-adult survival (SAS) rates were similar between the two groups (fallbrood = 1.84%, production = 1.80%) but the fallbrood strayed at a higher rate (7.64%) than the standard production groups (5.01%). However, due to their earlier return timing to the Grande Ronde River the fallbrood was harvested at a higher rate (9.4 adults harvested per 1,000 smolts released compared to 8.0 for the standard production). A paper that reports on first generation results of this experiment is in the journal *Fisheries Management and Ecology* (Clarke et al. 2017). We are still tabulating data from the second generation (brood years 2008-2011) of the experiment; however, preliminary results suggest that adult run timing was more similar between the fall brood and standard production lines, as were SAS, straying and harvest rates.

Compensation Area Goals

Goals for smolt-to-adult return (SAR) rates and the number of adults produced to the compensation area are 0.68% and 9,184 for the Grande Ronde basin (Wallowa stock) and 0.61% and 2,000 for the Imnaha basin (Imnaha stock). Compensation plan adult return goals are to the watershed above Lower Granite Dam. To provide a cumulative summary of disposition for all adults that returned to the compensation area, we expanded CWT recoveries to account for the non-CWT fish that returned.

For the Wallowa stock, we estimate that in the 2014-15 run year, 9,300 hatchery origin adults returned to the compensation area, representing 101.2% of the compensation area goal (Table 14). For the Imnaha stock, we estimate that 4,017 adults returned to the compensation area, representing 200.9% of the compensation area goal. Age composition of returning adults is shown in Table 15. Development of the compensation plan goals assumed that twice as many adult steelhead would be harvested in downriver fisheries as return to the compensation area (USACOE 1975); however, that harvest level was not reached for either stock.

There are three principal factors that influence success in meeting the compensation goals: number of smolts released for the brood years that produced the adults; SAS rates to the mouth of the Columbia River; and capture of fish below the compensation area in fisheries and as out-of-basin strays. Over the history of the LSRCP project, we have now reached our adult production compensation goal twelve times (1997-98, 2001-02, 2003-04, 2004-05, 2005-06, 2006-07, 2007-08, 2008-09, 2009-10, 2010-11, 2011-12 and 2014-15 run years) for the Wallowa program, and fourteen times for the Imnaha program (1992-93, 2001-02, 2002-03, 2003-04, 2004-05, 2005-06, 2006-07, 2007-08, 2008-09, 2009-10, 2010-11, 2011-12, 2013-14 and 2014-15 run years). For both the Grande Ronde and Imnaha programs, we have met our smolt production goals in most years. Returns in the 2014-15 run year represent the final returns of the 2010 BY. For the 2010 BY, SAS for the Wallowa and Imnaha stocks were below average at 0.46% and 0.79%, respectively (Figure 3). Smolt-to-adult return to the compensation area above Lower Granite Dam has reached our goal in only eleven of the last 26 brood years for Wallowa and twelve of the last 26 brood years for Imnaha stocks (Figure 4). This suggests that low SAS rates may be the primary factor for only occasionally achieving our adult compensation goals. However, the SAR compensation area goal has been reached in each of the last twelve years for Imnaha stock and in ten of the last twelve years for Wallowa stock. For the Wallowa stock, 29% of the CWT recoveries in the 2014-15 run year occurred downstream of the compensation area. For Imnaha stock, 15% of the recoveries in the 2014-15 run year occurred downstream of the compensation area.

The Imnaha steelhead supplementation program allows us to evaluate and compare productivity (adult progeny produced per parent) of hatchery and naturally spawning fish. Hatchery and natural origin fish are used both for hatchery spawning and they are passed above the weir to spawn naturally; therefore, progeny-per-parent ratios include both hatchery and natural origin parents. Progeny-per-parent ratios for naturally spawning fish were below 1.0 for completed brood years 1987-1994, 1998, and 2001-2009 and above 1.0 for completed brood years 1995-1997, 1999, and 2000 (Figure 5). Progeny-per-parent ratios for fish spawned in the hatchery (weir returns only) have been above 1.0 for all brood years except 1991. Hatchery

ratios exceeded natural ratios for all brood years except for the 1991 and 1997 broods. One purpose of the supplementation program is to enhance or stabilize natural fish abundance. Annual abundance of naturally-produced fish has been highly variable and in recent years adult returns have been lower; however, the long-term pattern suggests an increasing trend in natural returns (Figure 6).




Figure 1. Length-at-age relationships based on scale analysis for hatchery adult returns of one freshwater age (top) and two freshwater age (middle) Wallowa stock summer steelhead from 1991 to 2014, and in 2015 (bottom).




Figure 2. Length-at-age relationships based on scale analysis for hatchery adult returns of one freshwater age (top), and two and three freshwater age (middle) Imnaha stock summer steelhead from 1991 to 2014, and in 2015 (bottom).




Figure 3. Smolt-to-adult survival (SAS) for Wallowa and Imnaha stock summer steelhead, 1985-2010 brood years. Data is based on CWT recoveries.




Figure 4. Smolt-to-adult return (SAR) to the compensation area above Lower Granite Dam for Wallowa and Imnaha stock summer steelhead, 1985-2010 brood years. The Wallowa stock goal is 0.68% and the Imnaha stock goal is 0.61%. Data is based on CWT recoveries.




Figure 5. Progeny-to-parent ratios for Little Sheep Creek summer steelhead, 1987-2010 brood years. Both types of spawning include hatchery and natural origin parents. Dotted line represents replacement (P:P ratio = 1.0). Natural origin steelhead data for the 2010 brood year is not yet available.




Figure 6. Returns of naturally produced summer steelhead to Little Sheep Creek, run years 1984-85 to 2014-15.




Figure 7. Hatchery returns to the compensation area above Lower Granite Dam for Wallowa and Imnaha Stock summer steelhead. The compensation goal for Wallowa stock is 9,184 and the goal for Imnaha stock is 2,000 adults. Data is based on CWT recoveries.

Table 1. Summary of egg collection and juvenile survival for 2014 brood year summer steelhead released in the Grande Ronde and Imnaha river basins at LSRCP facilities in 2015.

Stock	Number of eggs taken	Eyed embryos	Total smolts released	Estimated survival rate	
				Egg-to-embryo	Embryo-to-smolt ^a
Wallowa	1,274,100	1,160,900 ^b	821,192	91.1	87.2
Imnaha	333,600	304,600 ^c	247,642	91.3	94.4

^a Embryos that were culled from or not part of production were subtracted from the calculation of embryo-to-smolt survival.

^b Includes 113,200 eggs lost due to shocking and 104,600 embryos that were euthanized because they were excess to program needs. Includes 1,600 embryos that were transferred to the Salmon and Trout Enhancement Program (STEP) Coordinator.

^c Includes 29,000 eggs lost due to shocking. Includes 39,500 eggs euthanized because they were excess to the program needs.

Table 2. Estimates of fin clip quality and coded-wire tag retention for 2014 brood year summer steelhead reared at Irrigon Fish Hatchery and released in 2015. Experimental group indicates treatment and rearing raceway number. Wallowa and Imnaha stocks were intended to be 100% adipose fin-clipped. Fall brood (Fall B., progeny of broodstock collected in early fall) were AdRV (adipose + right ventral fin-clipped) if they were released at Wallowa Hatchery and Ad only at Big Canyon. Wire tagged standard production (Prod.) groups were Ad+CWT, except for pond 8 which remained AdLV+CWT. Wire tagged fall brood were 100% AdRV+CWT if released at Wallowa and 100% Ad + CWT at Big Canyon.

Experimental Group, Raceway	Tag code	No. checked		Percent							
		CWT	Ad ^a	CWT+ Clip ^b	CWT+ noclip	NoCWT + clip	NoCWT + noclip	Ad	No Ad	RV	No RV
<i>Wallowa Stock</i>											
Fall B., 07	090806	511	-	98.6	0.6	0.8	0.0	-	-	97.4	2.6
Fall B., 21	090807	505	-	100.0	0.0	0.0	0.0	-	-	98.9	1.1
Fall B., 23	090815	512	-	99.6	0.4	0.0	0.0	-	-	-	-
Fall B., 15	090813	545	-	100.0	0.0	0.0	0.0	-	-	-	-
Prod., 08	090808	505	-	93.5	6.5	0.0	0.0	-	-	-	-
Prod., 10	090809	526	-	98.3	1.7	0.0	0.0	-	-	-	-
Prod., 14	090812	522	-	99.0	0.8	0.2	0.0	-	-	-	-
Prod., 16	090810	516	-	98.3	1.5	0.2	0.0	-	-	-	-
Prod., 18	090814	500	-	99.6	0.4	0.0	0.0	-	-	-	-
Prod., 22	090811	506	-	100.0	0.0	0.0	0.0	-	-	-	-
Average	-	515	304	98.7	1.2	0.1	0.0	99.6	0.4	96.8	2.9
<i>Imnaha Stock</i>											
Prod., 28	090805	524	300	99.6	0.4	0.0	0.0	100.0	0.0	-	-
Overall Avg.		516	303	98.8	1.1	0.8	0.0	99.6	0.4	98.1	1.9

^a Adipose fin (Ad) clip quality checks occurred during pre-release sampling at acclimation ponds.

^b A programmatic decision to discontinue ventral fin clipping to indicate the presence of a CWT began with brood year 2013. Fish in pond 8 were left ventral fin clipped (CWT+AdLV) to determine the effect of ventral fin clips on post release survival.

Table 3. Details of experimental and production groups of 2014 brood year summer steelhead released in the Grande Ronde (Wallowa stock) and Imnaha (Imnaha stock) river basins in 2015. Experimental group indicates release strategy and rearing raceway number(s). All groups were reared at Irrigon Hatchery and acclimated. Target size was 113 g (\pm SD) for Wallowa stock and 100 g for Imnaha stock. LGD indicates Lower Granite Dam; percent migration includes \pm 95% confidence intervals. This table does not include 39,803 smolts reared at Lyons Ferry Hatchery in Washington and released into Spring Creek.

Experimental group, Raceway	Transfer date	Release date	Tag code	Fork length (mm)	Weight (g)	Condition factor	Total fish Released	PIT tags released	Percent migration to LGD ^a
<i>Wallowa stock</i>									
<i>Spring Creek</i>									
Production, 8	Jan 20-22	April 5-6	090808	216 (24)	115.2 (32.6)	1.07 (0.08)	24,286	1,893	59.0 \pm 20.8
Production, 8,10,16	Jan 20-22	April 5-6	-	214 (24)	107.2 (33.2)	1.06 (0.07)	61,010	-	-
Production, 10 ^b	Jan 20-22	April 5-6	090809	N/A	N/A	N/A	26,964	1,202	78.7 \pm 48.0
Production, 16 ^b	Feb 20	April 5-6	090810	N/A	N/A	N/A	24,828	1,102	70.8 \pm 36.8
Fall Brood, 7 ^b	Jan 20-22	April 4-6	090806	N/A	N/A	N/A	25,586	4,374	87.7 \pm 21.2
Fall Brood,7,9,11,12	Jan 20-22	April 4-6	-	215 (22)	110.0 (41.9)	1.11 (0.07)	137,807	-	-
Production, 22 ^b	April 6-7	April 20-30	090811	N/A	N/A	N/A	27,383	2,491	109.5 \pm 56.0
Production, 22, 24	April 6-7	April 20-30	-	213 (20)	103.7 (30.4)	1.00 (0.05)	57,288	-	-
Fall Brood, 21	April 6-7	April 20-30	090807	N/A	N/A	N/A	26,707	2,398	95.1 \pm 35.9
Fall Brood, 19,21 ^b	April 6-7	April 20-30	-	215 (21)	100.0 (25.3)	1.00 (0.07)	61,123	-	-
<i>Deer Creek</i>									
Fall Brood, 15	Feb 23	April 15-20	090813	N/A	N/A	N/A	24,947	1,593	134.7 \pm 83.7
Fall Brood, 15,17 ^b	Feb 23	April 15-20	-	225 (19)	118.4 (31.0)	1.08 (0.08)	24,706	-	-
Production, 18	Feb 24	April 16-20	090814	N/A	N/A	N/A	26,329	2,982	94.3 \pm 32.9
Production, 13,17,18 ^b	Feb 24	April 16-20	-	218 (21)	116.5 (34.1)	1.10 (0.24)	95,617	-	-
Fallbrood, 23	April 20	May 1-12	090815	N/A	N/A	N/A	25,018	2,297	89.3 \pm 34.2
Fallbrood, 23,25 ^b	April 20	May 1-12	-	213 (18)	103.3 (28.1)	1.00 (0.06)	61,618	-	-
Production, 20,26	April 21	May 2-12	-	209 (19)	92.4 (22.8)	1.01 (0.08)	89,975	2,292	133.5 \pm 63.0
Total released							821,192	22,624	
<i>Cottonwood Creek, WA</i>									
Cottonwood, 14	Feb 2	Mar 23-29	090812	N/A	N/A	N/A	25,260	3,994	84.8 \pm 6.4
Cottonwood, 14	Feb 2	Mar 23-29	-	191 (18)	81.6 (23.8)	1.18 (0.07)	18,740	-	-
Total released							44,000	3,994	
<i>Imnaha stock</i>									
<i>Little Sheep</i>									
Production, 28	Feb 25-27	April 1-28	090805	N/A	N/A	N/A	23,692	6,245	85.5 \pm 11.4
Production, 27-32 ^c	Feb 25-27	April 1-28	-	216 (16)	105.0 (27.9)	1.01 (0.06)	223,950	8,653	84.4 \pm 8.9
Total released							247,642	14,898	

^a Percent of PIT tag release groups that migrated to Lower Granite Dam are Cormack-Jolly-Seber estimated survival probabilities. Values exceeding 100% occur when true survival is close to 100% and/or when PIT tag detection variability is high (Smith et al. 2000). ^bThis release group was indistinguishable based on external marks from other groups in the same acclimation pond, therefore unique size measurements could not be taken. ^cPercent migration to LGD was calculated from PIT tags in raceways 30 and 32.

Table 4. Density ($\pm 95\%$ confidence interval) and mean fork length (standard deviation in parentheses) of residual hatchery steelhead, wild rainbow trout/juvenile steelhead, and juvenile chinook salmon from index sites on Deer (Grande Ronde basin) and Little Sheep (Imnaha basin) creeks in 2015. Hatchery steelhead were classified as residuals after 20 June. HSTS indicates residual hatchery steelhead, WSTS indicates wild rainbow trout/juvenile steelhead for ages one and older, and WChS indicates juvenile (age 0+) spring chinook salmon.

Location ^a	Date	Species	Area (m ²)	Size of fish (mm)			Density ^b (fish/100m ²)
				N	Fork length	Range	
<i>Grande Ronde basin</i>							
Deer Cr.	23 July	HSTS	449.6	12	155.1 (33.9)	129-245	3.83 \pm 4.6
Deer Cr.	23 July	WSTS	449.6	22	99.1 (15.5)	84-136	6.5 \pm 3.2
Deer Cr.	23 July	WChS	449.6	137	77.6 (5.0)	68-88	39 \pm 7.7
<i>Imnaha basin</i>							
Little Sheep Cr.	21 July	HSTS	452.9	63	183.3 (40.5)	103-276	16.1 \pm 1.7
Little Sheep Cr.	21 July	WSTS	452.9	8	94.1 (20.2)	59-114	1.79 \pm 0.3

^a Index sites located on Deer Creek (Rkm 0.1) at Big Canyon Facility and on Little Sheep Creek (Rkm 8.0) at Little Sheep Creek Facility. Two adjacent sites were sampled at each location and each site typically included both riffle and pool habitat.

^b Density ($\pm 95\%$ confidence interval) was determined using a multiple pass removal method (Zippen 1958) with a backpack electrofisher (Smith-Root Model 12) and block seines.

Table 5. Travel days to Lower Granite Dam (LGD) and percent of the PIT tagged releases that were uniquely detected at the four lower Snake River dams for Little Sheep Creek smolts that departed the acclimation pond during the early, middle, and late periods of the volitional release. The early and late time periods were defined as the first and last 25% of the release period, which was typically a minimum of 28 days. The middle time period includes 50% of the release period or 14 days.

Brood Year	Travel Days to LGD			Percent Detected		
	Early	Middle	Late	Early	Middle	Late
2008	34.2	27.5	27.2	64.1	62.6	59.3
2009	41.7	33.4	32.8	39.1	40.9	39.5
2010	42.5	36.7	27.7	56.9	47.2	46.0
2011	35.4	28.8	22.3	56.5	47.0	49.3
2012	39.5	33.8	22.5	52.4	49.3	43.6
2013	42.0	37.0	22.0	63.5	57.9	63.0
2014 ^a	N/A	N/A	30.0	N/A	N/A	14.5
Mean	39.2	32.9	25.8	55.4	50.8	50.1

^a Due to equipment malfunction, PIT tag detection at Little Sheep acclimation pond was from April 23 to 28. As a result, data analysis was only able to be performed on the late group. This analysis included only 5 days of detections instead of the usual 7.

Table 6. Number of PIT tags released and unique adult PIT tag detections at Bonneville and Lower Granite dams during the 2014-15 run year by stock and brood year.

Brood year	PIT tags released	Age at return	Adult Detections	
			Bonneville Dam	Lower Granite Dam
<i>Wallowa Stock</i>				
2010	22,187	5	0	0
2011	22,381	4	201	131
2012	21,875	3	278	219
<i>Total</i>	66,443		479	350
<i>Imnaha Stock</i>				
2010	21,900	5	0	0
2011	21,937	4	166	119
2012	21,882	3	364	250
<i>Total</i>	65,719		530	369

Table 7. Timing of adult steelhead returns to LSRCF facilities in 2015 by location and origin.

Period	Week of the year	Number of fish trapped ^a					
		Wallowa		Big Canyon		Little Sheep	
		Hatchery	Natural	Hatchery	Natural	Hatchery	Natural
Jan 22-28	4	68	0	-	-	-	-
Jan 29-Feb 04	5	101	0	-	-	-	-
Feb 05-11	6	266	0	-	-	-	-
Feb 12-18	7	279	0	0	0	-	-
Feb 19-25	8	234	0	3	0	0	0
Feb 26-Mar 04	9	259	4	0	0	10	0
Mar 05-11	10	174	0	0	0	13	0
Mar 12-18	11	237	1	168	3	141	2
Mar 19-25	12	143	2	173	2	169	10
Mar 26-Apr 01	13	255	2	120	2	159	9
Apr 02-Apr 08	14	100	2	284	12	43	3
Apr 09-15	15	61	2	126	7	133	11
Apr 16-22	16	21	2	97	5	126	14
Apr 23-29	17	9	0	106	20	180	29
Apr 30-May 06	18	5	0	28	6	68	10
May 07-13	19	0	0	3	1	27	4
May 14-20	20	0	0	0	0	18	1
May 21-27	21	-	-	-	-	0	0
May 28-June 03	22	-	-	-	-	0	0
Jun 04-10	23	-	-	-	-	-	-
Jun 11-17	24	-	-	-	-	-	-
June 18-24	25	-	-	-	-	-	-
Total		2,212	15	1,108	58	1,087	93

^aThe ladder was opened on 27 January at Wallowa Fish Hatchery, and weirs were installed 17 February at Big Canyon Facility (Deer Creek) and 20 February at Little Sheep Creek Facility. Adult collections ended 18 May at Wallowa Fish Hatchery, 18 May at Big Canyon Facility, and 1 June at Little Sheep Creek Facility.

Table 8. Number, disposition, and mean fork length (mm) of adult steelhead that returned to LSRCP facilities in 2015 by stock, origin, estimated age (freshwater:saltwater), and gender. M indicates male and F indicates female.

Facility, stock, disposition	Hatchery ^a									Natural ^b									Grand total		
	1:1		1:2		2:1		1:3		Total	2:1		2:2		2:3		3:1		3:2		Total	
	M	F	M	F	M	F	M	F		M	F	M	F	M	F	M	F	M			F
<i>Wallowa Hatchery (Wallowa Stock-Production)</i>																					
Trapped	582	159	218	510	4	2	0	3	1,478	2	2	3	3	0	0	2	2	0	3	17	1,495
Passed	0	0	0	0	0	0	0	0	0	2	2	3	3	0	0	2	2	0	3	17	17
Outplanted	88	2	13	20	0	0	0	0	123	0	0	0	0	0	0	0	0	0	0	0	123
Kept	492	157	205	490	4	2	0	3	1,355	0	0	0	0	0	0	0	0	0	0	0	1,355
Mortality	1	0	5	1	0	0	0	0	7	0	0	0	0	0	0	0	0	0	0	0	7
Spawned ^c	49	3	63	109	1	1	0	0	226	0	0	0	0	0	0	0	0	0	0	0	226
Killed ^d	444	152	137	382	3	1	0	3	1,122	0	0	0	0	0	0	0	0	0	0	0	1,122
<i>Wallowa Hatchery (Wallowa Stock-Fallbrood)</i>																					
Trapped	396	105	75	149	2	5	0	0	732	0	0	0	0	0	0	0	0	0	0	0	732
Passed	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Outplanted	33	5	25	0	1	0	0	0	64	0	0	0	0	0	0	0	0	0	0	0	64
Kept	363	100	50	149	1	5	0	0	668	0	0	0	0	0	0	0	0	0	0	0	668
Mortality	4	1	2	0	0	0	0	0	7	0	0	0	0	0	0	0	0	0	0	0	7
Spawned	102	36	26	90	1	3	0	0	258	0	0	0	0	0	0	0	0	0	0	0	258
Killed ^d	257	63	22	59	0	2	0	0	403	0	0	0	0	0	0	0	0	0	0	0	403
<i>Wallowa Hatchery (Total Returns)</i>																					
Trapped	978	264	293	659	6	7	0	3	2,210	2	2	3	3	0	0	2	2	0	3	17	2,227
Fork length (mm)	559	565	719	671	-	-	-	-		-	-	-	-	-	-	-	-	-	-		
Standard deviation	40	34	40	36	-	-	-	-		-	-	-	-	-	-	-	-	-	-		
Sample size	42	35	39	39	-	-	-	-		-	-	-	-	-	-	-	-	-	-		

Table 8. Continued.

Facility, stock, Disposition	Hatchery ^a									Natural ^b									Grand Total		
	1:1		1:2		2:1		1:3		Total	2:1		2:2		2:3		3:1		3:2		Total	
	M	F	M	F	M	F	M	F		M	F	M	F	M	F	M	F	M	F		Total
<i>Big Canyon Facility (Wallowa stock)</i>																					
Trapped ^g	392	190	119	394	2	5	0	3	1,105	12	8	5	10	0	0	10	7	1	5	58	1,163
Passed	0	0	0	0	0	0	0	0	0	12	8	5	10	0	0	10	7	1	5	58	58
Outplanted	27	30	3	0	0	0	0	0	60	0	0	0	0	0	0	0	0	0	0	0	60
Kept	365	160	116	394	2	5	0	3	1,045	0	0	0	0	0	0	0	0	0	0	0	1,040
Mortality	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Spawned ^c	1	11	0	14	0	0	0	0	26	0	0	0	0	0	0	0	0	0	0	0	26
Killed ^d	364	149	116	380	2	5	0	3	1,019	0	0	0	0	0	0	0	0	0	0	0	1,019
Fork length (mm)	-	-	-	-	-	-	-	-	-	577	702	589	664	-	-	-	-	-	-	-	-
Standard deviation	-	-	-	-	-	-	-	-	-	47	40	70	7	-	-	-	-	-	-	-	-
Sample size	-	-	-	-	-	-	-	-	-	4	2	3	3	-	-	-	-	-	-	-	-
<i>Little Sheep Creek Facility (Imnaha stock)</i>																					
Trapped	523	246	46	262	3	0	0	0	1,080	25	24	11	22	0	0	5	4	0	2	93	1,173
Passed	30	21	2	18	0	0	0	0	71	15	17	7	16	0	0	0	2	0	0	57	128
Outplanted	175	80	17	82	1	0	0	0	355	0	0	0	0	0	0	0	0	0	0	0	355
Kept	318	145	27	162	2	0	0	0	654	10	7	4	6	0	0	5	2	0	2	36	690
Mortality	1	0	1	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	2
Spawned	35	15	11	34	1	0	0	0	96	10	7	4	6	0	0	5	2	0	2	36	132
Killed ^{d,e}	282	130	15	128	1	0	0	0	556	0	0	0	0	0	0	0	0	0	0	0	556
Fork length (mm)	568	562	692	680	-	-	-	-	-	553	561	697	667	-	-	-	549	-	-	-	-
Standard deviation	23	13	36	28	-	-	-	-	-	28	44	19	37	-	-	-	26	-	-	-	-
Sample size	14	6	3	11	-	-	-	-	-	7	18	4	15	-	-	-	2	-	-	-	-

^a Wallowa and Imnaha stock ages apportioned using CWT data and 165 (Wallowa stock) and 80 (Imnaha stock) scale samples collected in 2015. Mean fork lengths are from fish with scale samples collected in 2015.

^b Ages of natural steelhead from the Wallowa basin were apportioned using historical data (229 samples) and 2015 data (12 samples); Little Sheep Creek Facility natural steelhead ages apportioned using 2014 data (65 samples) and 2015 data (37 samples). Mean fork lengths are from fish with scale samples collected in 2015.

^c One male and twenty-five females were spawned for Washington Department of Fish and Wildlife.

^d For Wallowa stock steelhead, 952 fish that returned to Wallowa Hatchery and 243 fish that returned to Big Canyon Facility were euthanized and donated to local food banks. In addition, 38 fish from Wallowa Hatchery and 104 fish from Big Canyon Facility were euthanized and donated to local schools for educational purposes. For Imnaha stock steelhead, 261 fish that returned to Little Sheep Creek Facility were euthanized and donated to local food banks.

^e Includes 19 natural males that were live-spawned and passed above the weir.

^f Includes 6 hatchery males and 1 hatchery females that were initially outplanted to Big Sheep Creek. These 7 fish were subsequently recaptured at the weir and euthanized.

^g One Male and 4 female steelhead had an AdRV clip and were included in these numbers. With no CWT, we could not determine if they were strays or a mis-clip.

Table 9. Number of adult summer steelhead trapped at the Little Sheep Creek Facility weir that were either outplanted to Big Sheep Creek or passed above the weir, and were subsequently recaptured, 1999-2015.

Year	Big Sheep Creek			Little Sheep Creek		
	Number of fish		%	Number of fish		%
	Outplanted	Recaptured ^a	Recaptured ^b	Passed ^c	Recaptured ^a	Recaptured ^b
1999	42	6	14.3	80	1	1.3
2000	138	17	12.3	200	9	4.5
2001	354	48	13.6	784	89	11.4
2002	2,030	907	44.7	1,198	269	22.5
2003	1,403	439	31.3	387	36	9.3
2004	1,719	244	14.2	823	138	16.8
2005	1,555	109	7.0	461	37	8.0
2006	1,934	703	36.3	356	53	14.9
2007	1,315	168	12.8	241	14	5.8
2008	1,365	382	28.0	291	23	7.9
2009	869	394	45.3	281	15	5.3
2010	1,450	166	11.4	346	6	1.7
2011	401	154	38.4	306	2	0.7
2012	350	175	50.0	241	13	5.4
2013	58	5	8.6	245	20	8.2
2014	232	29	12.5	270	1	0.4
2015	362	10	2.8	147	1	0.7
Mean	-	-	22.6	-	-	7.3

^a Total number of recaptures, including multiple recaptures. For 1999-2002, recaptures were opercle punched at the weir and second and third time recaptures recorded.

^b Total recaptured divided by total outplanted.

^c Includes natural males that were live-spawned and passed above the weir.

Table 10. Summary of summer steelhead spawning surveys in Deer Creek above the Big Canyon Facility weir, 2002-2015. In prior years an index of redd visibility had been reported in this table; however, due to inconsistent methods for making calculations, a new method is being evaluated and will be available in the future. Note that data for 2005 has been updated to reflect an adjusted number of fish passed above the weir.

Year	Passed			Redds counted	Fish per redd	Females per redd	% Redds counted ^a	Redds per mile ^b
	Females	Males	Total					
2002	120	89	209	84	2.49	1.43	70	8.4
2003	92	48	140	64	2.19	1.44	70	6.4
2004	47	20	67	46	1.46	1.02	98	4.6
2005	42	35	76	35	2.20	1.20	83	3.5
2006 ^c	55	41	96	58	1.66	0.95	105	5.8
2007	27	21	48	41	1.17	0.66	152	4.1
2008	23	38	61	15	4.07	1.53	65	1.5
2009	42	38	80	21	3.81	2.00	50	2.1
2010	85	49	134	84	1.60	1.01	99	8.4
2011	75	58	133	28	4.75	2.68	37	2.8
2012	34	34	69	22	3.09	1.54	65	2.2
2013	41	22	63	33	1.91	1.24	80	3.3
2014 ^d	18	30	48	18	2.67	1.00	100	1.8
2015 ^e	34	32	66	49	1.35	0.69	144	4.9

^a Calculated as number of redds counted ÷ number of females passed x 100. Assumes each female built one redd.

^b Twelve miles of stream were surveyed in 2002, 2003, 2007-2010, and in 2012-2015. Ten miles of stream were surveyed in 2004-06 and in 2011. Redds per mile are based on the lower ten miles, since redds have not been observed between RM 10-12.

^c Includes an estimated seven additional hatchery steelhead (4 females and 3 males) that escaped above the weir prior to weir installation, based on marked and unmarked fallbacks at weir.

^d Includes an estimated 3 additional hatchery steelhead (1 female and 2 males) that escaped above the weir prior to weir installation. However, the total passed column does not include 3 steelhead passed above the weir after May 16, 2014 because stream surveys were discontinued prior to that date.

^e Estimate includes 9 additional steelhead (5 males, 4 females) that escaped above the weir prior to installation in February, based on marked and unmarked fallbacks recovered at the weir.

Table 11. Spawning summaries for summer steelhead at LSRCP facilities in 2015. The percent mortality is from green egg to eyed embryo after shocking.

Spawn date, lot number	Parental origin ^a	Number of females spawned ^b	Number of eggs	Eyed embryos ^c	% mortality
<i>Wallowa Hatchery (Wallowa stock)</i>					
3/4, WA520	Production	16	88,200	82,300	6.7
	Fall Broodstock	64	317,300	272,900	14.0
3/11, WA521	Production	22	116,900	109,500	6.3
	Fall Broodstock	38	184,100	165,900	9.9
3/18, WA522	Production	26	141,500	133,600	5.6
	Fall Broodstock	20	98,700	87,300	11.6
3/25, WA523	Production	22	125,700	117,200	6.8
	Fall Broodstock	0	0	0	0.0
4/1, WA524	Production	18	113,900	106,200	6.8
	Fall Broodstock	6	30,100	30,100	0.0
4/8, WA525	Production	10	57,700	55,900	3.1
	Fall Broodstock	0	0	0	0.0
Subtotal	Production	114	643,900	604,700	6.1
	Fall Broodstock	128	630,200	556,200	11.7
Total		242	1,274,100	1,160,900	8.9
<i>Little Sheep Creek Facility (Imnaha stock)</i>					
3/24, LI620	Hatchery	8	27,312	37,200	14.9
	Mixed		16,388		
3/31, LI621	Hatchery	8	16,537	42,700	3.2
	Mixed		27,563		
4/07, LI622	Hatchery	3	11,467	14,900	13.4
	Mixed		5,733		
4/14, LI623	Hatchery	14	42,571	68,000	8.7
	Mixed		31,929		
4/21, LI624	Hatchery	20	30,320	66,400	12.4
	Mixed		45,480		
4/28, LI625	Hatchery	17	36,847	75,400	3.7
	Mixed		41,453		
Subtotal	Hatchery	70	165,054	304,600	8.7
	Mixed		168,546		
Total		70	333,600	304,600	8.7

^a Family groups were usually one male x one female for Wallowa stock and were matrix spawned (three males x three females) for Imnaha stock. Mixed eggs include both natural and hatchery parents.

^b Number of males spawned equals the number of females spawned.

^c Includes 1,600 Wallowa production stock eyed embryos that were transferred to the Salmon and Trout Enhancement Program (STEP). Also includes 49,200 eyed embryos from Wallowa Production and 55,400 eyed embryos from Wallowa Fall broodstock and 39,500 eyed embryos from Little Sheep brood stock that were euthanized because they were excess to program needs.

Table 12. Summary of anadromous adult recoveries of coded-wire tagged (CWT) Wallowa stock summer steelhead for the 2014-15 run year. All CWT fish were of hatchery origin and were released into either Deer Creek (at Big Canyon Facility) or Spring Creek (at Wallowa Hatchery). Data were summarized as available through August 2017.

Brood year, release site	Experimental group ^a	CWT code	Recoveries at weirs ^b	Other in-basin recoveries ^c	Out-of-basin recoveries ^d	Total recoveries
2010						
Spring Cr.	Production, April	090321	1	0	0	1
2011						
Deer Cr.	Production, April	090429	38	76	110	224
	Production, May	090430	24	57	85	166
Spring Cr.	Production, April	090425	51	44	113	208
	Production, April	090426	47	53	114	214
	Production, April	090427	49	33	106	188
	Production, May	090428	42	7	124	173
	Fallbrood, April	090421	37	8	66	111
	Fallbrood, April	090422	28	8	74	110
	Fallbrood, April	090424	25	8	79	112
	Fallbrood, May	090423	22	7	114	143
2012						
Deer Cr.	Production, April	090561	36	43	89	168
	Production, May	090563	27	22	24	73
Spring Cr.	Production, April	090558	73	58	51	182
	Production, April	090559	71	48	73	192
	Production, April	090560	54	18	51	123
	Production, May	090562	51	7	45	103
	Fallbrood, April	090555	61	8	30	99
	Fallbrood, April	090556	51	4	47	102
	Fallbrood, April	090557	78	23	109	210
	Fallbrood, May	090554	42	7	56	105
	Total recoveries		908	539	1,560	3,007

^a Experimental groups include the release strategy. All releases were targeted for four fish per pound (113 g/fish). All fish were acclimated. April releases were forced (over a 24-hour period) and May releases were volitional (1-3 weeks) unless otherwise noted.

^b Actual number of CWT fish that were released into Spring Creek and recovered at the Wallowa Hatchery weir or released into Deer Creek and recovered at the Big Canyon Facility weir. The protocol was to collect all CWT fish at the weirs for sampling at the hatchery during spawning.

^c Estimated number (from creel surveys and harvest card returns) of CWT fish that were harvested in the Grande Ronde River basin fisheries, and in-basin stray recoveries.

^d Estimated number (from PSMFC and ODFW databases) of CWT fish that were recovered in the ocean, mainstem Columbia, Deschutes or Snake river fisheries, or in tributaries outside the Grande Ronde River basin. Unexpanded data were used when CWT expansion factors were 25 or greater due to low sampling rates.

Table 13. Summary of anadromous adult recoveries of coded-wire tagged (CWT) Imnaha stock summer steelhead for the 2014-15 run year. All CWT fish were of hatchery origin and were released into Little Sheep Creek at the Little Sheep Creek Facility. Data were summarized as available through August 2017.

Brood year, release site	Experimental group ^a	CWT code	Recoveries at weirs ^b	Other in-basin recoveries ^c	Out-of-basin recoveries ^d	Total recoveries
2011 Little Sheep	Production, April	090420	35	0	102	137
2012 Little Sheep	Production, April	090553	73	0	256	329
Total recoveries			108	0	358	466

^a Experimental groups include the release strategy. All Little Sheep fish were acclimated and volitionally released over a four-week period.

^b Estimated number of CWT fish recovered at the Little Sheep Creek Facility weir based on actual number recovered at the weir and estimated number either passed above the weir to Little Sheep Creek or outplanted to Big Sheep Creek to spawn naturally.

^c Estimated number (from creel surveys and harvest card returns) of CWT fish that were harvested in the Imnaha River basin fishery.

^d Estimated number (from PSMFC and ODFW databases) of CWT fish that were recovered in the ocean, mainstem Columbia, Deschutes or Snake river fisheries, or in tributaries outside the Imnaha River basin. Unexpanded data were used when CWT expansion factors were 25 or greater due to low sampling rates.

Table 14. Harvest and escapement distribution of adult summer steelhead by recovery location for the 2014-15 run year using the PSMFC and ODFW mark recovery databases. "C and S" indicates ceremonial and subsistence tribal fisheries. Data were summarized as available through August 2017. "-" indicates not sampled or undefined.

Location	Wallowa Stock			Imnaha Stock		
	Estimated CWT recoveries	Total return	Percent of total return	Estimated CWT recoveries	Total return	Percent of total return
Ocean harvest	0	0	0.0	0	0	0.0
Columbia River harvest						
Treaty net	296	974	8.1	8	73	1.6
C and S	0	0	0.0	0	0	0.0
Sport	378	1,186	9.9	42	393	8.5
Test	0	0	0.0	0	0	0.0
Tributary sport	145	440	3.7	17	150	3.2
Deschutes River harvest						
Sport	0	0	0.0	0	0	0.0
C and S	0	0	0.0	0	0	0.0
Strays						
Outside Snake R. basin	45	110	0.9	1	9	0.2
Within Snake R. basin*	3	15	0.1	0	0	0.0
Snake River sport, tribs. harvest*	693	2,321	19.3	290	2,688	57.9
Oregon tributary harvest* ^a	539	3,649	30.4	0	249	5.3
Hatchery weir* ^b	908	3,315	27.6	108	1,080	23.3
Total estimated return	3,007	12,010	100	466	4,642	100
Return to compensation area		9,300			4,017	
Percent of compensation goal		101.3			200.9	

* Indicates areas defining the compensation area. The compensation goal for Wallowa stock is 9,184 adults and the goal for Imnaha stock is 2,000 adults.

^a Harvest in Oregon tributaries are estimates based on angler surveys and harvest card returns.

^b Total returns to the hatchery weir are actual numbers, except for the Imnaha stock where we estimated the number of CWT fish recovered at the Little Sheep Creek Facility weir. This estimate is based on the actual number of CWT fish recovered at the weir and estimated number either passed above the weir to Little Sheep Creek or outplanted to Big Sheep Creek to spawn naturally.

Table 15. Harvest and escapement distribution of adult summer steelhead by age and recovery location for the 2014-15 run year using the PSMFC and ODFW mark recovery databases. "C and S" indicates ceremonial and subsistence tribal fisheries. Data were summarized as available through August 2017. "-" indicates not sampled or undefined.

Location	Total returns by age							
	Wallowa Stock				Imnaha Stock			
	Age 3	Age 4	Age 5	Total	Age 3	Age 4	Age 5	Total
Ocean harvest	0	0	0	0	0	0	0	0
Columbia River harvest								
Treaty net	257	717	0	974	38	35	0	73
C and S	0	0	0	0	0	0	0	0
Sport	414	772	0	1,186	349	44	0	393
Test	0	0	0	0	0	0	0	0
Tributary sport	175	265	0	440	0	150	0	150
Deschutes River harvest								
Sport	0	0	0	0	0	0	0	0
C and S	0	0	0	0	0	0	0	0
Strays								
Outside Snake R. basin	70	40	0	110	9	0	0	9
Within Snake R. basin*	2	13	0	15	0	0	0	0
Snake River sport, tribs. harvest*	775	1,546	0	2,321	2,018	670	0	2,688
Oregon tributary harvest** ^a	1,560	2,089	0	3,649	174	75	0	249
Hatchery weir** ^b	1,824	1,485	6	3,315	769	311	0	1,080
Total estimated return	5,077	6,927	6	12,010	3,357	1,285	0	4,642

* Indicates areas defining the compensation area. The compensation goal for Wallowa stock is 9,184 adults and the goal for Imnaha stock is 2,000 adults.

^a Total returns to Oregon tributaries are harvest estimates based on angler surveys and harvest card returns.

^b Total returns to the hatchery weir are actual numbers, except for the Imnaha stock where we estimated the number of CWT fish recovered at the Little Sheep Creek Facility weir. This estimate is based on the actual number of CWT fish recovered at the weir and estimated number either passed above the weir to Little Sheep Creek or outplanted to Big Sheep Creek to spawn naturally.

REFERENCES

- Carmichael, R.W. 1989. Lower Snake River Compensation Plan - Oregon evaluation studies, five-year study plan. Oregon Department of Fish and Wildlife, Fish Research Project, Portland.
- Carmichael, R.W., and E.J. Wagner. 1983. Evaluation of Lower Snake River Compensation Plan facilities in Oregon. Oregon Department of Fish and Wildlife, Fish Research Project 14-16-0001-83269, Annual Progress Report, Portland.
- Carmichael, R.W., and R.T. Messmer. 1985. Evaluation of Lower Snake River Compensation Plan facilities in Oregon. Oregon Department of Fish and Wildlife, Fish Research Project FRI/LSR-86-35, Annual Progress Report, Portland.
- Carmichael, R.W., B.A. Miller, and R.T. Messmer. 1986a. Lower Snake River Compensation Plan - Oregon evaluation studies. Oregon Department of Fish and Wildlife, Fish Research Project FRI/LSR-86-35, Annual Progress Report, Portland.
- Carmichael, R.W., R. Boyce, and J. Johnson. 1986b. Grande Ronde River Spring Chinook Production Report (U.S. v. Oregon). Oregon Department of Fish and Wildlife, Portland.
- Carmichael, R.W., R.T. Messmer, and B.A. Miller. 1987. Lower Snake River Compensation Plan - Oregon evaluation studies. Oregon Department of Fish and Wildlife, Fish Research Project FRI/LSR-88-16. Annual Progress Report, Portland.
- Carmichael, R.W., R.T. Messmer, and B.A. Miller. 1988a. Lower Snake River Compensation Plan - Oregon evaluation studies. Oregon Department of Fish and Wildlife, Fish Research Project AFFI/LSR-90-17. Annual Progress Report, Portland.
- Carmichael, R.W., B.A. Miller, and R.T. Messmer. 1988b. Summer steelhead creel surveys in the Grande Ronde, Wallowa, and Imnaha rivers for the 1987-88 run year. Oregon Department of Fish and Wildlife, Fish Research Project AFFI-LSR-89-02. Annual Progress Report, Portland.
- Carmichael, R.W., M.W. Flesher, and R.T. Messmer. 1989. Summer steelhead creel surveys in the Grande Ronde, Wallowa, and Imnaha rivers for the 1988-89 run year. Oregon Department of Fish and Wildlife, Fish Research Project AFFI-LSR-90-12. Annual Progress Report, Portland.
- Carmichael, R.W., M.W. Flesher, and R.T. Messmer. 1990. Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Imnaha rivers for the 1989-90 run year. Oregon Department of Fish and Wildlife, Fish Research Project AFFI-LSR-91-12. Annual Progress Report, Portland.
- Carmichael, R.W., D.L. Eddy, M.W. Flesher, M. Keefe, P.J. Keniry, S.J. Parker, and T.A. Whitesel. 1999. Lower Snake River Compensation Plan: Oregon evaluation studies. Oregon Department of Fish and Wildlife, 1994 Annual Progress Report, Portland.

- Carmichael, R.W., D.L. Eddy, M.W. Flesher, T.L. Hoffnagle, P.J. Keniry, and J.R. Ruzycki. 2004. Lower Snake River Compensation Plan: Oregon evaluation studies. Oregon Department of Fish and Wildlife, 1995 and 1996 Bi-Annual Progress Report, Portland.
- Carmichael, R.W., D.L. Eddy, M.W. Flesher, G.C. Grant, J.R. Ruzycki, and G.R. Vonderohe. 2005a. Lower Snake River Compensation Plan: Oregon summer steelhead evaluation studies. Oregon Department of Fish and Wildlife, 1997 and 1998 Bi-Annual Progress Report, Portland.
- Carmichael, R.W., D.L. Eddy, M.W. Flesher, G.C. Grant, J.R. Ruzycki and G.R. Vonderohe. 2005b. Lower Snake River Compensation Plan: Oregon Summer Steelhead Evaluation Studies. Oregon Department of Fish and Wildlife, 1999 and 2000 Bi-Annual Progress Report, Portland.
- Carmichael, R.W., and T.L. Hoffnagle. 2012. Lower Snake River Compensation Plan: Oregon evaluation studies. Work Statement submitted to the U.S. Fish and Wildlife Service, Contract Number F13AC00034. Lower Snake River Compensation Plan Office, Boise, ID.
- Carmichael, R.W., T.L. Hoffnagle, and L.R. Clarke. 2013. Lower Snake River Compensation Plan: Oregon evaluation studies. Work Statement submitted to the U.S. Fish and Wildlife Service, Contract Number F14AC00042. Lower Snake River Compensation Plan Office, Boise, ID.
- Clarke, L.R., M. W. Flesher, T. A. Whitesel, G. R. Vonderohe, and R. W. Carmichael. 2010. Post-release performance of acclimated and direct-released hatchery summer steelhead into Oregon tributaries of the Snake River. *North American Journal of Fisheries Management* 30:1098-1109.
- Clarke, L.R., M. W. Flesher, S. M. Warren, and R. W. Carmichael. 2011. Survival and straying of hatchery steelhead following forced or volitional release. *North American Journal of Fisheries Management* 31:116-123.
- Clarke, L.R., M. W. Flesher, and R. W. Carmichael. 2014. Hatchery steelhead smolt release size effects on adult production and straying. *North American Journal of Aquaculture* 76:39-44.
- Clarke, L.R., M.W. Flesher, S.M. Warren, D.L. Eddy, and R.W. Carmichael. 2014. Lower Snake River Compensation Plan: Oregon Summer Steelhead Evaluation Studies. Oregon Department of Fish and Wildlife, 2012 Annual Progress Report, Salem.
- Clarke, L.R., M.W. Flesher, H.M. Stanton, D.L. Eddy, and R.W. Carmichael. 2015. Lower Snake River Compensation Plan: Oregon Summer Steelhead Evaluation Studies. Oregon Department of Fish and Wildlife, 2013 Annual Progress Report, Salem.
- Clarke, L.R., M. W. Flesher, W.J. Knox, and R. W. Carmichael. 2017. Increased harvest of anadromous hatchery steelhead *Oncorhynchus mykiss* (Walbaum) through return timing manipulation. *Fisheries Management and Ecology* 24: 274-282.

- Flesher, M.W., R.W. Carmichael, and R.T. Messmer. 1991. Summer steelhead creel surveys in the Grande Ronde, Wallowa, and Imnaha rivers for the 1990-91 run year. Oregon Department of Fish and Wildlife, Fish Research Project, AFF1-92-09. Annual Progress Report, Portland.
- Flesher, M.W., M.A. Buckman, R.W. Carmichael, R.T. Messmer, and T.A. Whitesel. 1992. Summer steelhead creel surveys in the Grande Ronde, Wallowa, and Imnaha rivers for the 1991-92 run year. Oregon Department of Fish and Wildlife, Fish Research Project, AFF1-LSR-94-07. Annual Progress Report, Portland.
- Flesher, M.W., M.A. Buckman, R.W. Carmichael, R.T. Messmer, and T.A. Whitesel. 1993. Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Imnaha rivers for the 1992-93 run year. Oregon Department of Fish and Wildlife, Fish Research Project, AFF1-LSR-94-14. Annual Progress Report, Portland.
- Flesher, M.W., M.A. Buckman, R.W. Carmichael, R.T. Messmer, and T.A. Whitesel. 1994. Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Imnaha rivers for the 1993-94 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Portland.
- Flesher, M.W., R.W. Carmichael, and T.A. Whitesel. 1995. Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Imnaha rivers for the 1994-95 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Portland.
- Flesher, M.W., R.W. Carmichael, and T.A. Whitesel. 1996. Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Imnaha rivers for the 1995-96 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Portland.
- Flesher, M.W., R.W. Carmichael, and T.A. Whitesel. 1997. Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Imnaha rivers for the 1996-97 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Portland.
- Flesher, M.W., R.W. Carmichael, and T.A. Whitesel. 1999. Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Imnaha rivers for the 1997-98 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Portland.
- Flesher, M.W., R.W. Carmichael, T.A. Whitesel, and J.R. Ruzycski. 2000. Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Imnaha rivers for the 1998-99 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Portland.
- Flesher, M.W., R.W. Carmichael, and J.R. Ruzycski. 2001. Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Imnaha rivers for the 1999-2000 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Portland.

- Flesher, M.W., R.W. Carmichael, and J.R. Ruzycki. 2004a. Lower Snake River Compensation Plan: Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Imnaha rivers for the 2000-01 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Portland.
- Flesher, M.W., R.W. Carmichael, and J.R. Ruzycki. 2004b. Lower Snake River Compensation Plan: Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Imnaha rivers for the 2001-02 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Portland.
- Flesher, M.W., G.R. Vonderohe, G.C. Grant, D.L. Eddy, and R.W. Carmichael. 2005a. Lower Snake River Compensation Plan: Oregon Summer Steelhead Evaluation Studies. Oregon Department of Fish and Wildlife, 2001 and 2002 Bi-Annual Progress Report, Salem.
- Flesher, M.W., R.W. Carmichael, and G.C. Grant. 2005b. Lower Snake River Compensation Plan: Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Imnaha rivers for the 2002-03 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Salem.
- Flesher, M.W., R.W. Carmichael, and G.C. Grant. 2007. Lower Snake River Compensation Plan: Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Imnaha rivers for the 2003-04 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Salem.
- Flesher, M.W., R.W. Carmichael, G.C. Grant, and L.R. Clarke. 2008a. Lower Snake River Compensation Plan: Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Imnaha rivers for the 2004-05 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Salem.
- Flesher, M.W., R.W. Carmichael, G.C. Grant, and L.R. Clarke. 2008b. Lower Snake River Compensation Plan: Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Imnaha rivers for the 2005-06 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Salem.
- Flesher, M.W., S.M. Warren, D.L. Eddy, L.R. Clarke, and R.W. Carmichael. 2009a. Lower Snake River Compensation Plan: Oregon Summer Steelhead Evaluation Studies. Oregon Department of Fish and Wildlife, 2005 Annual Progress Report, Salem.
- Flesher, M.W., R.W. Carmichael, and L.R. Clarke. 2009b. Lower Snake River Compensation Plan: Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Imnaha rivers for the 2006-07 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Salem.

- Flesher, M.W., R.W. Carmichael, and L.R. Clarke. 2010. Lower Snake River Compensation Plan: Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Innaha rivers for the 2007-08 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Salem.
- Flesher, M.W., R.W. Carmichael, and L.R. Clarke. 2011. Lower Snake River Compensation Plan: Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Innaha rivers for the 2008-09 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Salem.
- Flesher, M.W., R.W. Carmichael, and L.R. Clarke. 2012. Lower Snake River Compensation Plan: Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Innaha rivers for the 2009-10 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Salem.
- Flesher, M.W., S.M. Warren, R.W. Carmichael, and L.R. Clarke. 2013. Lower Snake River Compensation Plan: Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Innaha rivers for the 2010-11 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Salem.
- Flesher, M.W., S.M. Warren, R.W. Carmichael, and L.R. Clarke. 2014. Lower Snake River Compensation Plan: Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Innaha rivers for the 2011-12 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Salem.
- Flesher, M.W., R.W. Carmichael, and L.R. Clarke. 2015. Lower Snake River Compensation Plan: Summer steelhead creel surveys on the Grande Ronde, Wallowa, and Innaha rivers for the 2012-13 run year. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Salem.
- Gee, S.A., G. R. Vonderohe, M.W. Flesher, D.L. Eddy, G.C. Grant, and R.W. Carmichael. 2007. Lower Snake River Compensation Plan: Oregon Summer Steelhead Evaluation Studies. Oregon Department of Fish and Wildlife, 2003 Annual Progress Report, Salem.
- Gee, S.A., M.W. Flesher, D.L. Eddy, L. R. Clarke, J. R. Ruzycki, and R.W. Carmichael. 2008. Lower Snake River Compensation Plan: Oregon Summer Steelhead Evaluation Studies. Oregon Department of Fish and Wildlife, 2004 Annual Progress Report, Salem.
- Jonasson, B.C., R.W. Carmichael, and T.A. Whitesel. 1994. Residual hatchery steelhead: Characteristics and potential interactions with spring Chinook salmon in northeast Oregon. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Portland.

- Jonasson, B.C., R.W. Carmichael, and T.A. Whitesel. 1995. Residual hatchery steelhead: Characteristics and potential interactions with spring Chinook salmon in northeast Oregon. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Portland.
- Jonasson, B.C., R.W. Carmichael, and T.A. Whitesel. 1996. Residual hatchery steelhead: Characteristics and potential interactions with spring Chinook salmon in northeast Oregon. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Portland.
- Messmer, R.T., R.W. Carmichael, and M.W. Flesher. 1989. Lower Snake River Compensation Plan: Oregon evaluation studies. Oregon Department of Fish and Wildlife, Fish Research Project, AFF1-LSR-91-1. Annual Progress Report, Portland.
- Messmer, R.T., R.W. Carmichael, and M.W. Flesher. 1990. Lower Snake River Compensation Plan: Oregon evaluation studies. Oregon Department of Fish and Wildlife, Fish Research Project, AFF1-LSR-91-1. Annual Progress Report, Portland.
- Messmer, R.T., R.W. Carmichael, M.W. Flesher, and T.A. Whitesel. 1991. Evaluation of Lower Snake River Compensation Plan facilities in Oregon. Oregon Department of Fish and Wildlife, Fish Research Project, AFF1-LSR-92-10. Annual Progress Report, Portland.
- Messmer, R.T., R.W. Carmichael, M.W. Flesher, and T.A. Whitesel. 1992. Evaluation of Lower Snake River Compensation Plan facilities in Oregon. Oregon Department of Fish and Wildlife, Fish Research Project, AFF1-LSR-94-06. Annual Progress Report, Portland.
- Messmer, R.T., R.W. Carmichael, M.W. Flesher, and T.A. Whitesel. 1993. Evaluation of Lower Snake River Compensation Plan facilities in Oregon. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Portland.
- Ruzycki, J.R., M.W. Flesher, R.W. Carmichael, and D.L. Eddy. 2003. Lower Snake River Compensation Plan: Oregon evaluation studies – steelhead life history, genetics, and kelt reconditioning. Oregon Department of Fish and Wildlife, Fish Research and Development. 1997-2001 Progress Report, Portland.
- USACOE (U.S. Army Corps of Engineers). 1975. Special report to the Lower Snake River Compensation Plan: Lower Snake River, Washington and Idaho. U.S. Army Engineer District, Walla Walla, Washington.
- Warren, S.M., M.W. Flesher, D.L. Eddy, L.R. Clarke, and R.W. Carmichael. 2009. Lower Snake River Compensation Plan: Oregon Summer Steelhead Evaluation Studies. Oregon Department of Fish and Wildlife, 2006 Annual Progress Report, Salem.
- Warren, S.M., M.W. Flesher, D.L. Eddy, L.R. Clarke, and R.W. Carmichael. 2010. Lower Snake River Compensation Plan: Oregon Summer Steelhead Evaluation Studies. Oregon Department of Fish and Wildlife, 2007 Annual Progress Report, Salem.

- Warren, S.M., M.W. Flesher, D.L. Eddy, L.R. Clarke, and R.W. Carmichael. 2011a. Lower Snake River Compensation Plan: Oregon Summer Steelhead Evaluation Studies. Oregon Department of Fish and Wildlife, 2008 Annual Progress Report, Salem.
- Warren, S.M., M.W. Flesher, D.L. Eddy, L.R. Clarke, and R.W. Carmichael. 2011b. Lower Snake River Compensation Plan: Oregon Summer Steelhead Evaluation Studies. Oregon Department of Fish and Wildlife, 2009 Annual Progress Report, Salem.
- Warren, S.M., M.W. Flesher, D.L. Eddy, L.R. Clarke, and R.W. Carmichael. 2012. Lower Snake River Compensation Plan: Oregon Summer Steelhead Evaluation Studies. Oregon Department of Fish and Wildlife, 2010 Annual Progress Report, Salem.
- Warren, S.M., M.W. Flesher, D.L. Eddy, L.R. Clarke, and R.W. Carmichael. 2013. Lower Snake River Compensation Plan: Oregon Summer Steelhead Evaluation Studies. Oregon Department of Fish and Wildlife, 2011 Annual Progress Report, Salem.
- Whitesel, T.A., B.C. Jonasson and R.W. Carmichael. 1993. Residual hatchery steelhead: Characteristics and potential interactions with spring Chinook salmon in northeast Oregon. Oregon Department of Fish and Wildlife, Fish Research Project. Annual Progress Report, Portland.
- Zippen, C. 1958. The removal method of population estimation. *Journal of Wildlife Management* 22:82-90.

APPENDIX A: Corrections to Tables 7 and 8 in the 2014 Annual Progress Report with revised numbers shown in bold.

Table 7. Timing of adult steelhead returns to LSRCP facilities in 2014 by location and origin.

Period	Week of the year	Number of fish trapped ^a					
		Wallowa		Big Canyon		Little Sheep	
		Hatchery	Natural	Hatchery	Natural	Hatchery	Natural
Jan 22-28	4	-	-	-	-	-	-
Jan 29-Feb 04	5	-	-	-	-	-	-
Feb 05-11	6	0	0	-	-	-	-
Feb 12-18	7	0	0	-	-	-	-
Feb 19-25	8	0	0	0	0	0	0
Feb 26-Mar 04	9	0	0	111	0	26	1
Mar 05-11	10	55	0	333	9	50	1
Mar 12-18	11	450	1	197	6	89	1
Mar 19-25	12	329	0	212	2	13	0
Mar 26-Apr 01	13	221	0	146	2	78	10
Apr 02-Apr 08	14	247	0	131	2	58	6
Apr 09-15	15	268	3	208	7	139	15
Apr 16-22	16	157	3	64	4	67	11
Apr 23-29	17	76	1	64	3	45	6
Apr 30-May 06	18	40	0	31	3	166	57
May 07-13	19	20	0	25	7	12	6
May 14-20	20	14	0	8	3	23	8
May 21-27	21	1	0	0	0	2	2
May 28-June 03	22	0	0	-	-	0	0
Jun 04-10	23	-	-	-	-	-	-
Jun 11-17	24	-	-	-	-	-	-
June 18-24	25	-	-	-	-	-	-
Total		1,878	8	1,530	48	768	124

^aThe ladder was opened on 4 February at Wallowa Fish Hatchery, and weirs were installed 19 February at Big Canyon Facility (Deer Creek) and 21 February at Little Sheep Creek Facility. Adult collections ended 2 June at Wallowa Fish Hatchery, 26 May at Big Canyon Facility, and 2 June at Little Sheep Creek Facility.

Table 8. Number, disposition, and mean fork length (mm) of adult steelhead that returned to LSRCP facilities in 2014 by stock, origin, estimated age (freshwater:saltwater), and gender. M indicates male and F indicates female.

Facility, stock, disposition	Hatchery ^a									Natural ^b									Grand total		
	1:1		1:2		2:1		1:3		Total	2:1		2:2		2:3		3:1		3:2		Total	
	M	F	M	F	M	F	M	F		M	F	M	F	M	F	M	F	M			F
<i>Wallowa Hatchery (Wallowa Stock-Production)</i>																					
Trapped	750	357	87	155	4	11	0	3	1367	4	1	1	0	0	0	1	0	1	0	8	1375
Passed	0	0	0	0	0	0	0	0	0	4	1	1	0	0	0	1	0	1	0	8	8
Outplanted	8	1	0	1	0	0	0	0	10	0	0	0	0	0	0	0	0	0	0	0	10
Kept	742	356	87	154	4	11	0	3	1357	0	0	0	0	0	0	0	0	0	0	0	1357
Mortality	3	0	1	0	0	0	0	0	4	0	0	0	0	0	0	0	0	0	0	0	4
Spawned	86	51	31	66	0	0	0	0	234	0	0	0	0	0	0	0	0	0	0	0	234
Killed ^c	653	305	55	88	4	11	0	3	1119	0	0	0	0	0	0	0	0	0	0	0	1119
<i>Wallowa Hatchery (Wallowa Stock-Fallbrood)</i>																					
Trapped	280	162	24	43	0	1	0	1	511	0	0	0	0	0	0	0	0	0	0	0	511
Passed	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Outplanted	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Kept	280	162	24	43	0	1	0	1	511	0	0	0	0	0	0	0	0	0	0	0	511
Mortality	1	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1
Spawned	104	83	14	34	0	1	0	1	237	0	0	0	0	0	0	0	0	0	0	0	237
Killed ^c	175	79	10	9	0	0	0	0	273	0	0	0	0	0	0	0	0	0	0	0	273
<i>Wallowa Hatchery (Total Returns)</i>																					
Trapped	1030	519	111	198	4	12	0	4	1878	2	1	1	0	0	0	3	0	1	0	8	1886
Fork length (mm)	572	561	698	684	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Standard deviation	29	35	34	27	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Sample size	37	39	16	22	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Wallowa Hatchery (Fall Broodstock)</i>																					
Transferred to WFH	12	8	0	2	0	0	0	0	22	0	0	0	0	0	0	0	0	0	0	0	22
Passed	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Outplanted	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Kept	12	8	0	2	0	0	0	0	22	0	0	0	0	0	0	0	0	0	0	0	22
Mortality	2	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	2
Spawned	9	6	0	2	0	0	0	0	17	0	0	0	0	0	0	0	0	0	0	0	17
Killed	1	2	0	0	0	0	0	0	3	0	0	0	0	0	0	0	0	0	0	0	3

Table 8. Continued.

Facility, stock, Disposition	Hatchery ^a									Natural ^b									Grand Total		
	1:1		1:2		2:1		1:3		Total	2:1		2:2		2:3		3:1		3:2		Total	
M	F	M	F	M	F	M	F	M		F	M	F	M	F	M	F	M	F	M		F
<i>Big Canyon Facility (Wallowa stock)</i>																					
Trapped	644	427	126	317	3	7	0	6	1530	10	5	3	5	0	0	13	6	3	3	48	1578
Passed	0	0	0	0	0	0	0	0	0	10	5	3	5	0	0	13	6	3	3	48	48
Outplanted	75	40	51	25	3	7	0	4	205	0	0	0	0	0	0	0	0	0	0	0	205
Returned to river ^d	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Kept	569	387	75	292	0	0	0	2	1325	0	0	0	0	0	0	0	0	0	0	0	1325
Mortality	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Spawned	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Killed ^c	569	386	75	292	0	0	0	2	1325	0	0	0	0	0	0	0	0	0	0	0	1325
Fork length (mm)	-	-	-	-	-	-	-	-	-	565	-	-	-	-	-	625	-	-	-	-	-
Standard deviation	-	-	-	-	-	-	-	-	-	7	-	-	-	-	-	-	-	-	-	-	-
Sample size	-	-	-	-	-	-	-	-	-	2	-	-	-	-	-	1	-	-	-	-	-
<i>Little Sheep Creek Facility (Imnaha stock)</i>																					
Trapped	370	305	49	44	0	0	0	0	768	35	43	8	22	0	0	8	7	0	1	124	892
Passed	52	84	11	9	0	0	0	0	156	27	39	6	16	0	0	7	7	0	1	103	259
Outplanted	108	77	18	8	0	0	0	0	211	0	0	0	0	0	0	0	0	0	0	0	211
Kept	210	144	20	27	0	0	0	0	401	8	4	2	6	0	0	1	0	0	0	21	422
Mortality	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Spawned ^e	47	41	8	15	0	0	0	0	111	8	4	2	6	0	0	1	0	0	0	21	132
Killed ^f	163	103	12	12	0	0	0	0	290	0	0	0	0	0	0	0	0	0	0	0	290
Fork length (mm)	567	577	659	642	-	-	-	-	-	557	553	634	659	-	-	556	561	-	-	-	-
Standard deviation	21	50	40	24	-	-	-	-	-	43	18	2	36	-	-	19	60	-	-	-	-
Sample size	10	17	4	9	-	-	-	-	-	19	21	2	13	-	-	5	6	-	-	-	-

^a Wallowa stock hatchery ages apportioned using CWT data and 121 scale samples collected in 2014; Imnaha stock hatchery ages apportioned using CWT data, 115 scale samples from 2014. Mean fork lengths are from fish with scale samples collected in 2014.

^b Wallowa stock (Wallowa Hatchery and Big Canyon Facility) natural steelhead ages apportioned using historical data (220 samples) and 2014 data (3 samples); Little Sheep Creek Facility natural steelhead ages apportioned using 2013 data (40 samples) and 2014 data (57 samples). Mean fork lengths are from fish with scale samples collected in 2014.

^c For Wallowa stock steelhead, 920 fish that returned to Wallowa Hatchery and 1,074 fish that returned to Big Canyon Facility were euthanized and donated to local food banks. In addition, 46 fish from Wallowa Hatchery and 120 fish from Big Canyon Facility were euthanized and donated to local schools for educational purposes. For Imnaha stock steelhead, 161 fish that returned to Little Sheep Creek Facility were euthanized and donated to local food banks.

^d Hatchery steelhead are no longer recycled to the Wallowa River fishery for additional angler opportunity.

^e Includes 11 natural males that were live-spawned and passed above the weir.

^f Includes 17 hatchery males and 4 hatchery females that were initially outplanted to Big Sheep Creek. These 21 fish were subsequently recaptured